Abstract #012345
Parallel sessions: 1,2,3,4
Plenary sessions: 1,2,3

Elastic ρ' Electroproduction at High Q^2

H1 Collaboration

An analysis is reported of the enhancement in the $\pi^+\pi^-\pi^+\pi^-$ invariant mass ($M_{4\pi}$) spectrum of the elastic electroproduction reaction $ep \to \pi^+\pi^-\pi^+\pi^- p$ at $M_{4\pi} \sim 1.550$ GeV. When interpreted in terms of the production of a single resonance ρ', a measurement is made of the ratio of the cross sections $ep \to e\rho' p$ to $ep \to e\rho p$ using $2.2\,\mathrm{pb}^{-1}$ of e^+p interaction data for the kinematic ranges $4 < Q^2 < 50$ GeV^2 and $40 < W < 140$ GeV. Significant $\rho(770)$ production is observed in the $\rho' \to \pi^+\pi^-\pi^+\pi^-$ decay products. Kinematical dependencies of the ρ' production cross section are presented. The data are also analysed in terms of the production of the vector meson resonances $\rho'(1450)$ and $\rho'(1700)$, which have been shown to provide an interpretation of e^+e^- annihilation and and elastic photoproduction of 4π. The results are discussed in terms of the production of $^3S_1$ and $^3D_1$ quark-antiquark radial recurrences with $I^G J^{PC} = 1^+1^{--}$ and in terms of models of their production at high Q^2.