Abstract #000, March 19, 1999

# Measurement of Dijet Cross-Sections at Low $Q^2$ in $ep$ Collisions and the Extraction of an Effective Parton Density for the Virtual Photon

M. Erdmann[1] (for the H1 Collaboration)

*(1) University Karlsruhe / DESY, Notkestr. 85, D-22607 Hamburg, Germany*

The triple-differential dijet cross-section, $\mathrm{d}^3\sigma_{ep}/\mathrm{d}Q^2\mathrm{d}\overline{E_t}^2\mathrm{d}x_\gamma^{jets}$, is measured with the H1 detector at HERA as a function of the photon virtuality $Q^2$, the fraction of the photon's momentum carried by the parton entering the hard scattering, $x_\gamma^{jets}$, and the square of the mean transverse energy, $\overline{E_t}^2$, of the two highest $E_t$ jets. Jets are found using a longitudinal boost-invariant $k_T$ clustering algorithm in the $\gamma^* p$ center of mass frame. The measurements cover the ranges $1.6 < Q^2 < 80$ GeV$^2$ in virtuality and $0.1 < y < 0.7$ in inelasticity $y$. The results are well described by leading order QCD models which include the effects of a resolved component to the virtual photon. Models which treat the photon as point-like fail to describe the data. An effective leading order parton density for the virtual photon is extracted as a function of the photon virtuality, the probing scale and the parton momentum fraction. The $x_\gamma$ and probing scale dependences of the parton density show characteristic features of photon structure, and a suppression of this structure with increasing $Q^2$ is seen.

[1] H1 Collaboration, C.Adloff et al.,
DESY-98-205 / submitted to Eur. Phys. J. C