Paper pa02-060
Submitted to the 28th International Conference on High Energy Physics
ICHEP'96, Warsaw, Poland, July 1996

A Measurement of the Production of $D^{*\pm}$ Mesons in Deep-Inelastic Diffractive Interactions at HERA

H1 Collaboration

Abstract

The first measurement of the cross section for the production of $D^{*\pm}$ mesons in deep-inelastic diffractive interactions at the ep collider HERA is presented. The data were taken during the 1994 running period and correspond to an integrated luminosity of 2.2 pb^{-1}. Using the decay channel $D^{*+} \to D^0\pi^+_{slow} \to (K^-\pi^+)\pi^+_{slow}$, a cross section of $(380 \pm {}^{150}_{120\,stat} \pm {}^{140}_{110\,sys})$ pb is measured in the kinematic region $10\ \mathrm{GeV}^2 < \mathrm{Q}^2 < 100\ \mathrm{GeV}^2$, $0.06 < y < 0.6$, $x_{I\!P} < 0.05$, $M_Y < 1.6$ GeV, $|\eta| < 1.25$ and $p_\perp > 1$ GeV, where M_Y is the mass of the diffractively scattered proton remnant system and η and $p_\perp$ are respectively the pseudorapidity and transverse momentum of the $D^{*\pm}$ in the H1 laboratory frame. A predicted cross section is calculated using a partonic model of $e I\!P$ collisions in which the parton density functions for the $I\!P$ are taken from a QCD analysis of the diffractive structure function $F_2^{D(3)}$. In this model a "leading" gluon distribution carries about 80% of the momentum of the diffractive exchange. The result (~ 200 pb) is consistent with the measured value. A prediction (less than ~ 10 pb) based on a quark-dominated $I\!P$ is inconsistent at the 2σ level with the measured value.

Introduction

The contribution of charm production to diffractive interactions is sensitive to the partonic structure of the diffractive exchange. Measurements of the diffractive structure function $F_2^{D(3)}$ suggest that this structure receives a substantial contribution from gluons, and a QCD analysis of $F_2^{D(3)}$ employing the DGLAP evolution equations yields a solution which tends towards a "leading" gluon distribution at the starting scale for the perturbative evolution $Q^2 = 2.5\,\mathrm{GeV}^2$ [1, 2, 3, 4]. Such a solution gives rise to the expectation of a large cross section for charm production through boson-gluon fusion.

In this paper we present a search for the production of $D^{*\pm}$ mesons which subsequently decay via the channel[1] $D^{*+} \to D^0\pi^+_{slow} \to (K^-\pi^+)\pi^+_{slow}$. The first observation of $D^{*\pm}$ production in deep-inelastic diffractive interactions is reported and a measurement is made of the $D^{*\pm}$ production cross section in the kinematic region 10 $\mathrm{GeV}^2 < \mathrm{Q}^2 <$ 100 GeV^2, $0.06 < y < 0.6$, $x_{I\!P} < 0.05$, $M_Y < 1.6$ GeV, $|\eta| < 1.25$ and $p_\perp > 1$ GeV. M_Y is the mass of the diffractively scattered proton remnant system; η and $p_\perp$ are respectively the pseudorapidity and transverse momentum of the $D^{*\pm}$ in the H1 laboratory frame. The measured cross section is compared with two predictions of a partonic model of diffractive scattering based on electron-pomeron interactions. In one case, the parton distributions ascribed to the pomeron are obtained from the QCD analysis of $F_2^{D(3)}$. In the other case, the pomeron is given a QCD-evolved quark-based structure [3].

This cross section measurement follows a comparison reported in [3] of the $D^{*\pm}$ yield in deep-inelastic diffraction with that expected based on the QCD analysis of $F_2^{D(3)}$.

Event Selection

The data used were collected at the H1 detector [5] in 1994, using 27.5 GeV positrons colliding with 820 GeV protons. After rejecting data taken when parts of the detector needed for this analysis were not operational, the integrated luminosity used is 2.24 pb^{-1}.

This analysis is restricted to events with a scattered positron detected in the backward electromagnetic calorimeter (BEMC). There must be a localized energy deposit of at least 10 GeV within the angular range $155^\circ < \theta < 173^\circ$ with respect to the direction of the proton beam. The deposit must lie within the efficient region of the BEMC and be associated with a hit in the backward proportional chamber (BPC), which is immediately in front of the BEMC. These requirements restrict the accessible range in Q^2 and y. The requirement that a $D^{*\pm}$ be produced in the central region imposes an additional limit at the lower end of the y range. The region covered is 10 $\mathrm{GeV}^2 < \mathrm{Q}^2 <$ 100 GeV^2 and $0.06 < y < 0.6$. The kinematic variables Q^2 and y are calculated using the "sigma" method [6], which minimises the QED radiative corrections to the Born cross section.

Diffractive events are recognized by their lack of energy flow in the region around the proton beam. Energy flow in this region is detected via charged particles produced in secondary scattering in the beam pipe and associated material, as well as those coming directly from the ep interaction vertex [7]. An event is rejected if it has more than 7.5 GeV in the plug calorimeter, more than one hit pair in the first half of the forward

[1] Charge conjugate states are implicitly included.

muon detector, or any hits in the first three layers of the proton remnant tagger. The pseudorapidity of the most forward cluster with more than 400 MeV, η_{max}, must be less than 3.2. The fraction of the proton's momentum carried by the diffractive exchange, $x_{I\!P} = (Q^2 + M_X^2)/(Q^2 + W^2)$, is reconstructed using tracks and calorimeter information. Only events with $x_{I\!P}$ less than 0.05 are accepted.

The scattered proton is not detected, so it is not possible to distinguish an event in which the proton scatters elastically from one in which it dissociates into a low-mass state. The cross section is instead quoted for the range $M_Y < 1.6$ GeV. Models of proton dissociation [8, 9, 10] imply that the correction from the measured cross section ($M_Y <$ 1.6 GeV) to one in which the proton vertex is truly elastic is about 5%.

Reconstruction of $D^{*\pm}$

The charmed meson D^{*+} is reconstructed in this analysis using its decay $D^{*+} \to D^0 \pi^+_{slow} \to (K^-\pi^+)\pi^+_{slow}$. Because of the small mass difference $\Delta M = M_{D^{*+}} - M_{D^0}$, the pion from the D^{*+} decay (π^+_{slow}) has a momentum of only 40 MeV in the D^{*+} rest frame. Due to this tight kinematic constraint, ΔM is better resolved than the D^{*+} mass itself and is used in the cross-section measurement. The combinatorial background is also suppressed because of the small available phase space near threshold.

The decay products are detected in the central tracker. The analysis is restricted to the region with a good detection efficiency: tracks are only used if they lie within the angular range $25° < \theta < 155°$ and have a $p_\perp$ of at least 200 MeV (K,π) or 100 MeV (π_{slow}). To protect against badly reconstructed tracks, each track is required to be at least 10 cm long in the projection perpendicular to the beam and to start in the inner section of the tracker.

All accepted tracks with $p_\perp > 200$ MeV are combined in oppositely charged pairs. No particle identification is used; each track is allowed to be either a kaon or a pion. If a pair has an invariant mass within 80 MeV of the D^0 mass, it is counted as a D^0 candidate. Each D^0 candidate is combined in turn with all remaining tracks with $p_\perp > 100$ MeV and the appropriate charge to reconstruct D^{*+} candidates. After cuts on the direction ($|\eta| < 1.25$) and transverse momentum ($p_\perp > 1$ GeV) of the $D^{*\pm}$, the mass difference ΔM is plotted as a histogram (fig. 1).

The number of $D^{*\pm}$ candidates is determined by fitting the histogram with a Gaussian peak plus a background function $a(\Delta M - M_{\pi^+})^b$, where a, b and the three parameters of the Gaussian are allowed to vary. The number of $D^{*\pm}$ detected is found to be 11.6 ± 4.0 (statistical error only).

The distribution of η_{max} for $D^{*\pm}$ candidates in the region of the peak, selected as described above (apart from the cut on η_{max}), is shown in fig. 2. It clearly shows the low-η plateau characteristic of diffractive events.

Acceptance and Background Corrections

The acceptance is determined by running the H1 detector simulation program on a sample of $D^{*\pm}$ events from the diffractive Monte Carlo generator RAPGAP [11], which is based



Figure 1: Distribution of the mass difference $\Delta M = M(K^-\pi^+\pi^+_{slow}) - M(K^-\pi^+)$, with a fitted curve of the form $a(\Delta M - M_{\pi^+})^b + Gaussian$ and a prediction from RAPGAP with the pomeron structure extracted from $F_2^{D(3)}$.



Figure 2: Distribution of η_{max} for diffractive $D^{*\pm}$ candidates in the region $|\Delta M - 145.4 \text{ MeV}| < 2.0$ MeV, with the expectation from the non-diffractive generator AROMA.

on a partonic model of $e\mathbb{P}$ interactions. The efficiency of the BEMC trigger is known to be better than 99% and is neglected.

The main sources of background are reflections (contributions from other D^0 decay modes), DIS background (events with $x_{\mathbb{P}} > 0.05$), and proton dissociation events (those with $M_Y > 1.6$ GeV). The contribution from reflections is estimated using RAPGAP. The contribution from DIS background is estimated using RAPGAP (for $0.05 < x_{\mathbb{P}} < 0.2$) and the heavy quark generator AROMA [12] (for $x_{\mathbb{P}} > 0.2$). The fraction of events in which the proton dissociates into a system with $M_Y > 1.6$ GeV is estimated using the diffractive vector-meson dominance (VMD) generator, DIFFVM [8].

Systematic Errors

The largest systematic errors on the cross section are those due to the uncertainty in the $p_\perp$ distribution of the $D^{*\pm}$ ($\pm^{24}_{14}\%$) and the correction for reflections ($\pm 19\%$). The effect of varying the $p_\perp$ distribution is estimated by reweighting the Monte Carlo and recalculating the acceptance. There is also an error of $\pm 11\%$ from the fitting procedure, including uncertainty in the shape of the combinatorial background.

The other sources of systematic uncertainty taken into account, each responsible for an error of no more than $\pm 5\%$, are Monte Carlo statistics, the branching ratios for the D^{*+} and D^0 decays, the corrections for proton dissociation events and DIS background, the effect of noise in the forward detectors, and the uncertainty in the distributions of t, $x_{\mathbb{P}}$ and β.

Result

The cross section[2] is calculated for the following kinematic region: 10 GeV^2 < Q^2 < 100 GeV^2, $0.06 < y < 0.6$, $x_{\mathbb{P}} < 0.05$, $M_Y < 1.6$ GeV, $|\eta(D^*)| < 1.25$ and $p_\perp(D^*) > 1.0$ GeV (η and $p_\perp$ in the H1 laboratory frame). The result is

$$\sigma(ep \to eD^{*\pm}XY) = (380 \pm ^{150}_{120} \pm ^{140}_{110})\ \text{pb},$$

where the errors are the statistical and experimental systematic uncertainties respectively.

The Monte Carlo generator RAPGAP, which models diffractive events as the interaction of the positron with a partonic pomeron, predicts a value of ~ 200 pb when the parton densities of the pomeron are taken from a QCD analysis [3, 4] of the diffractive structure function $F_2^{D(3)}$. When the parton densities are instead calculated on the assumption of a quark-based pomeron at 2.5 GeV^2, evolved to higher Q^2 [3], the predicted cross section is less than ~ 10 pb.

[2]The cross section is defined as $N/\mathcal{L}_{int}$, where N is the corrected number of $D^{*\pm}$ produced and $\mathcal{L}_{int}$ is the integrated luminosity.

Conclusions

A first measurement of the cross section for $D^{*\pm}$ production in deep-inelastic diffractive interactions is reported. The result for the kinematic region covered is $(380 \pm {}^{150}_{120\,stat} \pm {}^{140}_{110\,sys})$ pb. The prediction ($\sim$ 200 pb) of a model in which charm is produced by boson-gluon fusion and the partonic structure of diffractive exchange is taken from a QCD analysis of diffractive interactions is consistent with the measured cross section. In this model, the structure of the pomeron receives a substantial contribution from a hard gluon distribution. The smaller prediction (less than $\sim$ 10 pb) of a model in which the diffractive exchange is a quark-based pomeron is inconsistent at the 2σ level with the measured cross section.

Acknowledgments

We are grateful to the HERA machine group, whose outstanding efforts have made and continue to make this experiment possible. We thank the engineers and technicians for their work in constructing and now maintaining the H1 detector, our funding agencies for financial support, the DESY technical staff for continual assistance, and the DESY directorate for the hospitality which they extend to the non-DESY members of the collaboration.

References

[1] J. P. Phillips: *Rapidity Gap Events at HERA and the Structure of the Pomeron*, in Proc. of the International Workshop on Deep–Inelastic Scattering and Related Topics, Paris, France, April 1995, ed. J-F. Laporte and Y. Sirois, p.359;

J. B. Dainton: *Deep–Inelastic Physics with H1 at the HERA ep Collider*, in Proc. of the International Workshop on Deep–Inelastic Scattering and Related Topics, Paris, France, April 1995, ed. J-F. Laporte and Y. Sirois, p.37.

[2] H1 Collaboration: *The Partonic Structure of Proton Diffraction*, paper EPS–0491 submitted to the 1995 Europhysics Conference, Brussels, and to the 1995 Lepton Photon Symposium, Beijing.

[3] A. Mehta: *Deep–Inelastic Diffraction*, in Proc. of the Topical Conference on Hard Diffractive Processes, February 18–23 1996, Eilat, Israel.

[4] H1 collaboration: *A Measurement and QCD Analysis of the Diffractive Structure Function* $F_2^{D(3)}$, paper pa02-061 contributed to this conference.

[5] H1 Collaboration: *The H1 Detector*, DESY 93–103 (1993).

[6] U. Bassler, G. Bernardi: *On the Kinematic Reconstruction of Deep Inelastic Scattering at HERA: the* Σ *Method*, DESY 94–231 (1994).

[7] H1 Collaboration: Phys. Lett. B 348 (1995) 681.

[8] B. List: *Diffraktive J/psi-Produktion in Elektron-Proton-Stössen am Speicherring HERA*, Diploma thesis H1–10/93–319 (1993).

[9] R. Engel: Proceedings of the XXIXth Rencontre de Moriond (1994) 321.

[10] T. Sjöstrand: Comp. Phys. Comm. 39 (1986) 347;
T. Sjöstrand, M. Bengtsson: Comp. Phys. Comm. 43 (1987) 367;
T. Sjöstrand, M. Bengtsson: Comp. Phys. Comm. 46 (1987) 43.

[11] H. Jung: Comp. Phys. Comm. 86 (1995) 147.

[12] G. Ingelman, J. Rathsman, G.A. Schuler: *AROMA 2.2 – A Monte Carlo Generator for Heavy Flavour Events in ep collisions*, DESY 96–058 (1996).