Measurement of the forward jet cross–section in the low–x_B region at HERA as a signature of BFKL dynamics

Ewelina Maria Łobodzińska

PhD Dissertation
Institute of Nuclear Physics

Kraków, July 1997

Preface

Thesis performed at the INP in Kraków
and at the DESY laboratory in Hamburg
under the supervision of Prof Albert De Roeck
and of Prof Jacek Turnau.

Contents

1 Introduction

The main aim of high-energy physics is the search for fundamental constituents of matter and the nature of interactions between them. Research in this field has been carried out with various kinds of experiments. Scattering experiments turned out to be very successful in unraveling the structure of matter. In these experiments an energetic point–like test particle is scattered on the probe and its angular and energy distributions are measured. In 1911, while bombarding a golden foil with α particles Rutherford observed occasional large–angle scatterings and from this fact he deduced that a compact nucleus lay at the centre of the atom.

More recently, giant particle accelerators have enabled us to probe the structure of nucleon very finely. In mid–1950s elastic scattering of electrons at Stanford showed that the proton is an extended object with dimension of the order of 1 fm. Inelastic scattering of 20 GeV electrons on nucleons in the late 1960s and early 1970s at Stanford established point–like quark constituents of the proton and revealed approximate scaling of the structure functions. In 1975, higher beam energies allowed observations of logarithmic violations of scaling, which was instrumental for the formulation of Quantum Chromodynamics.

In 1992 the electron–proton collider HERA was commissioned at the Deutsches Elektronen–Synchrotron laboratory in Hamburg. With this accelerator, c.m. energies of 300 GeV can be reached compared to 30 GeV in current fixed–target experiments. This results in a vast increase in phase space, physics potential for lepton–nucleon scattering, and resolution of proton constituents (Q^2 rises by a factor of 100).

The HERA experiments, H1 and ZEUS, are therefore exploring a new kinematical region, which was suggested to be sensitive to new dynamic features of QCD. There are several "dedicated" measurements which are aimed at revealing QCD in the region where values of Bjorken x are very low. One of them is the analysis of deep inelastic scattering events containing, in the hadronic final state, a jet – different from the jet resulting from the quark struck by the virtual photon – emitted close to the proton remnant direction. This measurement is designed in such a way that the traditional DGLAP [2] parton evolution is suppressed and the BFKL evolution [4]– expected to become valid in the low–x area – enhanced. Thus, this analysis has a chance to find a distinct signature of BFKL dynamics in the HERA region.

In this thesis we present detailed studies of the forward jet measurement as a possible footprint of the BFKL evolution. We use data collected in 1994 by the H1 experiment.

In Section 2 the deep inelastic scattering process is briefly described and basic kinematical variables are introduced. We show general principles of the quark parton model and explain how it gets improved by Quantum Chromodynamics. Different evolution schemes used to estimate behaviour of parton densities in a proton are also discussed. Section 3 is devoted to BFKL signatures which may be observed in the HERA experiments. In that section the motivation for the forward jet studies are presented. A short description of the Monte Carlo models used in our analysis is given in Section 4. Main information points on HERA and some details about the H1 detector components relevant for our measurement are presented in Section 5. In Section 6 data selection, sources of background and methods of its removal are explained. Jet studies are reported in Section 7. In Section 8 the data correction method used in this analysis, i.e. Bayes unfolding, is explained. We present some tests performed in order to check correctness of this method and comparison of results obtained with this method and with the simple bin-to-bin unfolding. Section 9 is devoted to the presentation of the final results of the forward jet analysis and to the discussion of these results versus predictions of various models. A short summary is given is Section 10.

All information contained in Sections 2 – 5 is based on literature, while the analysis presented in the next sections has been performed by the author.

2 Deep inelastic scattering

In the electron–proton (ep) deep inelastic scattering (DIS) process – schematically shown in Fig.2.1 – the incoming electron couples to the electroweak current J which probes the structure of the proton. The neutral (NC) and charged (CC) components of the current can be distinguished by the final state electron or neutrino, respectively. The virtuality of the exchanged boson Q^2 determines the resolution with which the proton is probed. For Q^2 much larger than proton mass squared (m_p), the proton can be thought of as a group of quasi–free constituents, partons, one of which interacts with the current while the rest (i.e. the proton remnant) moves on unperturbed.



Figure 2.1: *Schematic diagram showing deep inelastic ep scattering as described by the Quark Parton Model.*

Let us label the incoming and the outgoing lepton four–momenta with k and k', respectively, and the proton four–momentum with p. The exchanged boson has four–momentum $q = k - k'$, and the squared centre–of–mass energy of the ep system $s = (k+p)^2$.

For a fixed value of energy $\sqrt{s}$, the inclusive $ep \rightarrow eX$ scattering, in which only the scattered electron is measured, is fully described by any two of the following variables:

- $Q^2 \equiv -q^2 = -(k-k')^2 > 0$ – square of the four–momentum transfer,
- $\nu = q \cdot p/m_p$ – energy of the current J in the rest frame of the incoming proton,
- $y = p \cdot q/p \cdot k = \nu/\nu_{max}$ – fraction of the incident electron energy transferred to the proton in the rest frame of the proton.
- $x = Q^2/(2q \cdot p) = Q^2/(2m_p\nu)$ – the Bjorken scaling variable; in the parton model of a deep inelastic collision, it represents the fraction of the proton momentum carried by the struck quark,
- $W^2 = (p+q)^2 = m_p^2 + Q^2(1/x - 1)$ – squared invariant mass of the final hadronic system X.

Three of these variables are related[1] by:

$$xys = Q^2 \ , \tag{2.1}$$

Since both variables, x and y, are bounded by the kinematical condition $0 < x, y < 1$, the maximum value of the momentum transfer is $Q^2_{max} = s$.

Choosing x and Q^2 as independent variables, the differential cross–section for inelastic $ep \to eX$ scattering has the form:

$$\frac{d^2\sigma}{dxdQ^2} = \frac{4\pi\alpha^2}{xQ^4}[y^2 xF_1(x,Q^2) + (1-y)F_2(x,Q^2)], \tag{2.2}$$

where F_1 and F_2 are structure functions of the proton. The $1/Q^4$ dependence results from virtual photon exchange.

In the Bjorken limit, i.e. Q^2 and $2p \cdot q \to \infty$ while x is fixed, structure functions depend approximately only on dimensionless variable x:

$$F_i(x,Q^2) \approx F_i(x) \ . \tag{2.3}$$

This is the famous Bjorken scaling, first observed in the DIS experiments at SLAC, which finds natural explanation in the quark parton model.

2.1 The quark parton model

While describing deep inelastic electron–proton scattering in the quark parton model (QPM) [1] it is important to refer to a particular frame, called the infinite momentum frame. In this frame the proton is moving very fast so that the relativistic time dilation slows down the rate with which the partons interact with each other. That makes characteristic time of the interaction between the virtual photon and a parton $(1/Q)$ much smaller than the time of parton–parton interactions. Thus, one may view the photon as being absorbed *instantaneously* by some quark in the proton and – in other words – the struck quark appears essentially *free* during the short time of interaction with the virtual photon.

Therefore, the electron–proton scattering may be written as an incoherent sum of probabilities of scattering from single free quarks:

$$\frac{d^2\sigma}{dxdQ^2} = \sum_q \int_0^1 d(\xi) f_q(\xi) \left(\frac{d^2\sigma}{dxdQ^2}\right)_{eq} , \tag{2.4}$$

where $f_q(\xi)$ is the probability of finding parton q in the proton carrying fraction ξ of its momentum.

The scattered and unscattered partons recombine then to form hadrons. Hadronization takes place over a long time scale in comparison with the scattering process, so it can be treated separately.

[1] High energy limit $s \gg m_P^2$ is assumed.

In the QPM both proton and partons are assumed to be massless in the high energy limit. In this case, fraction ξ of the proton momentum carried by the struck quark is equal to the Bjorken variable x, since the massless condition for the scattered quark gives:

$$0 = (\xi p + q)^2 \qquad \Rightarrow \qquad \xi = \frac{-q^2}{2p \cdot q} = x \; . \tag{2.5}$$

The differential electron–parton cross–section in (2.4) depends on the assumptions about parton structure. Identifying the partons with quarks, spin–1/2 constituents of proton, the elementary cross–section is of the form:

$$\left(\frac{d^2\sigma}{dxdQ^2}\right)_{eq} = \frac{2\pi\alpha_{em}^2 e_q^2}{Q^4}\left[1 + (1-y)^2\right]\delta(x-\xi), \tag{2.6}$$

where α_{em} is the fine structure constant and e_q are fractional charges of quarks. Let us stress that in this case quarks are elementary (pointlike) particles.

Upon inserting equation (2.6) into (2.4) and comparing it with (2.2), one can obtain the following QPM prediction:

$$F_2(x) = 2xF_1(x) = x\sum_q e_q^2 \int_0^1 d\xi f_q(\xi)\delta(x-\xi) = \sum_q e_q^2 x f_q(x). \tag{2.7}$$

This expression displays three important results:

- Structure functions of the proton directly "measure" partonic distributions $f_q(x)$.
- Proton structure functions exhibit the Bjorken scaling (F_2 does not depend on Q^2). This is the result of pointlike structure of quarks.
- Structure functions are not independent: $F_2 = 2xF_1$. This is known as the Callan–Gross relation which follows from the spin–1/2 nature of the charge–carrying constituents of the proton.

The Bjorken scaling is observed for values of $x \sim 0.1$, and the structure functions increase and decrease logarithmically in Q^2 for smaller and larger values of x, respectively. This profound experimental fact, which cannot be accounted for by the QPM, is explained by Quantum Chromodynamics (QCD), the fundamental theory of strong interactions.

2.2 QCD–improved parton model

One of the most crucial features of QCD is dependence of the strong coupling constant α_s on Q^2:

$$\alpha_s(Q^2) = \frac{\alpha_s(\mu^2)}{1 + [\alpha_s(\mu^2)/6\pi](33/2 - n_f)log(Q^2/\mu^2)}, \tag{2.8}$$

where n_f is the number of flavours and μ^2 is an arbitrary normalization point at which the coupling α_s has been measured. In contrast to the QED coupling constant, the QCD coupling $\alpha_s(Q^2)$ decreases with increasing Q^2. This leads to the fact that QCD is asymptotically free, that is $\alpha_s \to 0$ as $Q^2 \to \infty$. This phenomenon, called asymptotic freedom, puts the QPM on a sound theoretical basis, since deep inelastic scattering is a high Q^2 process.

Perturbative QCD can be used to calculate the Q^2 dependence of the deep inelastic structure functions of the proton.

The QPM formula (2.7) may be regarded as the zeroth–order term in the QCD perturbative expansion of the proton structure function. In the first–order approximation one has to add contribution from the processes schematically shown in Fig.2.2(b). After summing the infinite number of such emissions, structure function F_2 takes the following form:

$$F_2(x, Q^2) = \sum_{i=q,\bar{q}} e_i^2 x f_i(x, Q^2) \tag{2.9}$$

where f_i are Q^2–dependent **quark** and **antiquark** distributions in proton.



Figure 2.2: *Schematic diagrams of partonic subprocesses contributing to deep inelastic scattering: a) term of the zeroth order in α_s – the QPM, b) terms of the first order in α_s.*

QCD emissions described above lead to the Altarelli–Parisi (DGLAP) evolution equations [2]:

$$\frac{\partial f_i(x, Q^2)}{\partial \log Q^2} = \frac{\alpha_s(Q^2)}{2\pi} \sum_{j=q,\bar{q},g} \int_x^1 \frac{d\xi}{\xi} f_j(\xi, Q^2) \; P_{ij}(\frac{x}{\xi}) \; , \tag{2.10}$$

which determine how quark, antiquark and **gluon** distributions evolve with Q^2 once they are known at some initial scale Q_0^2. Functions P_{ij} describe probability of finding parton i with momentum fraction x inside parton j with momentum fraction ξ. Parton distributions are universal, which means that they describe quark (or gluon) composition of the proton in any hard interaction it undergoes.

Perturbative QCD cannot provide initial conditions, i.e. parton distributions at some initial scale Q_0^2, for integration of the DGLAP evolution equations. These have to be taken from DIS experiments. The basic procedure is to parametrize the x dependence of initial densities $f_i(x, Q^2)$ at some low scale Q_0^2 (but high enough to justify the use of perturbative QCD), and then evolve the initial distributions up in Q^2 to compute the cross–sections or structure functions at measured (x, Q^2) points. The initial parameters are tuned to get the best agreement between the computed and the measured quantities.

In this way, the QCD parton model has introduced a definite pattern of scaling violations. Structure functions $F_i(x, Q^2)$ are no longer simply scaling functions of x alone, but have acquired a logarithmic dependence on Q^2.

2.3 Altarelli–Parisi evolution equations

As already mentioned in the previous section, the traditional way to estimate the behaviour of parton densities in a proton is based on the Altarelli–Parisi (also called DGLAP, i.e. Dokshitzer–Gribov–Lipatov–Altarelli–Parisi) evolution equations [2], which in the lowest order resum the leading powers of $\log Q^2$. That is, for each additional factor of α_s only the leading $\log Q^2$ terms accompanying this α_s are kept.

As it was noticed by Dokshitzer [3], in an axial gauge[2] these leading logarithms are generated by "ladder" diagrams of the type shown in Fig.2.3. The diagram with n gluon rungs corresponds to the $(\alpha_s \log Q^2)^n$ contribution. In the DGLAP scheme this parton cascade follows strong ordering in the transverse momenta:

$$Q^2 \gg k_{nT}^2 \gg k_{n-1T}^2 \gg \cdots \gg k_{1T}^2 \tag{2.11}$$

and ordering in the longitudinal momenta:

$$x_n < x_{n-1} < \cdots < x_1 < 1. \tag{2.12}$$



Figure 2.3: *"n–rung" ladder diagrams which are effectively resummed by the DGLAP and BFKL evolution equations.*

2.4 Evolution equations in the low–x region

When x is small, the partons are mainly gluons, so attention is focused on the evolution of gluon densities $g(x, Q^2)$ in a proton. In the low–x and large Q^2 region, behaviour

[2] An axial gauge is one in which the gluon has only two physical polarization states; in such a gauge we do not need unphysical "ghosts" contributions which in general would be required to cancel the scalar polarization component of the gluon.

of $g(x, Q^2)$ can be calculated in the double leading logarithmic approximation (DLLA) of the DGLAP equations. This involves summation of the double leading logarithmic contributions proportional to $(\alpha_s \log Q^2 \log 1/x)^n$. It amounts to resummation of the ladder diagrams where both the transverse and fractional longitudinal gluon momenta are strongly ordered along the ladder (Fig.2.3):

$$Q^2 \gg k_{nT}^2 \gg k_{n-1T}^2 \gg \cdots \gg k_{0T}^2 \tag{2.13}$$

$$x_n \ll x_{n-1} \ll \cdots \ll x_1 \tag{2.14}$$

The sum of these graphs gives, up to slowly varying logarithmic factors, the following formula:

$$xg(x, Q^2) \sim exp\left[2\left(\frac{3\alpha_s}{\pi}\log(Q^2)\log\left(\frac{1}{x}\right)\right)^{\frac{1}{2}}\right] \tag{2.15}$$

However, this approximation does not take into account all leading terms in the small–x limit. Terms containing the leading power of $\log(1/x)$, but not accompanied by the leading powers of $\log Q^2$, are neglected. So, if we are interested in small x but moderate Q^2 values, the BFKL (Balitsky-Fadin-Kuraev-Lipatov) equation [4] is more appropriate. It sums leading $\alpha_s \log 1/x$ terms and keeps the full Q^2 dependence. The strong ordering in transverse momenta required by the DGLAP evolution equations is now relaxed and the integration is carried over the full k_T phase–space. This equation is also identified with summation of the ladder diagrams shown in Fig.2.3, but now the gluon cascade follows strong ordering in fractional longitudinal momenta:

$$x_n \ll x_{n-1} \ll \cdots \ll x_1 \tag{2.16}$$

The sum of these diagrams leads to a low–x gluon density behaviour of the form :

$$xg(x, Q^2) \sim x^{-\alpha_p+1} \tag{2.17}$$

with $\alpha_p - 1 = (12\alpha_s)/\pi \log 2$, where α_p is the intercept of the trajectory of the so–called Lipatov Pomeron. In the above, effects of the running α_s and of slowly varying logarithmic factors are omitted. For a typical value of $\alpha_s \approx 0.2$, we have $\alpha_p - 1 \approx 0.5$, so the strong rise of $xg(x, Q^2)$ should be observed for $x \to 0$.

The three different limits of $g(x, Q^2)$ in the $(\log Q^2$, $\log 1/x)$ plane are shown in Fig.2.4:

- **the leading** $log(Q^2)$ **region**: $\log Q2/Q_0^2 \gg 1$, $\log 1/x \sim 1$, parton densities are described by the Altarelli–Parisi evolution equations,

- **the double leading log region**: $\log Q_2/Q_0^2 \gg 1$, $\log 1/x \gg 1$, parton densities are described by the Altarelli–Parisi evolution, where only terms proportional to $(\alpha_s \log(Q^2)\log(1/x))^n$ are kept,

- **the leading** $log(1/x)$ **region**: $\log Q^2/Q_0^2 \sim 1$, $log 1/x \gg 1$, parton densities are described by the BFKL evolution equation.



Figure 2.4: *Three limits of $g(x,Q^2)$ in the ($\log(Q^2)$, $\log(1/x)$) plane.*

3 Possible BFKL signatures in the low–x region at HERA

The $e-p$ collider HERA entered a new kinematical region, where values of the Bjorken variable x_B are very small (10^{-2} – 10^{-4}) and virtuality Q^2 is large enough to justify the use of perturbative QCD, so one can expect that the effects predicted by the BFKL evolution equation may occur.

The proton structure function F_2, measured by two HERA experiments, H1 and ZEUS, really exhibits a steep growth with decreasing x_B [5] (Fig.3.1).

In this figure, experimental data are compared with various theoretical predictions, in particular with the MRS(D0) curve based on traditional scenarios without the Lipatov Pomeron and computed when no HERA data were available, and also with the MRS(D-) curve calculated with the assumption that gluon density for low–x is proportional to $x^{-0.5}$. It turned out that real data agree with neither of these extreme approximations but lie inbetween.

This behaviour of F_2 caused a heated debate on whether the HERA data are still in a region where the QCD evolution of gluon densities can be described by the DGLAP evolution equations, or whether they extend into a new region where QCD dynamics is described by the BFKL evolution equation. The final conclusion seems to be that present F_2 measurements do not yet allow us to discriminate between these two evolution schemes, and therefore other less inclusive measurements are necessary.



Figure 3.1: F_2 *proton structure function as measured by the HERA experiments in 1994.*

One of the simplest observables suggested as a test of the BFKL versus DGLAP hypothesis is the flow of transverse energy (E_T) as a function of rapidity η in the hadronic centre–of–mass frame.[3]

BFKL calculations [6] predict an increase of mean transverse energy $\langle E_T \rangle$ with decreasing x_B, while DGLAP calculations predict just the opposite. In addition, due to transverse momentum ordering, the DGLAP evolution is expected to produce less E_T in the region between the current and the remnant jets for low–x_B events, as compared to the BFKL evolution [6].

Analysis of transverse energy flow based on the H1 data collected in 1993 has produced very promising results. The E_T flow in the CMS as a function of rapidity has been reasonably well described by CDM Monte Carlo model (in this model gluons in the ladder are not ordered in transverse momenta, so it is sometimes called the BFKL–like model) in all x_B and Q^2 bins, while the MEPS Monte Carlo model (DGLAP–based model)[4] has followed the data for large x_B and Q^2 values only.

However, it appears that the recently improved version of the DGLAP–based model[5] provides a good overall description of the measured transverse energy flow. Also the $\langle E_T \rangle$ distribution is now fairly well described by both Monte Carlo models.

Nevertheless, the model based on the DGLAP evolution scheme needs to produce 60-80% of its transverse energy during the hadronization phase so as to be able to follow the data. Hadronization allows it not only to increase the amount of E_T significantly, but even to compensate for the wrong x_B dependence seen at the parton level.

[3]Particle masses are neglected so that rapidity is equal to pseudorapidity $\eta = -\ln \tan \Theta/2$.

[4]CDM – Colour Dipole Model, MEPS – Matrix Elements + Parton Shower. Monte Carlo models will be described in detail in the next chapter.

[5]Soft colour interactions were added and the sea–quark (remnant) treatment was improved.

Since analysis of transverse energy flow also turned out to be inconclusive, more attention has been paid to other possible signatures of BFKL dynamics, such as:

- Measurement of charged particles p_T spectra [7]. This measurement is not so sensitive to hadronization effects, so hard tails of the p_T distributions, as seen in the data, are well described by the CDM Monte Carlo model, while the MEPS model predicts much softer spectra.
- Di–jet production. It was suggested [8] that weakening of the azimuthal (back-to-back) correlation between jets produced in DIS at HERA may be used to identify BFKL dynamics. From the experimental point of view, such analysis seems to be quite difficult; nevertheless, serious attempts to test this effect have been made.
- Forward jet production [9], [10, 11], [12].
- Azimuthal decorrelation of the forward jet [13].

The last two items will be covered here since detailed analysis of the forward jets as a potential footprint of BFKL dynamics constitues the principal topic of this thesis.

3.1 Forward jets as a signature of BFKL dynamics at HERA region

A proposal of a measurement which would be able to probe the low–x region was made by Mueller [9] and thoroughly examined by the Durham [10, 11] and DESY [12] groups. The idea is to study deep inelastic scattering events containing, in a hadronic final state, a jet characterized by fractional longitudinal momentum $x_{jet} = p_{z_{jet}}/p$ (where p is proton momentum and $p_{z_{jet}}$ is longitudinal jet momentum) and transverse momentum $k_{T_{jet}}$ (Fig.3.2). Such events will be referred to as (DIS +) forward jet events.



Figure 3.2: *Schematic diagram of deep inelastic scattering events containing, in a hadronic final state, a forward jet that arises from either a quark or a gluon.*

In order to focus on the BFKL type of evolution, the two basic conditions have to be met:

- $k^2_{T_{jet}} \sim Q^2$
- $x_{jet} \gg x_B$

The former ensures suppression of the DGLAP type of evolution, since it leaves no room for strong ordering in transverse momenta. The latter requirement maximizes phase–space for the BFKL evolution.

As seen in Fig.3.2 the jet under consideration – in order to fulfill the second condition – is emitted close to the proton remnant direction defined as the forward direction in the HERA geometry.

The diagram relevant to the computation of the cross–section for the process illustrated in Fig.3.2 is shown below (Fig.3.3).



Figure 3.3: *Diagram illustrating the terms used for computation of the cross–section for the DIS + jet production process. The figure is taken from [11].*

F denotes the photon–gluon subprocess which embodies all the BFKL emissions and formally satisfies the BFKL equation,

$$F(z, Q^2, k_T^2) = F_0(k_T^2) + \int_z^1 \frac{dz'}{z'} \int dk_T'^2 K(k_T, k_T') F(z', Q^2, k_T'^2), \quad z = x_B/x_{jet} \qquad (3.1)$$

which is symbolically shown in Fig.3.4.



Figure 3.4: *Leading log$(1/x)$ approximation for F. The figure is taken from [11].*

F_0 is just the quark box (and the crossed quark box), i.e. the contribution where no gluon is emitted in the ladder – **the Born approximation**. The remaining term effectively sums the gluon ladder contribution arising from both virtual and real gluon emissions.

Since the photon is probing the parton (which produces a jet) rather than the proton itself, and the jet is expected to have quite large x_{jet}, we only need to provide parton distributions in the region where they are well known. This way the problem of having to provide parton distributions at small x is avoided. In terms of the jet variables the differential deep inelastic structure function has the form:

$$\frac{\partial F_2}{\partial x_{jet} \partial k_T^2} = \frac{3\alpha_s(Q^2)}{\pi k_T^4} \sum_a f_a(x_{jet}, k_T^2) F(z, Q^2, k_T^2) \tag{3.2}$$

where f_a denotes parton distributions.

Equations 3.1 and 3.2 are used to evaluate the differential cross–section for production of deep–inelastic + forward jet events. Calculations are done in two ways:

1. Only for the box diagram ($F = F_0$), so we get the cross–section without the BFKL effects. This is the Born approximation for the forward jet process.

2. Taking into account gluon emissions. Assuming the running coupling constant, equation 3.1 may be solved numerically by evolving down in z, starting at the boundary condition $F = F_0$ at some value of z_0 where the BFKL effect is expected to be small.
   Cross–section obtained in this way exhibits a steep rise with decreasing z.

Calculations results constitute the main motivation for the forward jet analysis and will be compared with the DIS + forward jet cross–section measured in the data.

In the end, let us stress again the ingenuity of Mueller's idea. In the forward jet measurement, phase–space for the BFKL type of evolution is maximized by the condition $x_{jet} \gg x_B$, and at the same time the DGLAP parton emissions are suppressed by the requirement $k_T^2 \sim Q^2$. Thus, this measurement has a chance to provide a distinct footprint of BFKL dynamics.

3.2 Azimuthal dependence of forward jets

Analysis of azimuthal dependence of forward jets was proposed [13] as a further signal of BFKL dynamics. In HERA reference frame, BFKL dynamics predicts that with increasing rapidity difference between the forward and the current jets, the forward jet "forgets" about the azimuthal direction defined by the outgoing electron, so the cross–section becomes $\Delta\phi$–independent. $\Delta\phi$ is defined in the HERA frame as the angle between transverse momenta of the outgoing lepton and the forward jet.

Azimuthal dependence of the forward jet production cross–section was calculated in [13] assuming the BFKL gluon evolution in the ladder. The result may be expressed by the following formula:

$$\frac{d\sigma}{d\Delta\phi} \sim A + C\cos(2\Delta\phi). \tag{3.3}$$

Coefficient C is negative, so at $\Delta\phi = \pi/2$ there is a maximum. However, with increasing x_{jet}/x_B the angular dependent part is suppressed while A increases. This confirms our expectations of the BFKL prediction to be $\Delta\phi$–independent for low–x_B.

Fig.3.5 shows prediction of azimuthal dependence for the Born and BFKL cases (plot on the left-hand-side). The Born calculations give only approximate predictions, so the results of the fixed–order matrix element numerical studies for various x_B ranges are also presented (the right-hand-side plot). For numerical calculations, the PROJET [14] Monte Carlo is used with the matrix element $g \to qq'g$, integrated over the full range of invariant mass squared $s_{q\bar{q}}$, where any of the partons may initiate the forward jet.



Figure 3.5: *Dependence on the difference in azimuthal angles of the outgoing electron and the forward jet, normalized to a common average. The left-hand figure: solid line – the BFKL result, dashed line – the approximative analytical Born prediction;* $0.2 \cdot 10^{-3} < x_B < 0.4 \cdot 10^{-3}$ *. The right-hand figure: fixed-order results,* $0.2 \cdot 10^{-3} < x_B < 0.4 \cdot 10^{-3}$ *– solid line,* $1.8 \cdot 10^{-3} < x_B < 2. \cdot 10^{-3}$ *– dotted line,* $3.4 \cdot 10^{-3} < x_B < 3.6 \cdot 10^{-4}$ *– dashed–dotted line.*

One may see that for the approximate analytical Born result the cross–section has a clear maximum at $\Delta\phi = \pi/2$. For the fixed–order results this cosine–like structure is deformed, as the maximum moves towards $\Delta\phi = \pi$. However, the biggest discrepancies between the numerical and Born predictions are observed for high values of x_B. In the BFKL case, any $\Delta\phi$–dependence is completely washed out.

In the analysis of the DIS + forward jet events we will also concentrate on the correlation in the azimuthal angle between the forward jet and the outgoing electron. Predictions of this section will be used for interpretation of our measurement results.

4 DIS Monte Carlo models

For comparison of the measurements with QCD predictions, models which describe the QCD processes at the parton level and the fragmentation into hadrons are indispensable.

In the H1 collaboration the Monte Carlo production is performed in three steps:

1. Event generation
   Various models generating ep scattering events are available. They differ mainly in the details of the parton cascade simulation and the phenomenological description of hadronization. Monte Carlo models considered in the analysis of the forward jet production are shortly described in the next section.

2. Detailed simulation of the H1 detector response
   Final state particles produced by Monte Carlo generators are traced through the H1 detector by means of the H1 simulation program – H1SIM. H1SIM works within the GEANT [21] frame and simulates responses of each part of the H1 detector.

3. Reconstruction of the simulated events
   Simulated events are reconstructed in the same way as the real data. For this purpose, the H1REC package is used.

In the following we describe MC programs used in this analysis.

LEPTO

In LEPTO [15] QCD processes are calculated up to $O(\alpha_s)$ according to exact first order matrix elements (ME). It contains the following processes:

- the leading–order parton–level process, $\gamma q \rightarrow q$ — shown by the diagram of the naive quark–parton model (QPM) in Fig.4.1 (a),
- gluon radiation (QCD–Compton), $\gamma q \rightarrow qg$ – Fig.4.1 (b),
- boson–gluon fusion (BGF), $\gamma g \rightarrow qq$ – Fig.4.1 (c).

Higher order contributions are simulated in the leading logarithmic approximation by the parton shower (PS).

In DIS the quark struck by the electroweak boson can emit partons either before or after the boson vertex giving rise to initial and final parton showers, respectively.
A parton in the incoming nucleon and close to the mass–shell can initiate a parton emission cascade. In each branching of the cascade, the parton becomes more off–shell with a space–like virtuality ($m^2 > 0$) due to radiation of on–shell or having a time–like virtuality ($m^2 < 0$) parton.
This initial state space–like shower results in a space–like quark which interacts with the electroweak boson that turns it into an outgoing quark which is either on–shell or has a time–like virtuality. In the latter case a final state time–like shower will result. Off–shell mass is reduced by branching into daughter partons with decreasing off–shell masses and decreasing opening angles. This shower continues until all partons are (essentially) on–shell. Any parton with a time–like virtuality from the initial state shower will develop similarly. General behaviours of initial and final state parton showers are similar since



Figure 4.1: *Basic parton processes implemented in LEPTO:*
a) diagram of naive quark–parton model,
b) diagrams contributing to the QCD–Compton process,
c) diagrams contributing to the BGF process.

they are both based on branching processes: $q \to qg$, $g \to q\bar{q}$, $g \to gg$ as described by the Altarelli–Parisi evolution equations in the leading logarithmic approximation of the QCD.
Combination of the two approaches (ME+PS) gives the first order parton emission plus the higher–order emissions through parton shower.
Fragmentation into hadrons is performed with the Lund string model as implemented in JETSET [20].

ARIADNE – CDM
ARIADNE [18] is a generator for QCD cascades only. Gluon radiation is performed in Ariadne by implementation of the Colour Dipole Model (CDM). In contrast to the bremsstrahlung–like parton shower model, the CDM model does not distinguish between initial and final state radiation and includes interference effects between them. The struck parton and the proton remnant form a colour dipole. When this dipole radiates a gluon, it splits into two radiating dipoles: one between the struck quark and the emitted gluon, and another one between the emitted gluon and the remnant. Further gluon radiations lead to a chain of such dipoles.

The CDM model, in contrast to LEPTO and HERWIG which are based on DGLAP evolution equation, is called BFKL–like model because gluons emitted in the cascade do not obey strong ordering in k_T.

DJANGO
The event generator DJANGO [16] simulates deep inelastic lepton–proton scattering. It includes both QED and QCD radiative effects and allows for generation of events ac-

cording to QCD cascade models or QCD matrix elements. These are matrix elements available which describe effects from perturbative QCD to first order in α_s due to virtual gluon emission and boson–gluon fusion. A combination of LLA QCD cascades with first–order matrix elements is possible. QCD cascades can be generated using either the parton shower (PS) model implemented in LEPTO, or the colour dipole model as implemented in ARIADNE [18].
Process $ep \rightarrow e'X$, including electroweak corrections of $O(\alpha)$ and the bremsstrahlung process $ep \rightarrow e'\gamma X$, is simulated with the help of HERACLES [17]. The full hadronic final state is generated using the Lund string model JETSET [20].

JETSET
The JETSET [20] Monte Carlo program is a generic package for jet fragmentation, particle decays, final–state parton showers, event–analysis routines, and other utilities.
Many Monte Carlo programs that generate parton configurations in different processes use JETSET for fragmentation. The fragmentation in JETSET is based on a string fragmentation model.
In the string model a string is stretched between colour–connected partons, so as to make colour singlet strings of minimum invariant mass. The string has a uniform energy per unit length, corresponding to a linear quark confining potential. Gluons are supposed to produce kinks on the string, each initially carrying localized energy and momentum equal to that of its parent gluon. The string breaks up into hadron–sized pieces through spontaneous $q\bar{q}$ pair production in its intense colour field. The string break–up process proceeds until only on–mass–shell hadrons remain.
A large fraction of the particles produced by fragmentation are unstable and subsequently decay into stable ones. The decay treatment is handled by JETSET, making use of a set of tables where branching ratios and decay modes are stored.

5 Experimental apparatus

5.1 HERA collider

Traditionally, lepton–nucleon interactions were investigated in fixed–target experiments where a stationary target was hit by a beam of high energy leptons. Colliding–beam accelerators allowed the energy accessible in the c.m. frame to be significantly increased as compared to fixed–target experiments. The electron–proton collider differs from electron–electron or proton–proton colliders mainly by the fact that it needs two independent magnet rings and accelerating systems. Electrons and protons have to be stored in two separate rings due to a big mass difference. Since electrons have small masses, energy loss by synchrotron radiation is very large ($\sim 1/\mathrm{m}_e^4$), and thus the electron beam needs much more power pumped in so as to compensate for the losses, which in consequence limits the practical energy achievable. Due to their bigger masses, protons may be accelerated to higher energies, but then stronger magnetic field is needed to keep them in the orbit. This leads to the fact that, in contrast to the electron ring, the proton one has to be equipped with superconducting magnets.

Hadron–Electron Ring Accelerator (HERA) is the first and only electron–proton collider. It was commissioned at the Deutsches Elektronen–Synchrotron (DESY) laboratory in Hamburg, Germany in 1992. HERA consists of two independent accelerators designed to store 820 GeV protons and 30 GeV electrons and to collide them head–on at four interaction points spaced uniformly around its 6.3 km – long underground tunnel. At present two multipurpose detectors are running at two of the interaction points – the H1 detector situated in the North Hall and the ZEUS detector in the South Hall.

At the beginning of 1994 HERA operated with $\mathrm{e}^-\mathrm{p}$ collisions, later the electron beam was replaced with a positron one[6]. The positron beam shows much better stability and is easier to guide than the electron one. Lifetime of the electron ring fills is almost 2 times higher for the positron than for the electron beams.

Integrated luminosity for physics analysis was 0.4 pb^{-1} for e^-p and 2.7 pb^{-1} for e^+p . The electron and proton beams were organized into 168 and 170 bunches, respectively. 153 bunches were colliding, separated from each other by 96 ns.

A summary of parameters designed and achieved in 1994 at the HERA collider is shown in Table 5.1.

5.2 H1 detector

The H1 detector is a nearly hermetic multi–purpose apparatus built to investigate inelastic high–energy interactions of electrons and protons at HERA. The H1 coordinate system is defined in such a way that its origin is placed at the nominal interaction point. Initial protons move in the positive z direction, the x coordinate points towards the centre of the HERA ring, and the y coordinate points upwards. Polar angle Θ is defined with respect to the proton beam direction.

The H1 detector is asymmetric. Asymmetry is required by the HERA event topology because the centre of mass of the collision is strongly boosted along the proton beam

[6] In this thesis the incident and scattered lepton will always be referred to as an electron.

HERA parameters	designed		running in 1994	
	electron	proton	electron	proton
Nominal Energy (GeV)	30	820	27.6	820
C.M. Energy (GeV)	314		301	
Maximum Q^2 (GeV^2)	9×10^4		9×10^4	
Luminosity (cm^{-2}s^{-1})	1.5×10^{31}		0.5×10^{31}	
Circumference (m)	6336			
Injection Energy (GeV)	14	40	14	40
Circulating Current (mA)	58	163	5 – 30	10 – 50
Number of Bunches	210		153	
Time between Crossings (ns)	96		96	
Horizontal Beam Size σ_x (mm)	0.26	0.29	0.27	0.18
Vertical Beam Size σ_y (mm)	0.07	0.07	0.06	0.06
Longitudinal Beam Size σ_z (cm)	0.8	11	0.8	11

Table 5.1: *Design and 1994–running values of the main HERA parameters.*

direction, and consequently high densities of energetic particles are expected in the forward region.

All *ep* collisions occur in the interaction region, the longitudinal dimension of which is determined by the length of the proton bunch and was found in 1994 to be $\sigma_{zvtx} = 11\ cm$.

General layout of the detector is shown in Fig.5.1. The interaction region is surrounded with tracking devices $\boxed{2,3}$ which measure momenta of charged particles and reconstruct the position of the interaction point. The tracking system consists of drift chambers interleaved with proportional chambers which enable fast trigger selection. All detectors are enclosed in calorimeter $\boxed{4,5}$ which permits to measure the energy and direction of charged and neutral particles. The liquid argon calorimeter covers central and forward regions. The very forward direction around the beam pipe is closed by plug calorimeter $\boxed{13}$ constructed of copper absorber and silicon pad readout. The calorimetric coverage in the backward region is completed with electromagnetic calorimeter $\boxed{12}$ located after proportional chamber (BPC)[7] and before two scintillator walls – time–of–flight detectors (TOF) – used to veto proton–induced background events. Superconducting coil $\boxed{6}$ surrounding the calorimeter produces a magnetic field of 1.2 T. The effect of the magnetic field on the beams is compensated by small solenoid $\boxed{7}$. Magnet yoke $\boxed{10}$ is made of iron stabs interleaved with streamer tube detectors for measuring hadronic energy escaping the main calorimeters. Muons are identified and measured by muon chambers $\boxed{9}$ inside and outside the iron. In the forward region muons are traced using forward muon spectrometer $\boxed{11}$ which consists of drift chamber planes mounted on either side of a toroidal magnet.

A detailed description of the H1 detector and its performance can be found in [22].

[7]The detectors in the backward region were upgraded in 1995 by replacement of the lead/scintillator tile calorimeter and a proportional chamber.