INAUGURAL-DISSERTATION
zur
Erlangung der Doktorwürde
der
Naturwissenschaftlich-Mathematischen
Gesamtfakultät
der
Ruprecht-Karls-Universität
Heidelberg

vorgelegt von

Dipl.-Phys. Thomas Nunnemann
aus Mainz

Tag der mündlichen Prüfung: 11.2.1999

Measurement of Leading Neutron Production in Deep–Inelastic *ep* Scattering at HERA

Gutachter: Prof. Dr. Bogdan Povh
Prof. Dr. Jürgen von Krogh

Bestimmung der Produktion vorwärts gestreuter Neutronen in tiefinelastischer ep-Streuung bei HERA

Beim H1 Experiment am HERA Beschleuniger wurden Neutronen, die unter einem kleinen Winkel ($0\,\mathrm{mrad} \leq \theta \lesssim 0.6\,\mathrm{mrad}$) in tiefinelastischer Streuung produziert wurden, nachgewiesen. Dazu wurde ein Kleinwinkel-Kalorimeter benutzt, welches 1996 eingebaut wurde. Vorgestellt wird die Messung des semi-inklusiven Wirkungsquerschnittes $\mathrm{d}\sigma(ep \to enX)/(\mathrm{d}x\mathrm{d}Q^2\mathrm{d}E')$ für die Produktion von Neutronen mit einem Transversalimpuls $p_T \leq 200\,\mathrm{MeV}$ im kinematischen Bereich von $2\,\mathrm{GeV}^2 \leq Q^2 \leq 50\,\mathrm{GeV}^2$, $6 \cdot 10^{-5} \leq x \leq 6 \cdot 10^{-3}$ und $160\,\mathrm{GeV} \leq E' \leq E_p$, wobei E' die Energie des Neutrons und $E_p = 820\,\mathrm{GeV}$ die des Protonstrahles bezeichnen. Verschiedene Produktionsmechanismen werden dem Ergebnis gegenübergestellt und ihre Vorhersagen überprüft. Die Neutronen-Daten werden mit einer ähnlichen Messung der Produktion von führenden Protonen verglichen. Für Energiebruchteile $0.7 \leq E'/E_p \leq 0.9$ stimmen beide semi-inklusive Wirkungsquerschnitte mit einem Regge-Modell überein, welches die Produktion von führenden Baryonen durch Pion-, Pomeron- und Reggeonaustausch beschreibt. Die Messung des Wirkungsquerschnittes für die Produktion von Neutronen ermöglicht die erste Abschätzung der Pion-Strukturfunktion bei kleinen Werten der Bjorkenvariable x.

Measurement of Leading Neutron Production in Deep–Inelastic ep Scattering at HERA

Deep–inelastic scattering events with a leading neutron have been observed by the H1 experiment at HERA using a forward neutron calorimeter. The calorimeter has been installed in 1996 and covers scattering angles between $0\,\mathrm{mrad}$ and $\lesssim 0.6\,\mathrm{mrad}$. For the first time at HERA, the semi–inclusive cross section $\mathrm{d}\sigma(ep \to enX)/(\mathrm{d}x\mathrm{d}Q^2\mathrm{d}E')$ for the production of leading neutrons with transverse momenta $p_T \leq 200\,\mathrm{MeV}$ has been measured in the kinematic region $2\,\mathrm{GeV}^2 \leq Q^2 \leq 50\,\mathrm{GeV}^2$, $6 \cdot 10^{-5} \leq x \leq 6 \cdot 10^{-3}$ and neutron energy $160\,\mathrm{GeV} \leq E' \leq E_p$, where $E_p = 820\,\mathrm{GeV}$ is the proton beam energy. The data are used to test different production models. The neutron data are compared to a similar measurement of leading proton production. For fractional energies $0.7 \leq E'/E_p \leq 0.9$, both semi–inclusive cross sections can be described by a Regge model of leading baryon production which consists of pion, pomeron and reggeon exchanges. The leading neutron data are used to estimate for the first time the structure function of the pion at small Bjorken–x.

Animadverti jam ante aliquot annos quàm multa, ineunte aetate, falsa pro veris admiserim, & quàm dubia sint quaecunque istis postea superextruxi, ac proinde funditus omnia semel in vitâ esse evertenda [...]

Several years have now elapsed since I first became aware that I had accepted, even from my youth, many false opinions for true, and that consequently what I afterward based on such principles was highly doubtful; and from that time I was convinced of the necessity of undertaking once in my life to rid myself of all the opinions I had adopted [...]

René Descartes, *Meditationes De Prima Philosophia*, 1641, Meditatio I.1,
English Translation by John Veitch, 1901.

Contents

Introduction

Revealing the structure of matter is one of the key issues in modern physics. The construction of systems, e. g. galaxies, solid states or elementary particles, gives information about the underlying physical forces and laws. Whereas in the macroscopic world the structure can be often seen directly with the naked eye, in the microscopic world of subatomic and subnuclear particles, most of the information is obtained by means of scattering experiments with highly energetic beams.

At the HERA accelerator, the structure of the proton is probed using high–energy beams of electrons and protons. The process of deep–inelastic scattering of the electron off the proton allows to study the partonic structure of the nucleon. The proton, as we understand it nowadays, is composed of quarks, antiquarks and gluons, which are assumed to be point–like objects. After many experiments, which started in the late 1960's, the structure of the proton can be parameterized in terms of parton distribution functions, which give the probability of finding an (anti–)quark or gluon with a certain momentum fraction inside the proton. Are these partons homogeneously and independently distributed over the full size of the proton or are there correlations between the different type of components? Hadronic correlations in the proton in form of virtual pions are believed to play a very important dynamical role in the nucleon structure, e. g. within the time allowed by the uncertainty principle, the proton might fluctuate into a state consisting of a pion and a neutron. Consequentially the question arises, is it possible to probe the structure of the pion inside the proton? Numerous theoretical works have suggested, that the production of highly energetic, forward neutrons in deep–inelastic *ep* scattering is sensitive to the structure of the pion. The produced neutron tags the virtual pion, which is probed by the exchanged photon.

In order to study neutron production, a *Forward Neutron Calorimeter* was installed in the H1 experiment at the *ep* collider HERA in the beginning of 1996. The calorimeter is located inside the HERA tunnel, approximately 100 m away from the central part of the H1 detector. This position allows to measure forward neutrons which are produced with very small scattering angles.

This work will focus on the measurement of the semi–inclusive production cross section of leading neutrons in deep–inelastic *ep* scattering using data obtained with the H1 detector in 1996. It is organized as follows: Chapter 1 gives an overview over the theory of semi–inclusive deep–inelastic scattering and discusses the pion exchange mechanism and alternative approaches for the production of leading neutrons. The HERA collider and the H1 detector are briefly described in Chapter 2. The construction of the forward neutron calorimeter is covered in Chapter 3, where special focus is put on the acceptance, calibration and resolution of the detector. The cross section measurement is presented in Chapter 4: The event selection and reconstruction, in particular the determination of the neutron's energy spectrum, are discussed in detail. In the last Chapter the result of this cross section measurement is presented and compared to the predictions of Monte Carlo simulations, which are based on different production mechanisms.

The neutron data are compared to a similar analysis of leading proton production. For high energies of the leading neutron or proton, both data sets can be described by a Regge Model of leading baryon production, which consists of pion, reggeon and pomeron exchanges. The leading neutron data are used to estimate for the first time the structure function of the pion at small Bjorken–x.

Chapter 1

Deep–Inelastic Scattering with a Leading Baryon

High energy scattering experiments of leptons on protons have a long and successful tradition in resolving the internal structure of the proton. Starting in the 1960's the measurement of elastic electron–proton scattering was used to determine the electromagnetic form–factor of the proton, which is directly related to the charge distribution and the magnetic moment of the proton.

In the late sixties, deep–inelastic scattering events, where the scattered electron looses a large amount of energy, were observed at the *Stanford Linear Accelerator Center* (SLAC). The experiments done at SLAC revealed, that the structure function of the proton only depends on the scaling variable x, and shows nearly no dependence on the four–momentum transfer Q^2 between the electron and the proton target [Blo69, Bre69]. This famous scaling behaviour had been predicted by Bjorken already in 1967 [Bjo67]. The *Quark Parton Model* (QPM), which was formulated by Feynman, gave an intuitive explanation for the observation of scaling [Fey69]. The scaling variable x was interpreted as the fractional momentum of the struck, point–like partons which were identified with the spin 1/2 quarks. Further experimental and theoretical progress showed that the proton does not consist only of quarks (valence and sea quarks) but that approximately half of the proton's momentum is carried by gluons, which in the framework of *Quantum Chromodynamics* (QCD) are identified as the carrier of the strong interaction. Due to the fact that the strong force between two partons is growing with increasing distance (the so called *confinement*), long range correlations between the quarks and gluons inside the proton are prominent and can lead to the formation of colourless hadronic fluctuations inside the proton.

Scattering processes on a long range scale, e. g. extended objects, are typically characterized by little transfer of transverse momentum. In *Regge* theory the scattering off a colourless hadronic fluctuation is described by the exchange of this colour–singlet object leaving the remainder of the target in a non–excited state, mostly a proton or a neutron.

The H1 and ZEUS experiment at the *ep* collider HERA discovered *large rapidity gap* events [ZEU93, H1C94], in which no particles are produced in the forward direction except a low mass state, presumably a high–energy proton, which escapes through the beam pipe undetected in the central detector. These events are usually interpreted



Figure 1.1: Schematic representation of inclusive deep–inelastic scattering in the QPM. The electron (e) interacts via an exchanged photon (γ^*) with a parton inside the proton (p). In the infinite momentum frame, xp can be identified with the initial momentum of the struck quark.

in terms of *pomeron* exchange [H1C95a, H1C97b, ZEU95, ZEU98]. The pomeron is a virtual particle which carries the quantum number of the vacuum and does not exist as a free particle. Also mesonic fluctuations inside the proton contribute to the total observed cross section. The most prominent contribution is the pion, since it is the lightest hadron.

Having these fluctuations, the deep–inelastic scattering process can take place on a parton of the virtual pion inside the proton rather than on a parton, which belongs to the "bare" proton. The pion might be probed by the exchanged photon offering the possibility of measuring the pion structure function in deep–inelastic *ep* scattering at HERA.

1.1 Inclusive Deep–Inelastic Scattering

Nowadays inclusive deep–inelastic scattering (DIS) is extensively studied at the HERA *ep* collider. In the Quark Parton Model, this reaction can be understood as the elastic scattering of the electron with a quark belonging to the proton. The distributions of the partons inside the proton can be determined by measuring only the energy and the scattering angle of the outgoing electron. Therefore this kind of measurement is called inclusive. At HERA one has the possibility of detecting in addition the hadronic final state, providing a more accurate determination of the kinematic variables.

1.1.1 Kinematics of Inclusive DIS

The deep–inelastic scattering reaction is sketched in Figure 1.1. Neglecting the contribution of the weak interaction, the scattering of the electron off the proton is, to lowest order, mediated by the exchange of a virtual photon. The struck parton hadronizes into the *current jet*, whereas the rest of the target proton constitutes the *proton remnant*.

The kinematics of this process can be naturally described in the four–dimensional *Minkowski* space. The following notation is of common use[1]:

k, k'	four–momentum of the incoming and outgoing electron,
E_e, E'_e	energy of the electron before and after the scattering process[2],
p	four–momentum of the proton in the initial state,
E_p	energy of the incoming proton[2],
m_e, m_p	masses of the electron and proton respectively,
P	sum of the four–momenta of the hadronic final state,
$q = k - k'$	four–momentum of the virtual photon,
θ_e	scattering angle of the electron[2,3],
$s = (k+p)^2$	squared centre–of–mass energy of the proton–electron system,
$W^2 = (q+p)^2$	squared centre–of–mass energy of the photon–proton system.

At fixed centre–of–mass energy $\sqrt{s}$, the inclusive event kinematics are completely defined by two of the following three Lorentz invariant variables:

$$\begin{aligned} Q^2 &= -q^2 = -(k-k')^2 \ , \\ x &= \frac{Q^2}{2p\cdot q} \ , \\ y &= \frac{p\cdot q}{p\cdot k} \ , \end{aligned} \tag{1.1}$$

which are related by $Q^2 = sxy$, if mass terms are neglected. In the HERA laboratory frame, the invariant variables can be expressed in terms of the beam energies and the kinematic quantities of the scattered electron:

$$\begin{aligned} s &= 4E_eE_p \ , \\ Q^2 &= 4E_eE'_e\cos^2\frac{\theta_e}{2} \ , \\ y &= 1 - \frac{E'_e}{E_e}\sin^2\frac{\theta_e}{2} \ . \end{aligned} \tag{1.2}$$

The equations are exact only to lowest order, e. g. in the absence of initial or final state radiation (see Section 4.7).

The quantities Q^2, x and y can be intuitively illustrated. *Heisenberg's* uncertainty principle relates Q^2, the virtuality of the exchanged photon, to the minimal distance, which can be resolved by the photon: $\hbar/\sqrt{Q^2}$. Thus, with increasing Q^2 smaller distances within the proton can be probed. In a frame where the momentum of the proton is much larger than its mass (*infinite momentum frame*), the scaling variable x can be interpreted as the fraction of the proton momentum carried by the struck quark. In the proton rest frame, $y = (E_e - E'_e)/E_e$, which can be illustrated as the part of the energy, which is transferred by the photon. Therefore, y is called "inelasticity".

[1] In this work, a system of natural units is used, whereby $\hbar = c = 1$.
[2] Quantities defined in the laboratory frame.
[3] In the HERA convention, scattering angles are defined with respect to the proton beam direction.

1.1.2 The Structure of the Proton

For not too high Q^2 ($Q^2 \ll m^2_{Z^0}$), the contribution of the weak interaction to the deep–inelastic scattering process can be neglected. In this region of Q^2 the double–differential cross section can be parameterized by two structure functions F_1 and F_2 [4]:

$$\begin{aligned} \frac{\mathrm{d}^2\sigma^{ep}}{\mathrm{d}x\,\mathrm{d}Q^2} &= \frac{4\pi\alpha^2}{x\,Q^4}\left[xy^2F_1(x,Q^2)+(1-y)F_2(x,Q^2)\right] \\ &= \frac{4\pi\alpha^2}{x\,Q^4}\left[1-y+\frac{y^2}{2[1+R(x,Q^2)]}\right]F_2(x,Q^2)\ , \end{aligned} \tag{1.3}$$

where α is the electromagnetic coupling constant and $R = (F_2 - 2xF_1)/(2xF_1)$ is the ratio between the absorption cross sections for longitudinally and transversely polarized virtual photons. The expectation of the QPM is, that R vanishes for spin 1/2 partons (*Callan–Gross* relation [Cal69]). QCD predicts deviations from zero, which is in accordance with recent measurements [Tao96, NMC97, Yu98].

In the Quark Parton Model the structure function F_2 is expected to be independent of Q^2 and can be related to the parton momentum distributions $f_i(x)$:

$$F_2(x) = \sum_{i=q_j,\bar{q}_j} e_i^2 x f_i(x)\ , \tag{1.4}$$

where the sum is over the quarks and antiquarks, and e_i is their electric charge.

1.1.3 QCD Evolution and the Violation of Scaling

In the QPM the normalization of the parton distribution functions $f_i(x)$ is constrained by the momentum sum rule:

$$\sum_i \int_0^1 \mathrm{d}x\, x f_i(x) = 1\ . \tag{1.5}$$

Measurements made in the seventies demonstrated that this sum rule is not fulfilled: Quarks and antiquarks can only account for approximately half the proton momentum. The experiments also showed that the scaling behaviour is only true for $x \approx 0.2$. Outside this x region F_2 depends on Q^2: the scaling hypothesis is violated. These deviations from the predictions of the QPM are explained by Quantum Chromodynamics.

QCD is the field theory of the strong interaction. Quarks, which carry a colour charge, interact by the exchange of a gluon. Since also the gluons have a colour charge, they interact with each other as well as with quarks. The self–coupling of the gluon field creates an anti–screening effect leading to a logarithmic decrease of the strong coupling constant $\alpha_s(Q^2)$ with increasing Q^2. Since $\alpha_s(Q^2) \to 0$ for $Q^2 \to \infty$, the quarks appear to be free at very large Q^2. This property is called *asymptotic freedom.* For low values of Q^2, $\alpha_s(Q^2)$ is large: The partons are confined in hadrons.

QCD gives an intuitive interpretation of the violation of the scaling behaviour. As the virtuality Q^2, with which the proton is probed by the photon, increases, finer

[4] A derivation of this formula can be found in numerous textbooks, e. g. [Rob90, Nac90, Per87, Pov95]



(a) (b) (c)

Figure 1.2: Vertices relevant to parton splitting functions: (a) Quark–gluon vertex determining P_{qq} and P_{gq}; (b) Annihilation vertex which induces P_{qg}; (c) Three gluon vertex determining P_{gg}.

structures within the proton are resolved. A quark, which is seen at a certain $Q^2 = Q_0^2$, might have radiated a gluon, when it is probed at a higher $Q^2 > Q_0^2$. The quark has lost momentum. With increasing Q^2 more and more splitting processes occur. Thus, as Q^2 increases, the quark density at low x grows and the probability to find a quark at large x decreases.

This physical picture can be quantitatively expressed by perturbative Quantum Chromodynamics (pQCD): For sufficiently small values of the strong coupling constant, solutions to QCD may be expanded in a perturbation series of powers of $\alpha_s(Q^2)$.

In pQCD the (anti–)quark ($q_i(x)$, $\bar{q}_i(x)$) and gluon ($g(x)$) distributions evolve as a function of Q^2. The dynamics of this evolution is described by the DGLAP equations (Dokshitzer, Gribov, Lipatov, Altarelli, Parisi) [Gri72, Dok77, Alt77], which are given in leading order (LO) by

$$\frac{\mathrm{d}q_i(x,Q^2)}{\mathrm{d}\ln Q^2} = \frac{\alpha_s(Q^2)}{2\pi}\int_x^1 \frac{\mathrm{d}y}{y}\left[\sum_j q_j(y,Q^2)P_{q_iq_j}\left(\frac{x}{y}\right) + g(y,Q^2)P_{q_ig}\left(\frac{x}{y}\right)\right] \quad (1.6)$$

$$\frac{\mathrm{d}g(x,Q^2)}{\mathrm{d}\ln Q^2} = \frac{\alpha_s(Q^2)}{2\pi}\int_x^1 \frac{\mathrm{d}y}{y}\left[\sum_j q_j(y,Q^2)P_{gq_j}\left(\frac{x}{y}\right) + g(y,Q^2)P_{gg}\left(\frac{x}{y}\right)\right] .$$

The splitting functions $P_{ba}(x/y)$ denote the probability for a parton a with fractional proton momentum y radiating a parton b, which has a fractional momentum $x < y$. The relevant LO vertices, which determine the splitting functions, are shown in Figure 1.2.

The DGLAP formalism permits, parton distributions, which are assumed to be known for all x at a certain Q_0^2, to be predicted for all x at larger $Q^2 > Q_0^2$. However, the equations cannot predict the x behaviour of the structure function.

In leading order[5] pQCD the structure function F_2 is like in the case of the QPM closely connected to the quark and antiquark distribution functions:

$$F_2(x,Q^2) = \sum_i e_i^2 x\left[q_i(x,Q^2) + \bar{q}_i(x,Q^2)\right] , \quad (1.7)$$

[5] In the DIS regularization scheme, Eq. 1.7 is valid to all orders.



Figure 1.3: Measurement of the structure function $F_2(x, Q^2)$ as a function of Q^2 at different values of x. The figure combines data from H1 [H1C96b, H1C98a], NMC [NMC95], BCDMS [Ben89] and SLAC [Whi92].

where the sum is over the different quark flavours. In next–to–leading order (NLO) in the commonly used $\overline{\mathrm{MS}}$ regularization scheme[6], F_2 is also directly related to the gluon density.

The high centre–of–mass energy at HERA allows one to measure the proton structure function in a wide kinematic region not explored by the preceding fixed–target experiments. Q^2 ranges from $\sim 1\,\mathrm{GeV}^2$ to $\sim 10,000\,\mathrm{GeV}^2$, and x values of approximately 10^{-5} can be reached at small Q^2. Figure 1.3 shows the measurement of $F_2(x, Q^2)$ made by the H1 collaboration together with the data from previous fixed–target experiments.

[6]The modified minimal subtraction scheme, e. g. see [Rob90].

1.2 Semi–Inclusive DIS with a Leading Baryon

In a semi–inclusive measurement, the properties of an additional final state particle besides the scattered electron are determined. In part of the DIS events a high–energy baryon (most likely a proton or neutron) is produced with a very small scattering angle. These very forward baryons are subsequentially referred to as leading particles.

The kinematics of the semi–inclusive process $ep \to eNX$, where N denotes a leading nucleon, can be intuitively described in the particle–exchange picture. The proton exchanges a virtual particle which interacts with the electron, whereas the rest of the proton composing the leading baryon acts as "spectator" (see Figure 1.4). The definition of the kinematics is however general and does not depend on the underlying physical model.



Figure 1.4: Production of a leading proton or neutron in the particle–exchange picture. A pion, reggeon or pomeron is exchanged.

The four–momentum, energy, mass and transverse momentum of the leading baryon is denoted by p', E', m_N and p_T respectively. Besides the two inclusive kinematic variables (e. g. x, Q^2) the kinematics of the semi–inclusive process are described by two additional variables[7]. A Lorentz invariant choice of the kinematic variables is

$$\begin{aligned} z &= 1 - x_{\mathbb{P}} = 1 - \frac{q \cdot (p - p')}{q \cdot p} \simeq \frac{E'}{E_p} \,, \qquad (1.8) \\ t &= (p - p')^2 \simeq -\frac{p_T^2}{z} - (1 - z)\left(\frac{m_N^2}{z} - m_p^2\right) \,. \end{aligned}$$

For $p_T \ll \sqrt{p'^2}$, which is in general true for tiny scattering angles, z is equivalent to the fractional energy of the leading baryon. $x_{\mathbb{P}}$ can be identified with the fractional

[7] The angle between the electron and baryon scattering plane is not considered here.

momentum of the exchanged object. t, which is always a negative quantity, corresponds to the four–momentum transferred between the proton and the final state baryon. In the particle–exchange picture t can be identified with the squared four–momentum of the exchanged particle. Another useful variable is

$$\beta = \frac{Q^2}{2q \cdot (p - p')} = \frac{x}{1-z} , \tag{1.9}$$

which can be interpreted as the Bjorken variable of the exchanged particle.

The mass of the central hadronic system X is related to the variable z, since

$$M_X^2 = (q + p - p')^2 = (1-z)(W^2 + Q^2 - m_p^2) + t - Q^2 \simeq (1-z)W^2 - zQ^2 . \tag{1.10}$$

In analogy to the inclusive proton structure function F_2, the four–fold differential cross section for baryon production can be parameterized by a semi–inclusive structure function, $F_2^{LB(4)}$, defined by

$$\frac{\mathrm{d}^4\sigma(ep \to eNX)}{\mathrm{d}x\,\mathrm{d}Q^2\,\mathrm{d}z\,\mathrm{d}t} = \frac{4\pi\alpha^2}{x\,Q^4}\left[1 - y + \frac{y^2}{2[1 + R(x, Q^2, z, t)]}\right] F_2^{LB(4)}(x, Q^2, z, t) . \tag{1.11}$$

Since $\beta = x/(1-z)$ this equation can be rewritten with respect to β:

$$\frac{\mathrm{d}^4\sigma(ep \to eNX)}{\mathrm{d}\beta\,\mathrm{d}Q^2\,\mathrm{d}z\,\mathrm{d}t} = \frac{4\pi\alpha^2}{\beta\,Q^4}\left[1 - y + \frac{y^2}{2[1 + R(\beta, Q^2, z, t)]}\right] F_2^{LB(4)}(\beta, Q^2, z, t) . \tag{1.12}$$

The structure function $F_2^{LB(4)}$ is denoted by $F_2^{LP(4)}$ and $F_2^{LN(4)}$ for the semi–inclusive processes which have final state protons and neutrons respectively.

1.3 Hadron–Hadron Interactions

In hadronic interactions the appropriate QCD scale is given by the momentum transferred. For large momentum transfer, which is referred to as hard interaction, the strong coupling constant is small, which allows one to apply pQCD. However, the bulk of the hadronic interactions take place at low momentum transfer, where a perturbative calculation is not possible due to large $\alpha_s(Q^2)$. In this non–perturbative regime, one has to rely on phenomenological approaches. Regge theory provides a powerful apparatus being able to describe soft hadronic cross sections.

1.3.1 Regge Theory in a Nutshell

Since the postulation of the pion by Yukawa in 1935 [Yuk35], the interaction between hadrons, in particular nucleons, has been attempted to be described by meson exchange models. In the One–Pion–Exchange (OPE) model (see [Dre61]) the scattering amplitude corresponds to the exchange of a particle with fixed angular momentum. The Regge theory of potential scattering [Reg59] provides a further development of this concept. The angular momentum is treated as a continuous complex variable and the scattering amplitude is associated with the exchange of a Regge trajectory of composite particles. This formalism permits to avoid divergences which are present in the OPE model. Detailed and comprehensive reviews of Regge theory can be found in, for example, [Col84, Kai79, Alb81, Gou83]. Here only some relevant results are briefly summarized.



(a) (b)

Figure 1.5: (a) Diagrammatic representation of a generalized s and t channel process. (b) Reggeon trajectory. Shown is the spin of the resonances as a function of their squared mass, which is identified as t (from [Lan94]).

The Regge Trajectory

The Regge trajectory is represented by a linear function in t:

$$\alpha(t) = \alpha_0 + \alpha' \cdot t \ , \tag{1.13}$$

where α_0 and α' are referred to as intercept and slope (shrinkage) parameter. One distinguishes two regions of t. For $t < 0$ (s channel process), the trajectory is exchanged between the two scattering hadrons. In the t channel process ($t > 0$), t can be related to the squared mass of the channel resonances, which occur at integer (meson trajectory) or half integer (baryon trajectory) values of $\alpha(t)$, which correspond to the spin of the resonances.

A single trajectory is characterized by a definite set of quantum numbers (isospin I, parity P, charge–conjugation C). In elastic scattering the ρ, ω, a_2 and f_2 trajectories are degenerate, so that they can be combined in a single *reggeon* trajectory as demonstrated in Figure 1.5b. The most important trajectories are:

- pomeron ($\mathbb{P}$), $\alpha_{\mathbb{P}}(t) \simeq 1.08 + 0.25\,\mathrm{GeV}^2 \cdot t$, $I^{PC} = 0^{++}$;
- reggeon ($\mathbb{R}$), $\alpha_{\mathbb{R}}(t) \simeq 0.5 + 0.9\,\mathrm{GeV}^2 \cdot t$;
- pion (π), $\alpha_\pi(t) \simeq 0 + 1\,\mathrm{GeV}^2 \cdot t$, $I^{PC} = 0^{-+}$ (π^0).

The pomeron trajectory has the quantum numbers of the vacuum. Its intercept has been determined on the basis of total cross section measurements for pp and $p\bar{p}$ scattering [Don92]. The slope parameter results from an analysis of the elastic scattering data [Don86].

Elastic Cross Sections of Hadronic Interactions

In the asymptotic limit of $s \to \infty$ and $t/s \to 0$, the contribution of a Regge trajectory $\alpha_k(t)$ to the amplitude $A_k^{ij}(s,t)$ of the elastic scattering process $ij \to ij$ is

$$A_k^{ij}(s,t) \sim \hat{\beta}(t) \left(\frac{s}{s_0}\right)^{\alpha_k(t)} , \tag{1.14}$$

where $\hat{\beta}(t)$ is the function specifying the coupling of the trajectory to the external particles, and $s_0 \approx 1\,\mathrm{GeV}^2$ is the hadron mass scale. The differential elastic cross section can then be expressed as

$$\frac{\mathrm{d}\sigma_{el}^{ij}}{\mathrm{d}t} \sim \frac{1}{s^2} \left| A_k^{ij}(s,t) \right|^2 \sim F(t) \left(\frac{s}{s_0}\right)^{2[\alpha_k(t)-1]} . \tag{1.15}$$

Generalizing the findings to the case of the exchange of several trajectories k, the differential cross section for the elastic reaction $ij \to ij$ is given by[8]

$$\frac{\mathrm{d}\sigma_{el}^{ij}}{\mathrm{d}t} = \sum_k \frac{\beta_{ik}^2(t)\beta_{jk}^2(t)}{16\pi} \left(\frac{s}{s_0}\right)^{2[\alpha_k(t)-1]} , \tag{1.16}$$

where $\beta_{ik}(t)$, $\beta_{jk}(t)$ are form–factors parameterizing the coupling of the trajectory $\alpha_k(t)$ to the particles i and j respectively.

In the region $|t| \lesssim 1\,\mathrm{GeV}^2$ the form–factors can be well approximated by an exponential:

$$\beta_{ik}(t) \simeq \beta_{ik}(0) e^{B_{ik}t} . \tag{1.17}$$

For large centre–of–mass energies $\sqrt{s}$ the differential cross section is dominated by pomeron exchange. Due to the linearity of the trajectory (see Equation 1.13) the cross section can be written as

$$\frac{\mathrm{d}\sigma_{el}^{ij}}{\mathrm{d}t} \propto \left(\frac{s}{s_0}\right)^{2[\alpha_{\mathbb{P}}(0)-1]} e^{b(s)t} , \tag{1.18}$$

where

$$b(s) = b_0 + 2\alpha'_{\mathbb{P}} \ln(s/s_0) . \tag{1.19}$$

With increasing s the forward peak of the cross section ($t \to 0$) becomes sharper, since the slope b is growing. This is commonly called *shrinkage*.

Total Cross Sections of Hadronic Interactions

The total cross section is closely related to the forward elastic scattering amplitude via the *Optical Theorem* (see Figure 1.6):

$$\sigma_T^{ij} \sim \frac{1}{s} \mathrm{Im} A_k^{ij}(s, t=0) \sim s^{\alpha_k(0)-1} . \tag{1.20}$$

[8] The notation of [Gou83] is used.



Figure 1.6: Regge diagrams for the total cross section, for elastic scattering and for single dissociation. The sketch for the total cross section σ_T^{ij} illustrates the optical theorem.

The generalization to the case of several exchanged trajectories yields

$$\sigma_T^{ij} = \sum_k \beta_{ik}(0)\beta_{jk}(0)s^{\alpha_k(0)-1} \ . \tag{1.21}$$

Donnachie and Landshoff showed that the total cross section of any hadronic reaction (including photoproduction $\gamma p \to X$) can be fitted to the form [Don92]

$$\sigma_T^{ij}(s) = X_{\mathbb{P}} s^{\alpha_{\mathbb{P}}(0)-1} + Y_{\mathbb{R}} s^{\alpha_{\mathbb{R}}(0)-1} \ , \tag{1.22}$$

with the normalizations $X_{\mathbb{P}}$ and $Y_{\mathbb{R}}$ for the pomeron and reggeon contribution respectively. The quantity of $X_{\mathbb{P}}$ is the same for reactions, where one particle has been replaced by its antiparticle, e. g. pp and $\bar{p}p$, leading to a convergence of both cross sections at high centre–of–mass energies. This is in agreement with a prediction from quantum field theory, known as the Pomerančuk theorem [Pom58], that the cross sections should become the same for particle and antiparticle at infinite high energies.

Cross Section of Single Dissociation

The cross section of the process $ij \to Xj$, in which the particle i dissociates into the system X with invariant mass M_X, is given by (compare Figure 1.6) [Gou83]

$$\frac{\mathrm{d}^2\sigma^{ij}}{\mathrm{d}t\,\mathrm{d}M_X^2} = \sum_{k,l} \frac{\beta_{ik}(0)\beta_{jl}^2(t)g_{kll}(t)}{16\pi s^2} \left(\frac{s}{M_X^2}\right)^{2\alpha_l(t)} \left(M_X^2\right)^{\alpha_k(0)} \ , \tag{1.23}$$

where g_{kll} denotes the couplings between the trajectories. In the case of diffractive dissociation, where $M_X^2 \ll s$, the pomeron dominates the cross section, so that

$$\begin{aligned}\frac{\mathrm{d}^2\sigma_D^{ij}}{\mathrm{d}t\,\mathrm{d}M_X^2} &= \frac{\beta_{i\mathbb{P}}(0)\beta_{j\mathbb{P}}^2(t)g_{\mathbb{P}\mathbb{P}\mathbb{P}}(t)}{16\pi s^2}\left(\frac{s}{M_X^2}\right)^{2\alpha_{\mathbb{P}}(t)}\left(M_X^2\right)^{\alpha_{\mathbb{P}}(0)} \\ &\propto s^{2[\alpha_{\mathbb{P}}(0)-1]}\left(M_X^2\right)^{-\alpha_{\mathbb{P}}(0)}\exp\left[b_D(s,M_X^2)t\right] ,\end{aligned} \tag{1.24}$$

where the last proportionality is valid in the limit of small $|t|$, where the couplings can be approximated by an exponential. The slope parameter b_D is

$$b_D(s,M_X^2) = b_{D,0} + 2\alpha'_{\mathbb{P}}\ln\left(\frac{s}{M_X^2}\right) , \tag{1.25}$$

giving a logarithmic shrinkage in s/M_X^2. The t dependence of the cross section parameterization in Equation 1.24 was experimentally verified in various reactions [Coo81].

1.4 Regge Phenomenology in DIS

If factorization holds, the semi–inclusive DIS reaction $ep \to eNX$ can be split into a soft and hard part (compare [Ing85]). The non–perturbative dynamics at the proton vertex is described by a factor $f_{\mathcal{P}/p}(z,t)$, which parameterizes the flux of the particle[9] $\mathcal{P}$ in the proton. The hard deep–inelastic interaction occurs between the electron and the exchanged particle. The four–fold differential cross section for nucleon production can therefore be expressed by

$$\begin{aligned}\frac{\mathrm{d}^4\sigma(ep\to eNX)}{\mathrm{d}x\,\mathrm{d}Q^2\,\mathrm{d}z\,\mathrm{d}t} &= \sum_{\mathcal{P}} f_{\mathcal{P}/p}(z,t)\frac{\mathrm{d}\sigma^{e\mathcal{P}}}{\mathrm{d}x\,\mathrm{d}Q^2} \\ &= \sum_{\mathcal{P}} f_{\mathcal{P}/p}(z,t)\frac{4\pi\alpha^2}{x\,Q^4}\left[1-y+\frac{y^2}{2(1+R)}\right]F_2^{\mathcal{P}}(\beta,Q^2) ,\end{aligned} \tag{1.26}$$

where $F_2^{\mathcal{P}}$ is the structure function of the exchanged particle. Comparing this expression with Equation 1.12, one obtains for the semi–inclusive structure function

$$F_2^{LB(4)}(\beta,Q^2,z,t) = \sum_{\mathcal{P}} f_{\mathcal{P}/p}(z,t)F_2^{\mathcal{P}}(\beta,Q^2) . \tag{1.27}$$

The flux factors $f_{\mathcal{P}/p}$ can be parameterized by means of Regge phenomenology. Consider the reaction $Ap \to XN$. Since the fractional energy z of the leading nucleon N is related to M_X^2 by $z \simeq 1 - M_X^2/s$ (compare Equation 1.10)[10], Equation 1.23 can be rewritten with respect to z and t:

$$\begin{aligned}\frac{\mathrm{d}^2\sigma^{Ap\to XN}}{\mathrm{d}t\,\mathrm{d}z} &\simeq -s\frac{\mathrm{d}^2\sigma^{Ap\to XN}}{\mathrm{d}t\,\mathrm{d}M_X^2} \\ &\simeq -\sum_{k,\mathcal{P}}\frac{\beta_{Ak}(0)\beta_{p\mathcal{P}}^2(t)g_{k\mathcal{P}\mathcal{P}}(t)}{16\pi}(1-z)^{1-2\alpha_{\mathcal{P}}(t)}\left[(1-z)s\right]^{\alpha_k(0)-1} .\end{aligned} \tag{1.28}$$

[9] The trajectories are identified with the corresponding particles.
[10] s is the squ. centre–of–mass energy of the Ap system, which in DIS ($A=\gamma^*$) corresponds to W^2.

This expression can be splitted into flux factors $f_{\mathcal{P}/p}$ and the Ak cross sections σ^{Ak}:

$$\frac{\mathrm{d}^2\sigma^{Ap\to XN}}{\mathrm{d}t\,\mathrm{d}z} = \sum_{\mathcal{P}}\left[\sum_k \sigma^{Ak}\left([1-z]s\right)\right] f_{\mathcal{P}/p}(z,t) \ , \tag{1.29}$$

with

$$f_{\mathcal{P}/p}(z,t) \ \propto \ \beta^2_{p\mathcal{P}}(t)\,(1-z)^{1-2\alpha_{\mathcal{P}}(t)} \ , \tag{1.30}$$

$$\sigma^{Ak}\left([1-z]s\right) \ \propto \ \beta_{Ak}(0)\,\left[(1-z)s\right]^{\alpha_k(0)-1} \ . \tag{1.31}$$

It should be emphasized, that the z shape of the flux factor is controlled by the trajectory $\alpha(t)$ of the exchanged particle. The pomeron flux factor strongly rises with increasing z. For $z \gg 0.9$, pomeron exchange is the dominant contribution to the reaction $Ap \to Xp$. At fixed t the pion flux factor increases with decreasing z. Thus, pion exchange becomes an important production mechanism for the process $Ap \to XN$ at intermediate values of z ($0.6 \lesssim z \lesssim 0.9$).

In Section 5.7 it will be demonstrated that for $z \geq 0.7$ the deep–inelastic production of leading protons and neutrons with small transverse momentum can be described by Regge phenomenology. Neutron production, which demands charge exchange, is dominated by pion exchange, whereas the observed proton cross section requires pion, reggeon and pomeron exchange.

1.5 DIS on a Virtual Pion

Pion exchange is believed to play a significant role in semi–inclusive hadronic interactions [Pum73]. As demonstrated in the previous section, its contribution to the cross section of the reaction $Ap \to XN$ can be parameterized by means of an universal flux factor. Thus, the question arises, if the proton itself, i. e. its structure function, comprises contributions of virtual pions. In the following section experimental indications for the pion content of the proton are briefly discussed.

1.5.1 The Pion Content of the Proton

Assuming isospin invariance the parton distribution functions of the neutron $q_n(x,Q^2)$ are directly related to the proton quark densities $q_p(x,Q^2)$:

$$u_n = d_p,\ d_n = u_p,\ \bar{u}_n = \bar{d}_p,\ \bar{d}_n = \bar{u}_p,\ s_n = s_p,\ \text{and etc.}$$

Thus, the difference between the proton and neutron structure function, expressed in terms of valence and sea quark distributions in the proton, is given by

$$F_2^p(x,Q^2) - F_2^n(x,Q^2) = \frac{1}{3}x[u_v(x,Q^2) - d_v(x,Q^2)] + \frac{2}{3}x[\bar{u}(x,Q^2) - \bar{d}(x,Q^2)] \ . \tag{1.32}$$

Integrating this equation over x and considering, that the integrated valence quark distributions have to fulfil the sum rules

$$\int_0^1 \mathrm{d}x\,u_v(x,Q^2) = 2 \ , \quad \int_0^1 \mathrm{d}x\,d_v(x,Q^2) = 1 \ , \tag{1.33}$$



Figure 1.7: The ratio of $\bar{d}/\bar{u}$ in the proton as a function of x, obtained by the E866/NuSea Collaboration. The data are compared to the CTEQ4M parameterization of the proton parton distributions [Lai97]. Also plotted is the result from NA51 (from [E866-98]).

one obtains

$$I_G = \int_0^1 \mathrm{d}x \, \frac{1}{x} \left[F_2^p(x,Q^2) - F_2^n(x,Q^2)\right] = \frac{1}{3} + \frac{2}{3} \int_0^1 \left[\bar{u}(x,Q^2) - \bar{d}(x,Q^2)\right] . \quad (1.34)$$

If the sea is flavour symmetric, $\bar{u} = \bar{d}$, the integral over the sea distribution vanishes, yielding the *Gottfried* sum rule [Got67]:

$$I_G = \frac{1}{3} . \quad (1.35)$$

Measurements of deep–inelastic scattering of muons on hydrogen and deuterium by the NMC Collaboration revealed a significant and large violation of the Gottfried sum rule [NMC91, NMC94]. Extrapolating the measurement of $F_2^p - F_2^n$, obtained in the region $0.004 \leq x \leq 0.8$, to the full range in x, they evaluated the integral I_G [NMC94]:

$$I_G = 0.235 \pm 0.026 . \quad (1.36)$$

The violation of the flavour symmetry of the sea has been confirmed in semi–inclusive DIS [HER98] and in dimuon production in Drell–Yan processes [NA51-94, E866-98]. Figure 1.7 shows the recent measurement of the $\bar{d}/\bar{u}$ ratio in the range $0.020 < x < 0.345$, obtained by the E866/NuSea Collaboration [E866-98].

It has been shown that perturbative mechanisms or effects due to the finite quark mass cannot account for the observed large violation of the Gottfried sum rule [Kum98]. The pion cloud model, however, can provide an explanation for the origin of the asymmetry of the flavour sea [Tho83, Hol96a, Kum98]. The pion fluctuates in multi–particle

Fock states within the time allowed by the uncertainty principle. The predominant fluctuations are the πN states due to the low mass of the associated particles. The physical proton can be approximated by the following superposition[11]:

$$|p>_{phys} \simeq (1-a)|p>_{bare} + a\left(\frac{1}{3}|\pi^0 p> + \frac{2}{3}|\pi^+ n>\right) \ . \tag{1.37}$$

The contribution of $|\pi^+ n>$ is twice the one of $|\pi^0 p>$ due to the different Clebsch–Gordon coefficients. Looking at the quark content of the fluctuations, one recognizes immediately that $\bar{d}$ is favoured over $\bar{u}$.

The E866/NuSea Collaboration has demonstrated that their measurement of the light antiquark flavour asymmetry in the nucleon sea is in good agreement with the virtual pion model [E866-98a].

It is interesting to note, that also the inclusive proton structure function can be successfully described by a model, in which the sea partons mainly arise from pionic fluctuations [Edi98].

1.5.2 Pion Exchange in DIS

The experimental results indicate, that the cloud of virtual pions contained in the proton are an important source of the sea quarks. Since deep–inelastic *ep* scattering at low x probes the sea of partons inside the proton, one hopes to obtain information about the structure of the pion at HERA, provided evidence for the virtual pion is experimentally observed [Hol94, Lev95, Kop96].

Already in 1972 Sullivan pointed out, that π^+ exchange plays an important role for the deep–inelastic electro–production of neutrons [Sul72]. Since the production of a neutron requires the exchange of charge for the transition $p \to n$, neutral trajectories ($\mathbb{P}$, f_2, ω) cannot contribute to the cross section via the process sketched in Figure 1.4. It has been shown, that for $0.7 \lesssim z \lesssim 0.9$ and $|t| \lesssim 0.25\,\mathrm{GeV}^2$ neutrons are predominantly produced by pion exchange [Kop96]. Thus, in this kinematic region, the measurement of the semi–inclusive structure function $F_2^{LN(4)}$, as defined in Equation 1.12, might probe the pion content, since Equation 1.27 now simplifies to

$$F_2^{LN(4)}(\beta, Q^2, z, t) \simeq f_{\pi^+/p}(z,t) F_2^{\pi}(x, Q^2) \ . \tag{1.38}$$

Provided the flux factor $f_{\pi^+/p}$ is known, the measurement allows to extract the pion structure function F_2^{π}. The pion flux factor has been constrained using hadron–hadron data [Hol94, Hol96] [12]. In the light–cone representation it is given as a function of z and p_T by [Hol94]

$$f_{\pi^+/p}(z, p_T) = \frac{2}{3}\frac{3}{4\pi}\frac{g^2_{p\pi N}}{4\pi}\frac{[(1-z)^2 m_n^2 + p_T^2]}{z^2(1-z)\left[m_n^2 - M^2_{\pi n}(z, p_T^2)\right]^2}|F(z, p_T^2)|^2 \ , \tag{1.39}$$

where m_n is the mass of the neutron and the non–perturbative strong coupling constant is known from low–energy physics: $g^2_{p\pi N}/(4\pi) = 13.6$ [Tim91]. $M_{\pi n}$ is the invariant

[11] Some models also include $\pi\Delta$ fluctuations.

[12] For more details see Section 3.5.1.



Figure 1.8: The pion flux factor $f_{\pi^+/p}$ integrated over the region $0 \leq p_T \leq 200\,\mathrm{MeV}$. Shown is the parameterization of Holtmann et al. [Hol94] compared to the one by Kopeliovich et al. [Kop96]. The flux–factor, which is implemented in the original version of the RAPGAP generator, has a much lower magnitude.

mass of the intermediate two–body pion–neutron Fock state

$$M^2_{\pi n}(z, p_T^2) = \frac{m_n^2 + p_T^2}{z} + \frac{m_\pi^2 + p_T^2}{1-z} \ . \tag{1.40}$$

The vertex light–cone form factor $F(z, p_T^2)$ is parameterized as [Zol92, Hol96]

$$F(z, p_T^2) = \exp\left[-\frac{R^2_{\pi n}}{2}(M^2_{\pi n}(z, p_T^2) - m_n^2)\right] \ , \tag{1.41}$$

where $R_{\pi n} = 0.93\,\mathrm{GeV}^{-1}$ [Hol96]. The flux factor can be equivalently expressed with respect to z and t, the squared four–momentum transfer between the initial state proton and the final state neutron [Prz97]:

$$f_{\pi^+/p}(z, t) = \frac{2}{3}\frac{3}{4\pi}\frac{g^2_{p\pi N}}{4\pi}(1-z)\frac{-t}{(m_\pi^2 - t)^2}\exp\left[-R^2_{\pi n}\frac{m_\pi^2 - t}{1-z}\right] \ . \tag{1.42}$$

Another evaluation of the flux factor using a reggeized form yields [Kop96]

$$f_{\pi^+/p}(z, t) = \frac{2}{3}\frac{3g^2_{p\pi N}}{16\pi^2}\,(1-z)^{1-2\alpha'_\pi t}\,\frac{-t}{(m_\pi^2 - t)^2}\exp\left[-2R_\pi^2(m_\pi^2 - t)\right] \ , \tag{1.43}$$

where the slope of the pion trajectory $\alpha'_\pi = 1\ \mathrm{GeV}^{-2}$ and the non–perturbative radius is $R_\pi^2 = 0.3\ \mathrm{GeV}^{-2}$. Figure 1.8 demonstrates that the shape and the magnitude of both parameterizations of the pion flux factor are very similar.

Also neutral pions can be exchanged, leading to the production of forward protons. The flux factor $f_{\pi^0/p}$ is exactly half of the charged pion flux factor $f_{\pi^+/p}$:

$$f_{\pi^0/p}(z,t) = \frac{1}{2} f_{\pi^+/p}(z,t) \ , \tag{1.44}$$

because of the different Clebsch–Gordon coefficient (1/3 instead of 2/3). The leading proton, however, cannot provide a conclusive signature for the pion content, since other exchanges ($\mathbb{P}$, f_2, ω) dominate the cross section (see Section 5.7).

The possibility to study the pion structure at low x was the key idea, which inspired the ZEUS and H1 Collaboration to install a forward neutron calorimeter [Bha97, Bec95]. The parton distributions of the pion have previously been constrained only in the $x \gtrsim 0.2$ region by Drell–Yan and prompt photon production measurements of πN scattering (see Section 5.9.1). The determination of $F_2^{LN(4)}$ at HERA would allow to constrain the pion structure function at values of x which are more than an order of magnitude smaller.

In this work, the measurement of $F_2^{LN(4)}$ integrated over $p_T \leq 200\,\mathrm{MeV}$ is carried out. This data is used to constrain F_2^{π} for the first time at HERA (see Section 5.9.2).

1.5.3 Monte Carlo Implementation of the π Exchange Model

The pion exchange formalism is implemented in the RAPGAP and POMPYT Monte Carlo programs [Jun95, Bru96]. Both Monte Carlo generators have been originally developed to simulate diffractive scattering via the exchange of a pomeron.

RAPGAP is a full event generator for deep–inelastic scattering with the capability of simulating diffraction and pion exchange. For the Monte Carlo studies shown in this work, the parameterization of the pion flux factor, which is implemented in the original program (see Figure 1.8), was replaced by the parameterization given in Equation 1.42. The fragmentation in RAPGAP is performed with the Lund string model [And83] as implemented in the JETSET [Sjö94] Monte Carlo program (see Section 1.6.1).

POMPYT also simulates the complete hadronic final state. The same parameterization of the pion flux factor as in the modified RAPGAP version is used. The hard scattering is treated based on the PYTHIA program [Sjö94] and the subsequent hadronization is performed with JETSET.

1.6 An Alternative Approach of Leading Baryon Production

If a DIS process is mediated by a colour–singlet exchange, the proton remnant and the central hadronic system X are colourless, leading to a separation in rapidity[13]. This is not the case in standard DIS. Figure 1.9a shows a typical DIS event initiated by boson–gluon–fusion, which dominates the ep cross section at $x \lesssim 10^{-3}$. The struck gluon takes away colour, which produces a coloured proton remnant as well as a coloured hadronic

[13] The rapidity of a produced particle can be approximated by its pseudo–rapidity $\eta = -\ln\tan(\theta/2)$, where θ is the polar angle of the particle.



Figure 1.9: Rearrangement of the colour topology in the final state. Shown is the string configuration in a DIS boson–gluon–fusion event: (a) conventional Lund string connection of partons, and (b,c) after reconnection due to soft colour interaction. (b) leads to the dissociation of the remnant into the invariant mass M_R, and (c) results in a leading proton (from [Edi96]).

system, originating from the hard photon–gluon interaction. Thus, both systems are connected via colour strings, which results in the subsequential production of hadrons during the fragmentation process in the entire rapidity region between both systems. Therefore the probability of producing a high–energy baryon or a large rapidity gap is strongly suppressed.

1.6.1 The Fragmentation Process

The *Lund string model* [And83] is nowadays the most successful model to describe hadronization processes in the photon fragmentation region. The JETSET program, which is the Monte Carlo realization of this model [Sjö94], is used by most Monte Carlo event generators to simulate hadronization.

The Lund string model is built on the physical picture of a colour flux tube, which is being stretched between the partons moving apart from the common interaction vertex. Due to confinement, the potential energy of the string rises linearly with its dimension, leading to a break–up of the flux tube and the creation of a quark–antiquark pair. This consequentially results in the formation of new strings. Mesons can be produced by a combination of a quark and an antiquark of two adjacent string breakings. In some cases a colour string also breaks up into an antidiquark–diquark pair, leading to the potential production of baryons.

The production cross section for leading baryons predicted by the Lund string model is, however, much lower than in reality, in particular no baryons with very high z, which cause large rapidity gaps, are produced.

In order to explain those events, without using the pomeron exchange concept, alternative models were introduced, which are based on soft non–perturbative interactions occurring before the hadronization process [Buc95, Edi96].

1.6.2 Soft Gluon Interactions

The model of Buchmüller and Hebecker [Buc95] is based on boson–gluon–fusion as underlying partonic process (compare Figure 1.9a). The quark–antiquark pair, which is originally produced in a colour octet state, changes its colour randomly because of additional soft interactions with the colour field of the proton remnant. With a certain probability the original colour octet state might be transformed into a colour singlet state, so that no colour field is located between the quark–antiquark pair and the proton remnant, yielding a gap in rapidity. For a sufficiently fast rotation of the colour spin, the probability for the different colour states should be statistically distributed. The quark–antiquark pair would be in the colour octet state with a probability of 8/9 and in the colour singlet state with a probability of 1/9. This prediction is in agreement with the observation, that a large rapidity gap event is produced in $\sim 10\%$ of all DIS events [ZEU93, H1C94].

Soft Colour Interactions in the LEPTO Monte Carlo Model

A similar approach is the *soft colour interaction* (SCI) model [Edi96, Edi97]. Before the hadronization process, any pair of partons, which take part in the hard interaction, plus the remaining quarks in the proton remnant, can undergo a soft interaction, exchanging only the colour but no momentum. The probability for an interaction, which can be viewed as the exchange of a soft non–perturbative gluon, is described by a single phenomenological parameter R. The soft colour interaction can change the colour topology of the ep scattering event, such that colour singlet subsystems arise separated in rapidity (compare Figure 1.9b,c), which may lead to the formation of a baryon out of the proton remnant. In the case of boson–gluon–fusion this baryon is most likely a leading proton.

A leading neutron can be produced, if a sea antiquark enters the hard interaction. A partner sea quark in the proton remnant is assigned, which is given a momentum based on the assumption, that both quarks originate from the same gluon splitting.



Figure 1.10: z distributions for leading protons and neutrons produced by Lepto. *Shown are the nucleon spectra obtained without ($R = 0$) and with soft colour interaction ($R = 0.5$).*

The proton remnant contains also three valence quarks, which are split into a quark and a diquark. Thus, the remnant partons, together with the interacting sea antiquark, form two colour singlet systems. The situation in which the remnant sea quark is paired with the valence diquark might lead to the production of a neutron.

The SCI model is implemented in the LEPTO Monte Carlo program [Ing97]. LEPTO is a general event generator for deep–inelastic lepton–nucleon scattering, based on leading order electroweak cross sections. QCD matrix elements for gluon radiation and boson–gluon fusion are implemented to first order, and higher order QCD radiation is simulated by parton showers. The fragmentation, which takes place after the soft colour interaction is performed using the JETSET program.

Figure 1.10 demonstrates the effect of the soft colour interaction mechanism. Shown are the z spectra for leading protons and neutrons generated by LEPTO, obtained with and without SCI. The SCI mechanism increases the rate of produced nucleons in the region $z \gtrsim 0.6$. SCI also generates the diffractive peak in the proton spectrum at $z \approx 1$.

Chapter 2

The H1 Experiment at HERA

The physics program outlined in the previous chapter defines the experimental requirements which have to be fulfilled in order to study the production of forward neutrons in deep–inelastic scattering.

First of all, a particle accelerator has to provide the high–energy beams of both electrons (or positrons)[1] and protons. Very high centre–of–mass energies are necessary to study the kinematic region of large values of Q^2 and low values of x. The electron–proton collider HERA (*Hadron–Elektron–Ringanlage*) at DESY (*Deutsches Elektronen Synchrotron*) in Hamburg is dedicated for this purpose.

The scattered electron and most of the hadronic final state in a DIS event have to be measured by a system of nearly hermetic detectors which cover the interaction region. This is the task of the multi–purpose H1 detector. For the detection of leading neutrons and protons, which escape the main detector unseen through the beam pipe, a dedicated *Forward Neutron Calorimeter* (FNC) and a *Forward Proton Spectrometer* (FPS) is used.

In this chapter, the HERA collider and the H1 experiment will be shortly described. A detailed discussion of the FNC will be given in the following chapter.

2.1 The HERA collider at DESY

HERA, which was commissioned in 1992, is composed of two separate storage rings with a circumference of 6.3 km [Wii92]. Electrons and protons are separately accelerated up to energies of 27.5 GeV, respectively 820 GeV [2]. The two beam pipes are brought together at the position of the H1 and ZEUS experiments, allowing the two beams to cross at an angle of 0°.

Besides these collider experiments, which have been taking ep scattering data since 1992, HERA accommodates two additional *fixed target* experiments. HERMES, which started operation in 1995, makes use of the polarized electron beam to study the spin structure of nucleons. HERA–B is a high rate experiment using the halo of the proton beam for collisions with a target made of tungsten wires. It is currently being set up and will begin to study the expected CP violation in the B^0–$\bar{B}^0$ system in 1999.

[1] Throughout this work the incident and scattered electron or positron are referred to as electron.
[2] In 1998 the proton beam energy was increased to 920 GeV.



Figure 2.1: The HERA storage ring and its injection and preacceleration system at DESY. The locations of the H1 and ZEUS detector are indicated.

The HERA beams are both structured into 220 bunches separated by 96 ns, of which on average 190 are filled with particles. This leads to a bunch crossing rate of 10.4 MHz. "Crossings" of empty or unpaired bunches can be used for background studies.

2.2 The H1 Experiment

The design of the H1 detector (see Figure 2.2) is comparable to similar detectors being operated at other collider facilities. Tracking detectors, (2) and (3) [3], are positioned close to the nominal interaction point, followed by an electromagnetic and hadronic calorimeter, (4) and (5) , a superconducting coil, (6), and the iron yoke which is instrumented with muon chambers, (10). In contrast to the detectors at e^+e^- (LEP) and $p\bar{p}$ (Tevatron) colliders, which are build nearly symmetrical with respect to the interaction point, the H1 detector has a large forward backward asymmetry due to the different types of particle beams and due to the difference in the beam energies. The instrumentation of the H1 detector in the forward region, which is defined by the direction of the proton beam, is more complex, allowing to detect most of the hadronic final state of the deep–inelastic reaction, which is boosted along the direction of the proton beam. The backward region is designed to measure the energy and the angle of the scattered electron covering a kinematic area of four–momentum transfers of $1\,\mathrm{GeV}^2 \lesssim Q^2 \lesssim 100\,\mathrm{GeV}^2$. The detector is nearly hermetic, allowing particles only to escape through the forward and backward beam pipe. However a substantial part of the fragments of the target, and therefore also a large part of the energy flow, escapes

[3] Numbers in brackets refer to Figure 2.2.