DESY-THESIS-1999-011
January 1999

INAUGURAL - DISSERTATION

zur
Erlangung der Doktorwürde
der
Naturwissenschaftlich-Mathematischen
Gesamtfakultät
der
Ruprecht-Karls-Universität
Heidelberg

vorgelegt von
Thorsten Wengler, MA
aus Elsterwerda

Tag der mündlichen Prüfung: 27. Januar 1999

Measurement of π°-Meson Cross Sections at low Bjorken-x in Deep-Inelastic ep - Collisions at $\sqrt{s} = 300$ GeV

Gutachter: Prof. Dr. Franz Eisele
Prof. Dr. Otto Nachtmann

Abstract

High p_T π°-meson production cross sections for small polar angles θ_π with respect to the incident proton direction have been measured in deep-inelastic positron-proton scattering at low Bjorken-x. The analysis is based on an integrated luminosity of 5.8 pb^{-1} taken with the H1 detector at HERA in 1996. The π° cross sections are measured as a function of Q^2, as a function of x, and as a function of the transverse momentum $p_{T,\pi}$ and pseudorapidity η_π of the π°-mesons. The measurement extends over two orders of magnitude in x down to $x \approx 5{\cdot}10^{-5}$. High p_T particle production is shown to be strongly correlated to the underlying parton dynamics. The π° cross sections exhibit a strong rise towards low x and Q^2. The relative rate of π°-meson production in DIS events is determined and shows no dependence on x, but decreases with decreasing Q^2 for fixed x. Calculations using $\mathcal{O}(\alpha_s)$ QCD matrix elements and DGLAP-like parton cascades cannot describe the π° cross sections at low x. Including processes where the virtual photon is resolved leads to an improved description of the data. Renormalization and factorization scale uncertainties however limit the precision of the predictions especially for the resolved contributions. A calculation within the BFKL formalism is in good agreement with the data, implying that this approach is a valid approximation of QCD in the considered phase space.

Messung von π°-Meson-Wirkungsquerschnitten bei kleinen Bjorken-x in tief-inelastischen ep - Kollisionen mit $\sqrt{s} = 300$ GeV

Zusammenfassung

Vorgestellt werden Wirkungsquerschnitte für die Produktion von π°-Mesonen mit hohem transversalen Impuls und unter kleinen Winkeln zur Richtung des einlaufenden Protons in tief-inelastischer Positron-Proton Streuung bei kleinen Bjorken-x. Die Messung basiert auf einer Datenmenge von 5.8 pb^{-1}, die 1996 mit dem H1-Detektor an der Beschleunigeranlage HERA aufgezeichnet wurde. Die π° Wirkungsquerschnitte wurden gemessen als Funktion von Q^2 und Bjorken-x sowie als Funktion des transversalen Impulses $p_{T,\pi}$ und der Pseudorapidität η_π der π°-Mesonen. Die Produktion von Teilchen mit hohem Transversalimpuls ist stark korreliert zur Dynamik des partonischen Endzustandes. Die Messung erreicht Werte von $x \gtrsim 5{\cdot}10^{-5}$ und erstreckt sich über zwei Größenordnungen in x. Die Wirkungsquerschnitte zeigen einen starken Anstieg zu kleinen Q^2 und x. Die relative Rate der π°-Produktion in tief-inelastischen Streuereignissen ist nicht von x abhängig, sinkt aber mit kleiner werdendem Q^2 für festes x. Berechnungen unter Benutzung von $\mathcal{O}(\alpha_s)$ Matrixelementen und DGLAP-Partonkaskaden sind nicht kompatibel mit den Daten. Die zusätzliche Berücksichtigung von Prozessen, bei denen die hadronischen Eigenschaften des virtuellen Photons aufgelöst werden, führt zu einer besseren Beschreibung der gemessenen Verteilungen. Die Aussagekraft wird hierbei aber durch eine starke Abhängigkeit dieser Beiträge von der Wahl der Renormierungs- und Faktorisierungsskala begrenzt. Eine Rechnung im Rahmen des BFKL-Formalismus ist in guter Übereinstimmung mit den gemessenen Werten und impliziert damit die Gültigkeit dieser Näherungslösung der QCD im betrachteten Phasenraum.

To my parents and to Joannah.

Uncertainty,
in the presence of vivid hopes and fears,
is painful,
but must be endured
if we wish to live
without the support
of comforting fairy tales.

Bertrand Russel

Contents

List of Tables

List of Figures

Introduction

For the structure of matter, the current understanding is that all matter is made up of point-like particles (leptons and quarks). Their interactions are mediated by exchange particles (bosons). The theoretical framework of particle physics is provided by the *Standard Model*, which describes all the known fundamental forces except gravity. Gravity appears to be much too weak to be significant for ordinary particle processes. The electromagnetic and the weak force are mediated by the massless photon and the massive $Z^\circ/W^\pm$-bosons respectively. This thesis is mainly concerned with the strong force which is responsible for the interactions of quarks and gluons. Quarks are confined in hadrons and cannot be isolated for investigation. It is therefore necessary to study the structure of hadrons.

In this analysis deep-inelastic positron-proton scattering events are used to investigate the properties of the strong interaction. The experimental environment is provided by the accelerator facility HERA which collides counter-rotating positron and proton beams at a center of mass energy of $\sqrt{s} = 300$ GeV. The reaction products of scattering events are recorded by the H1 detector built around the northern of the four collision regions of HERA.

Two problems immediately appear if one tries to compare the measurements to the predictions made by the theory. Firstly, calculations are made for the interacting quarks and gluons, but only the hadrons in which they are confined can be observed. Secondly, although the basic equations of the theory are available, they can only be solved in various approximations. The region of validity for these approximations is not known a priori. It is hence very interesting to measure observables which a) retain the characteristics of the partonic final state after hadronization, and b) explore regions of the phase space where the approximations become critical.

The large amount of data taken by H1 in recent years provides the possibility to study an observable which fulfills both requirements: The production of high transverse momentum ($\gtrsim 2$ GeV) particles at low values of the kinematical quantity Bjorken-x ($x \gtrsim 5 \cdot 10^{-5}$). Especially interesting is the region close to the proton remnant (called the 'forward' region), since here predictions of different approximations differ strongly. Experimentally π°-mesons turn out to be the most suitable particle for this measurement. The observables measured in this analysis are therefore differential forward high p_T π°-meson production cross sections at low Bjorken-x.

The structure is as follows: In Chapter 1 the theoretical framework is developed to the necessary extent. The theory provides the motivation for this measurement and guides the interpretation of the results. The experimental results themselves, however, are of course independent of this framework. The experimental apparatus is introduced in Chapters 2 and 3. Chapter 4 is devoted to the selection of deep-inelastic scattering events from the available data. The necessary information is mostly obtained from the identification and measurement of the scattered positron. The selection of π°-mesons is described in Chapter 5. Neutral pions decay almost exclusively into two photons. The shape of the resulting electromagnetic shower in the calorimeter is analyzed in detail and used to statistically select π°-mesons over the often dense hadronic background. This strategy takes advantage of the very fine granularity of the H1 main calorimeter. In Chapter 6 the neutral pion production cross sections are determined. The distributions are corrected for detector effects and for the influence of QED radiation. Sources of systematic uncertainties are studied to estimate the resulting uncertainty for the neutral pion cross sections. The results are then given including the statistical and the systematic uncertainties associated with each data point. Finally in Chapter 7 the results are interpreted in terms of the available theoretical models. Last but not least the dependency of the predictions given by the various models on some of their respective input parameters is studied in the appendix.

Chapter 1

The Theory of Deep-Inelastic Scattering

The framework for the present analysis is provided by the theory of deep-inelastic positron-proton scattering (DIS). Both the motivation for the measurement and the interpretation of the results are based on this theory. Extensive reviews of theoretical issues in DIS are available [1, 2]. In this chapter the formalism is developed to the extent necessary to position the measurement within the framework available today.

First a number of kinematical quantities are defined which are commonly used to describe DIS processes. The cross section for deep-inelastic positron-proton scattering is introduced and interpreted in terms of the quark-parton model (QPM). Subsequently necessary modifications when considering the influence of the strong interaction as described by Quantum Chromodynamics (QCD) are discussed. Emphasis is placed on the treatment of parton emissions at small values of the Bjorken scaling variable x. The basic features of several parton emission mechanisms are given. The need for observables sensitive to the dynamical features of the emitted partons directly motivates the measurement of high p_T particle cross sections. The sensitivity of high p_T particles to the underlying parton dynamics is demonstrated.

Based on the mechanisms for emitting partons mentioned above, different models are available to predict the properties of the partonic final state. The main features of these models are discussed in some detail. Finally the partons have to be converted into hadrons to allow direct comparisons to the observed particle production. The main features of two hadronization models are described.

1.1 Kinematical Variables in DIS

The kinematics of deep-inelastic ep scattering can be described by two independent variables (such as the polar angle and the energy of the scattered positron) once



Figure 1.1: Four-vectors used to construct Lorentz-invariant kinematical quantities in deep-inelastic scattering.

the center of mass energy is given by the energies of the incident particles. In many cases it is convenient to use a set of Lorentz invariant quantities.

From the four-vectors of the incoming positron (k) and proton (P) (see Figure 1.1) one first of all obtains the center of mass energy squared

$$s = (P + k)^2 . \tag{1.1}$$

With the four-vector of the outgoing lepton (k') the invariant mass squared of the exchanged virtual boson is

$$Q^2 = -q^2 = -(k - k')^2 . \tag{1.2}$$

The two dimensionless variables x and y are also very commonly used. They are constructed using Q^2, P and k such that their values are confined to the range $0 \leq x, y \leq 1$

$$x = \frac{Q^2}{2 \cdot P \cdot q} \tag{1.3}$$

$$y = \frac{P \cdot q}{P \cdot k} . \tag{1.4}$$

The invariant mass squared of the outgoing hadronic system X in Figure 1.1 is given by

$$W^2 = (q + P)^2 . \tag{1.5}$$

Neglecting the lepton and nucleon masses, the following relations hold between these quantities:

$$Q^2 = s \cdot y \cdot x , \qquad W^2 = Q^2 \frac{1 - x}{x} . \tag{1.6}$$

1.2 The Cross Section for Deep-Inelastic ep Scattering

The basic process of deep-inelastic positron-proton scattering is

$$e + P \rightarrow e' + X\,, \tag{1.7}$$

where e and e' are the incoming and the outgoing lepton respectively, P is the incoming proton and X denotes the hadronic final state resulting from the scattering process. The interaction proceeds via the exchange of a neutral (γ/Z°) or charged ($W^\pm$) boson. In the latter case the final state positron is replaced by a neutrino. This analysis is only concerned with neutral current (NC) processes. The NC cross section can be written as

$$\sigma_{\rm NC} = \sigma(\gamma) + \sigma(Z^\circ) + \sigma(\gamma Z^\circ). \tag{1.8}$$

The first and second term describe the contribution from γ and from Z° exchange respectively, the last term is the γZ° interference. Contributions from Z° exchange, however, are suppressed due to the large mass of the Z°-boson:

$$\frac{\sigma(\gamma Z^\circ)}{\sigma(\gamma)} = \frac{Q^2}{Q^2 + M_{Z^\circ}^2} \qquad \text{and} \qquad \frac{\sigma(Z^\circ)}{\sigma(\gamma)} = \left(\frac{Q^2}{Q^2 + M_{Z^\circ}^2}\right)^2 . \tag{1.9}$$

The Z° mass squared $M_{Z^\circ}^2$ is about 84000 GeV^2 [3]. In this analysis only scattering events with $Q^2 < 100$ GeV^2 are considered. The contribution from Z° exchange and γZ° interference can therefore be neglected.

The differential cross section for inelastic positron-proton scattering may be then written as [1]:

$$\frac{d^2\sigma^{\rm NC}_{\rm e+P}(x, Q^2)}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4}\left[xy^2 F_1(x, Q^2) + (1-y)F_2(x, Q^2)\right]\,, \tag{1.10}$$

where α is the electromagnetic coupling constant. A third structure function F_3 takes the parity violating Z° exchange into account, which can be neglected here for the reasons given in (1.9). The other two structure functions $F_1(x, Q^2)$ and $F_2(x, Q^2)$ are required since the cross section has two independent contributions from the absorption of transversely ($\sigma_{\rm T}$) and longitudinally ($\sigma_{\rm L}$) polarized virtual photons. $F_1(x, Q^2)$ describes $\sigma_{\rm T}$ and $F_2(x, Q^2)$ describes the sum $\sigma_{\rm T} + \sigma_{\rm L}$. A longitudinal structure function $F_L(x, Q^2)$ and the ratio R

$$F_L := F_2 - 2xF_1 \qquad R := \frac{F_L}{F_2 - F_L} = \frac{F_2 - 2xF_1}{2xF_1} = \frac{\sigma_{\rm L}}{\sigma_{\rm T}} \tag{1.11}$$



Figure 1.2: The quark-parton model: The interaction at the hadronic vertex is taken to be the sum of incoherent scatterings from the point-like constituents (quarks) of the Proton.

are commonly used, resulting in the following three equivalent expressions for the differential cross section

$$\frac{d^2\sigma^{\rm NC}_{\rm e+P}(x,Q^2)}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4}\left[xy^2F_1(x,Q^2) + (1-y)F_2(x,Q^2)\right] \tag{1.12}$$

$$= \frac{4\pi\alpha^2}{xQ^4}\left[\left(1-y+\frac{y^2}{2}\right)F_2(x,Q^2) - \frac{y^2}{2}F_L(x,Q^2)\right] \tag{1.13}$$

$$= \frac{4\pi\alpha^2}{xQ^4}\left[1-y+\frac{y^2}{2}\frac{1}{1+R}\right]F_2(x,Q^2)\,. \tag{1.14}$$

As yet the structure functions thus defined cannot be calculated but have to be determined from experimental data.

1.3 The Quark-Parton Model

The quark-parton model (QPM) was introduced by Feynman [4] in an attempt to interpret the scaling behavior of the structure functions. According to a prediction by Bjorken [5], the structure functions in the limes of $Q^2 \to \infty$ and $P \cdot q \to \infty$ depend only on the dimensionless variable $x = Q^2/(2{\cdot}P{\cdot}q)$:

$$F_1(x,Q^2) \longrightarrow F_1(x) \tag{1.15}$$

$$F_2(x,Q^2) \longrightarrow F_2(x)\,. \tag{1.16}$$

In the first high energy deep-inelastic electron scattering experiments at SLAC [6] F_2 was indeed observed to be independent of Q^2 for the range $1 < Q^2 < 10$ GeV^2. The quark-parton model is based on two assumptions:

- The hadron participating in the scattering process is made of point-like constituents (partons) which travel in the direction of the hadron. The momentum of the hadron is distributed among the partons. The partons were soon taken to be equivalent to the quarks introduced by Gell-Mann [7] to explain the structure of hadrons.

- At large Q^2 the quarks behave as if they were free inside the hadron during the interaction. The photon-hadron interaction can therefore be expressed as the sum of incoherent scatterings from the point-like constituents (see Figure 1.2).

The argument is that the short time-scale of order $1/\sqrt{Q^2}$ causes the photon to see a frozen state of non-interacting quarks and that the final hadronization process occurs long afterwards. If this is the case, the differential cross section for $e + P \rightarrow e' + X$ is of the form

$$\frac{d^2\sigma^{\mathrm{NC}}_{\mathrm{e+P}}(x,Q^2)}{dxdQ^2} = \sum_i \int d\xi f_i(\xi) \frac{d^2\sigma_i(\xi,Q^2)}{d\xi dQ^2}\,, \tag{1.17}$$

where $f_i(\xi)$ is the probability of finding the quark of flavor i in the proton with a fraction ξ of its momentum. The cross section for electron-quark elastic scattering is [1]

$$\frac{d^2\sigma_i(\xi,Q^2)}{d\xi dQ^2} = \frac{4\pi\alpha^2}{Q^4} e_i^2 \frac{1}{2}\left[(1-y)+\frac{y^2}{2}\right]\delta(x-\xi)\,, \tag{1.18}$$

where the δ-function arises from energy-momentum conservation and from neglecting the mass of the outgoing quark, that is $(\xi P + q)^2 = 0$ which then gives $\xi = Q^2/(2{\cdot}P{\cdot}q) = x$. e_i is the electric charge of the respective quark. Inserting (1.18) into (1.17) and integrating out the δ-function yields

$$\frac{d^2\sigma^{\mathrm{NC}}_{\mathrm{e+P}}(x,Q^2)}{dxdQ^2} = \frac{4\pi\alpha^2}{Q^4}\left[(1-y)+\frac{y^2}{2}\right]\sum_i e_i^2 f_i(x)\,. \tag{1.19}$$

Several important results can now be obtained for the quark-parton model:

- The comparison of (1.19) to (1.12) reveals

$$\frac{F_2}{x} = 2{\cdot}F_1 = \sum_i e_i^2 f_i(x). \tag{1.20}$$

  The QPM thus provides a straightforward relationship between the structure functions and the parton density functions $f_i(x)$.

- The structure functions do not depend on Q^2, but only on x, consistent with the predictions made by Bjorken and the measurements at SLAC.

- $F_2 = 2{\cdot}x{\cdot}F_1$. This is known as the Callan-Gross relationship. It is valid as long as one considers the quarks to be the only constituents of the proton and is a direct consequence of the spin $\frac{1}{2}$ character of the quarks. Looking at (1.11) one gets $R = 0$ and $F_L = 0$, and the longitudinal absorption cross section for virtual photon scattering on quarks vanishes.

If quarks and anti-quarks are indeed the only constituents of the proton, the sum of the momenta they carry should be equal to the momentum of the proton (i.e. the sum of their momentum fractions should be equal to one). Measurements however have demonstrated that

$$\sum_q \int_0^1 x(q(x) + \bar{q}(x))dx \simeq 0.5 \tag{1.21}$$

which indicates that the quarks only carry about half of the momentum of the proton. In Quantum Chromodynamics this deficit in momentum is carried by the gluons.

1.4 Partons in Quantum Chromodynamics

The deviation from unity in (1.21) is a clear indication that quarks cannot be the only constituents of the proton. In addition, measurements covering a larger range of four-momentum transfers Q^2 soon indicated that the scaling behavior of the structure functions is violated. F_2 clearly depends on Q^2 at values of x smaller and larger than accessible in the original SLAC experiments (see Figure 1.3). The quark-parton model therefore cannot be the whole truth.

Quantum Chromodynamics

An explanation of these effects is given by Quantum Chromodynamics (QCD). The essential properties of this theory are [1]:

- Quarks carry the additional degree of freedom *color* as well as electric charge. There are three colors (Red, Green, Blue) which are the charge of the strong interaction.

- Color is exchanged by eight bi-colored gluons. The eight gluons are massless and have spin 1 (bosons). They do not carry electric charge.

- Color interactions are assumed to be a copy of electromagnetic interactions. Quark-gluon interactions can therefore be computed by the rules of QED with the substitution $\sqrt{\alpha} \rightarrow \sqrt{\alpha_s}$ at each vertex and the introduction of color factors. A necessary condition is, of course, that $\sqrt{\alpha_s}$ is small enough to make perturbation theory (as used in QED) applicable.



Figure 1.3: The proton structure function $F_2^p(x, Q^2)$ from H1 and from fixed target experiments [8]. The values are shifted for easier presentation by $c_i(x) = 0.6 \cdot i(x) - 0.4$ starting from $i = 1$ for $x = 0.65$.

- Gluons themselves carry color charge and can interact with other gluons.

The self-coupling of gluons manifests the non-Abelian nature of QCD. It leads, in contrast to the Abelian QED, to an *anti*-screening effect as a result of vacuum polarization around a color charge – the closer one probes this charge the less strong it appears.

To calculate cross sections in QCD one needs to integrate over the entire phase space of (real and virtual) quarks and gluons. These integrals, however, turn out to be divergent. A scheme is therefore defined to leave out the divergent parts of the integrals (regularization). The calculated cross sections become dependent on the energy scale μ_r^2 used in the regularization. This unphysical dependence of the cross sections on the scale μ_r^2 is compensated by defining an effective coupling constant and effective parton masses which absorb the divergent contributions (renormalization). The scale μ_r^2 is referred to as renormalization scale. The coupling constant is defined by the renormalization scheme used and depends on the scale μ_r^2. Requiring the cross sections to be independent of μ_r^2 leads to the renormalization group equation (RGE), which describes the dependence $\alpha_s(\mu_r^2)$. The RGE can be written as a perturbative expansion in α_s. The solution for the coupling constant when calculating to $\mathcal{O}(\alpha_s)$ is

$$\alpha_s(\mu_r^2) = \frac{4\pi}{\beta_0 \ln(\mu_r^2/\Lambda^2)}\,, \tag{1.22}$$

with $\beta_0 = 11 - 2n_f/3$ and n_f the number of quark flavors. Λ is a free parameter of QCD which has to be determined from experimental results. The value of α_s is usually given at the mass squared of the Z°. The current world average is $\alpha_s(M_Z^2) = 0.119 \pm 0.002$ [3]. The behavior of the coupling constant α_s has two immediate consequences:

- At high energies α_s is small and the quarks are 'asymptotically free'. In this kinematic region perturbation theory may be used to perform calculations within QCD.
- At large distances the coupling constant increases, leading to the confinement of quarks within hadrons. Here non-perturbative methods are needed.

It also means that in order to use perturbation theory at all, it must be possible to separate short-range from long-range processes. This is the idea of factorization, which will be important in the next section.

QCD corrections to the structure of the proton

In QCD the picture of the proton as used in the quark-parton model is modified by allowing gluon radiation. The QPM in this sense can be regarded as the lowest order process of QCD. Introducing the gluons provides ways to overcome the shortcomings of the QPM in describing experimental observations:



Figure 1.4: Gluon radiation and scaling violations: The higher resolution in (b) causes the virtual photon to 'see' more partons which on average carry a smaller fraction of the proton's momentum.

- The fractional momentum sum no longer deviates from unity. The missing momentum is carried by (electrically) uncharged gluons.
- The violation of Bjorken-scaling at large and small Bjorken-x can be explained by allowing the quarks to radiate gluons which in turn can split into $q\bar{q}$-pairs. The resolution increases with increasing energy (e.g. Q^2 in DIS). More and more gluon radiation processes and $q\bar{q}$-splittings can therefore be resolved, so that the apparent number of partons which share the proton's momentum increases (see Figure 1.4). Hence there is an increased chance of finding a quark at small x and a decreased chance of finding one at high x when moving towards larger Q^2. The result is a scale-violating Q^2 dependence of the structure functions.

Another consequence is, of course, that the simple relationship between the structure functions and the probability distribution of the quarks in momentum space no longer holds in its simple form (1.20). It was, after all, based on the assumption of an incoherent sum of γq scatterings. The separation of long range and short range processes according to the idea of factorization however allows a more general formulation for the structure functions

$$F_2(x,Q^2) = \sum_{i=q,g} \int_x^1 d\xi \; C_i^V \left(\frac{x}{\xi}, \frac{Q^2}{\mu_r^2}, \mu_f^2, \alpha_s \right) \cdot q_{i,h}\left(\xi, \mu_r^2, \mu_f^2, \alpha_s\right) \,, \tag{1.23}$$



Figure 1.5: (a) The ep cross section is factorized into positron-parton cross sections σ_{ei} and parton densities $q_{i,p}$. (b) shows the lowest order diagram contributing to σ_{ei} in (a).

where the sum extends over all (anti-)quark and gluon densities. C_i^V are the coefficient functions for each type of parton i and exchanged boson V (here the virtual photon). μ_r^2 is the renormalization scale introduced above. The factorization scale μ_f^2 defines the point of separation between long range and short range phenomena (see Figure 1.5). If the relevant scale for a process is larger than μ_f^2, this process is considered calculable in perturbative QCD. In the example above Q^2 is taken to be the relevant scale. Everything below μ_f^2 is absorbed into the parton densities $q_{i,h}$. This definition determines the properties of the coefficient functions C_i^V and the parton densities $q_{i,h}$:

- C_i^V includes the 'hard' subprocess, which can be calculated using perturbation theory. The form of the coefficient functions depends on the parton-flavor i and the exchanged boson V. It does, however, not depend on the type of hadron entering the process. C_i^V in (1.23) can therefore be used to calculate the structure function of any hadron such as the proton, neutron or pions.

- The parton densities $q_{i,h}$ are specific to the hadron h. They are, however, independent of the 'hard' subprocess under consideration. Once they have been extracted from the comparison with experimental data, the parton densities can therefore be used to describe any hard interaction. In this sense they are refered to as *'universal parton densities'*.

One should note that while C_i^V and $q_{i,h}$ depend on the choice of renormalization and factorization scale, $F_2(x, Q^2)$ does not. Any dependence on μ_f^2 and μ_r^2 therefore has to cancel when the sum and the integral are carried out in (1.23). In practice the perturbative series for the coefficients C_i^V has been calculated completely only to next-to-leading order in α_s for the inclusive ep cross section. Some dependence on μ_f^2 and μ_r^2 hence remains due to higher order effects which then also limit the notion of universality for the parton densities determined from experimental data.

Since these equations can not be solved completely so far, several approximations of QCD are commonly used which are valid in different regions of the phase space. In the perturbative expansion terms of the form $(\alpha_s \ln(Q^2/Q_0^2))$, $(\alpha_s \ln(1/x))$ and mixed terms $(\alpha_s \ln(Q^2/Q_0^2) \ln(1/x))$ appear. In the DGLAP scheme $(\alpha_s \ln(1/x))$ terms are neglected. The BFKL approach on the other hand takes the $(\alpha_s \ln(1/x))$ terms to be dominant over the $(\alpha_s \ln(Q^2/Q_0^2))$ terms used in the DGLAP approximation. As x decreases the mixed terms might be dominant if Q^2 is not too small. The CCFM equation finally promises a smooth transition between the domains of small and large x. These approximations will be treated in more detail in the following sections.

1.5 The DGLAP Evolution Equations

It has not been possible so far to calculate the parton densities $q_{i,h}$ from first principles in QCD. However, towards large Q^2 (the coupling constant α_s decreases) one can employ perturbative calculations to predict the evolution of the parton densities. The evolution of the quark and the gluon densities in this region is given by the DGLAP (Dokshitzer, Gribov, Lipatov, Altarelli, Parisi) equations [9]:

$$\frac{dq_i(x,Q^2)}{d\ln Q^2} = \frac{\alpha_s}{2\pi}\int_x^1 \frac{d\xi}{\xi}\left[q_i(\xi,Q^2)P_{qq}\left(\frac{x}{\xi}\right) + g(\xi,Q^2)P_{qg}\left(\frac{x}{\xi}\right)\right], \tag{1.24}$$

$$\frac{dg(x,Q^2)}{d\ln Q^2} = \frac{\alpha_s}{2\pi}\int_x^1 \frac{d\xi}{\xi}\left[\sum_i q_i(\xi,Q^2)P_{gq}\left(\frac{x}{\xi}\right) + g(\xi,Q^2)P_{gg}\left(\frac{x}{\xi}\right)\right]. \tag{1.25}$$

The index i denotes the quarks and anti-quarks of all flavors. The splitting functions $P_{i,j}(z)$ give the probability for parton branchings $q \to qg$, $g \to gg$ and $g \to q\bar{q}$, where the daughter parton i carries a fraction $(1-z)$ of its mother's (j) momentum (see Figure 1.6). The splitting functions are calculable by perturbative expansion

$$P_{ij}(z,\alpha_s) = \frac{\alpha_s}{2\pi}P_{ij}^0 + \left(\frac{\alpha_s}{2\pi}\right)^2 P_{ij}^1 + \cdots \tag{1.26}$$

In leading order the splitting functions are:

$$\begin{aligned}
P_{qq}^0(z) &= P_{gq}^0(1-z) = \frac{4}{3}\left[\frac{1+z^2}{(1-z)_+}\right] + 2\cdot\delta(1-z) \\
P_{qg}^0(z) &= \frac{1}{2}(z^2+(1-z)^2) \\
P_{gg}^0(z) &= 6\left[\frac{z}{(1-z)_+} + \frac{1-z}{z} + z(1-z)\right] + (11 - \frac{n_f}{3})\cdot\delta(1-z)\,.
\end{aligned} \tag{1.27}$$



Figure 1.6: Feynman graphs for splitting functions. The following processes are shown (from left to right): $q \to q'g$ (P_{qq}, P_{gq}), $g \to q\bar{q}$ (P_{qg}) and $g \to gg$ (P_{gg}).

The subscript '+' indicates that the singularities at $z = 1$ (corresponding to the collinear emission of partons) are regularized by using the prescription

$$\int_0^1 dz \frac{f(z)}{(1-z)_+} := \int_0^1 dz \frac{f(z) - f(1)}{1-z} \tag{1.28}$$

for any function $f(z)$. The DGLAP equations can be obtained using the Leading-Log Approximation (LLA). In leading order α_s the dominant contributions are here given by $(\alpha_s \ln(Q^2/Q_0^2))^n$ terms which can be summed for all orders n. The leading role of the Q^2 terms is inherent in the perturbative ansatz. Dokshitzer [9] showed that in an axial gauge (where the gluon only has the two physical polarization states) the LL(Q^2) is equivalent to the sum of ladder graphs as shown in Figure 1.7. The LL(Q^2) contribution comes from the region of phase space given by the following two conditions:

- The transverse momenta $k_{T,i}^2$ of the emitted partons are strongly ordered such that $k_{T,i}^2 \ll k_{T,i+1}^2 \ll \cdots \ll Q^2$.
- The longitudinal momenta x_i of the emitted partons satisfy the constraint $x_i > x_{i+1} > \cdots > x$.

The region where the LLA is applicable at all is given by the use of perturbation theory and by neglecting $\ln(1/x)$ terms in the LLA sum:

$$\begin{aligned} \alpha_s(Q^2)\ln(Q^2/Q_0^2) &\sim 1 \\ \alpha_s(Q^2)\ln(1/x) &\ll 1 \\ \alpha_s(Q^2) &\ll 1 \end{aligned} \tag{1.29}$$

Hence, if the terms including $\ln(1/x)$ become larger, the DGLAP equations as described above are no longer applicable. At HERA values of $x < 10^{-4}$ have been reached, where the influence of these logarithms might well be visible. It should be noted that in the DGLAP scheme no statement is made about the x dependence



Figure 1.7: Ladder diagram to visualize the parton evolutions in QCD. The gluon longitudinal and transverse momenta are labeled x_i and $k_{T,i}$.

of the parton distributions. One can therefore use input parton distributions which rise at small x to describe the data measured at HERA to the lowest available values of $Q^2 \simeq 0.5$ GeV^2 [10]. When moving towards smaller x, contributions $\alpha_s \ln(1/x)$ and $\alpha_s \ln(1/x) \ln(Q^2/Q_0^2)$ become important. If the latter are dominant the double-leading-log approximation (DLLA) can be used, which in addition to the strong ordering in $k_{T,i}^2$ of the emitted partons also requires the longitudinal momenta to be strongly ordered: $x_i \gg x_{i+1} \gg \cdots \gg x$.

1.6 The BFKL Equation

When x becomes small but Q^2 is not large enough for the DLL scheme to be applicable, the DGLAP approximations cease to be valid. In this region one needs to sum diagrams which are leading in $\ln(1/x)$ independent of $\ln Q^2$. This is carried out by the BFKL (Balitsky, Fadin, Kuraev, Lipatov) equation [11]. Here the gluon ladder need not be ordered in k_T. The gluon distribution therefore is not integrated over k_T. This unintegrated gluon density $f(x, k_T^2)$ relates to the more familiar gluon density as follows

$$xg(x,Q^2) = \int_0^{Q^2} \frac{dk_T^2}{k_T^2} f(x,k_T^2). \tag{1.30}$$

The BFKL equation can then be written as

$$\frac{df(x,k_T^2)}{d\ln(1/x)} = \int dk_T'^2\, K(k_T^2,k_T'^2)\, f(x,k_T'^2) = K \otimes f = \lambda f\,, \tag{1.31}$$

which describes the evolution in $\ln(1/x)$ of the unintegrated gluon density. The solution of this equation is controlled by the largest eigenvalue λ of the kernel K [12]. In leading order in $\ln(1/x)$ and for fixed α_s one obtains a very steep power law

$$xg(x,Q^2) \sim f(Q^2)x^{-\lambda}\,, \quad \text{with} \quad \lambda = \alpha_s \frac{12\cdot\ln 2}{\pi} \simeq 0.5\,. \tag{1.32}$$

Recent developments however indicate that the inclusion of the running coupling and the next-to-leading order contributions have a large effect on λ [13]. The different scope of the BFKL evolution leads to an ordering scheme for the parton emissions in LL$(1/x)$ which is very different from the DGLAP-type ordering:

- The transverse momenta $k_{T,i}^2$ of the emitted partons are unordered. The parton emissions perform a *random walk* in k_T^2 space.

- The longitudinal momenta x_i of the emitted partons are strongly ordered: $x_i \gg x_{i+1} \gg \cdots > x$.

The fact that the BFKL evolution is not ordered in transverse momentum leads to a diffusion of k_T^2, which can extend into the infra-red region. This is usually avoided by introducing a cut-off parameter k_0^2, below which gluon emissions are not allowed. An unwanted side effect of this procedure is, however, that the results thus obtained show a dependence on this parameter k_0^2.

1.7 The CCFM Equation and the Linked Dipole Chain Model

As stated above, the BFKL and DGLAP equations are valid in quite separate regions of phase space, depending on which set of logarithms is dominant. Catani, Ciafaloni, Fiorani and Marchesini (CCFM) [14] have developed a theoretical framework which treats both the small and the large x region in a unified way. The CCFM equation is based on the coherent radiation of gluons, which leads to an angular ordering of the gluons along a chain of multiple emissions. The CCFM equation is defined in



Figure 1.8: The LDC model: fan diagram for a DIS event with n initial emissions. q_i denote the emitted ISB partons, k_i denote the propagators.

terms of a scale dependent unintegrated gluon density $F(x, k_T^2, Q^2)$ quite analog to the BFKL equation above. The integrated gluon density can be recovered by

$$xg(x, Q^2) = \int^{Q^2} \frac{dk_T^2}{k_T^2} f(x, k_T^2, Q^2)\,. \tag{1.33}$$

The angular ordering introduces the additional scale which is needed to specify the maximum angle for a gluon emission. At very small x the angular ordering does not constrain the transverse momenta along the chain and F becomes the Q^2-independent gluon of the BFKL equation. Moving towards larger x the angular ordering becomes equivalent to an ordering in the gluon transverse momenta and one recovers the behavior of the DGLAP evolution [15].

Recently the CCFM approach has been reformulated to obtain the Linked Dipole Chain Model (LDC). The possible parton emissions are separated into Initial State (ISB) and Final State Bremsstrahlung (FSB). The ISB set in the CCFM model is defined by requiring that the emissions are ordered in rapidity and energy. In the LDC model an additional constraint on the transverse momenta in each emission is introduced [16] (see Figure 1.8 for the definition of the variables)

$$q_{T,i} > \min\left[(k_{T,i}), (k_{T,i-1})\right]\,. \tag{1.34}$$

Emissions which violate this condition are moved into the FSB set. The FSB kinematical region in the LDC thus formulated is consistent with the Color Dipole Model [17] for parton cascades. The initial ISB partons are assumed to form a chain of linked color dipoles which can radiate partons independently.



Figure 1.9: The transverse energy flow as a function of pseudorapidity in the hadronic cms [18]. The proton direction is to the left. The errors represent the statistical error only. Three QCD models are overlaid for comparison.

1.8 The Merits of High p_T Particle Observables

If there is one thing to be learned from the previous sections, it is that observables are needed to provide cornerstones on the road to further theoretical development. The most insight into the mechanisms at work will be provided by the approach which predicts the widest range of observations with the least number of starting assumptions. One of the most important measurements driving the understanding



Figure 1.10: The forward jet cross sections as a function of Bjorken-x as measured by H1 [19]. The jets fulfill the condition $0.5 < p_{T,j}^2/Q^2 < 2.0$. The errors shown are the statistical (inner) and the systematic error added in quadrature. The overlaid lines represent QCD model predictions (see [19] for a discussion of the models).

of the structure of hadrons and of processes involving multi-parton radiation has been the determination of the hadron structure functions such as $F_2^P(x, Q^2)$. Data is available down to $x \sim 10^{-5}$ and up to $Q^2 \sim 10^4$ GeV^2. The following situation has now emerged:

- QCD analyses based on the DGLAP evolution (see Section 1.5) in next-to-leading order α_s give a good description of the data down to very low values of Q^2 and x (see Figure 1.3), where in fact $\ln(1/x)$ contributions should become important.
- The problem in identifying the underlying small x dynamics is not due to the insufficient kinematical reach of the experiments, but rather due to the parametric freedom in specifying the initial conditions of the theory [10].
- The measurement of F_2 is too inclusive. One should explore properties of the hadronic final state in deep-inelastic scattering.

When trying to identify the dynamics leading to the *partonic* final state by measuring observables of the *hadronic* final state, another problem presents itself: hadronization. The process of converting the partonic properties of an event (as predicted by a theoretical model) into observable hadrons may wash out differences between the various models. A striking example of such an effect is the measurement of the transverse energy flow by H1. Although different models predict very different E_T-flows for the partonic final state [20], they are compatible with the data and with each other for the hadronic final state (see Figure 1.9). One therefore needs observables which retain the characteristics of the underlying parton level even after hadronization. The classical way of measuring the properties of an outgoing hard parton is based on the idea that whatever the final observable particles are, they will be close to the original parton in the appropriate space (e.g. transverse momentum relative to the original direction). Various algorithms are then used to collect all these particles into a 'jet'. An elegant way to probe the small x behavior of QCD was proposed by Mueller [21]. He suggested the study of deep-inelastic events (x, Q^2) which contain an identified energetic jet (x_j, k_{Tj}^2) close to the proton remnant:

- The jet should satisfy $k_{Tj}^2 \simeq Q^2$, which leaves only little phase space for the DGLAP evolution (see Section 1.5).
- At the same time k_{Tj}^2 has to be sufficiently large to avoid diffusion into the infra-red region as described in Section 1.6.
- The parton densities for this process are to be evaluated at (x_j, k_{Tj}^2), where they are well known from the global analyses, thereby reducing the ambiguities from the choice of the non-perturbative parton input.

This kind of measurement has been carried out by ZEUS [22] and H1 [19]. Part of the H1 results are shown in Figure 1.10. A detailed discussion of the results can be found in [19]. The differences between the various QCD models are clearly visible for this observable (defined for the hadronic final state).

From the above one can conclude that the interesting phase space is to be found close to the proton remnant and at high transverse momenta. It is mostly experimental arguments which lead the way from here to the measurement of high p_T π°-mesons. The first requirement, however, is that the π°-mesons are strongly correlated to the underlying parton dynamics. This is demonstrated in Figure 1.11. The Monte Carlo event generator LEPTO [23] is used to generate DIS events. These events are subjected to hadronization as implemented in the package JETSET [24] to convert the partons into observable hadrons. Events with a π°-meson of $p_{T,\pi} >$ 2 GeV are selected if the meson has a polar angle of $5^\circ < \theta_\pi < 25^\circ$ with respect to the incident proton direction. A parton is not necessarily a well defined object in an event generator. The π°-mesons are therefore compared to jets of partons found in the same event. A k_T-type jet-algorithm [25] was used to define the jets. In nearly all ($> 99\,\%$) selected π° events a jet was found. Figure 1.11 (a) shows the correlation in pseudorapidity ($\eta = -\ln(\tan(\theta/2))$) between the π°-meson and the parton jet. The correlation is indeed very strong, indicating that the direction of the initial hard parton(-jet) is in most cases not altered significantly by hadronization effects. Figure 1.11 (b) demonstrates how much of the transverse momentum of the parton jet is carried by the π°-meson. Since the mean value is about 0.4, the initial parton jet on average carries 2.5 times the transverse momentum of the π°-meson measured.



Figure 1.11: The angular correlation (a) and p_T-ratio (b) between π°-mesons and parton-jets in simulated deep-inelastic scattering events. Events are selected to contain a π°-meson with $p_{T,\pi} > 2$ GeV and a polar angle of $5^\circ < \theta_\pi < 25^\circ$ with respect to the incident proton direction. Parton jets are defined using a k_T-type jet algorithm [25].

It now remains to be argued why the π°-mesons are the most suitable particles to be used in such a measurement and what their advantages over the commonly used jet observables are:

- Due to their integrating nature, jet algorithms will pick up an increasing amount of unwanted energy from the proton remnant when moving towards smaller polar angles. The π°-meson as a single particle does not have this problem.
- A jet is defined by the jet algorithm. The properties of the jet algorithm will therefore influence the result. The notion of a particle is much more independent.

Experimentally the π°-meson is the most suitable particle to be measured at high transverse momenta because:

- π-mesons are the lightest hadrons and therefore the most numerous in the hadronic final state.
- The measurement of charged particles in tracking chambers at high momenta is difficult since the tracks are essentially straight.
- The π°-mesons decay into two photons, which cause a purely electromagnetic shower in the calorimeter. Electromagnetic showers are much more compact than showers of hadronic origin. It will be shown in the course of this analysis



Figure 1.12: Tree level diagrams for γP scattering (b) contributing to σ_{ei} in (a).

that a π°-signal can be selected based on the different nature of electromagnetic and hadronic showers in the calorimeter.

This method can be used up to very high π° energies and down to small angles with respect to the proton remnant – this is the phase space of interest defined above!

To summarize the discussion above: The measurement of π°-meson production provides observables which are strongly correlated to the underlying parton dynamics. Experimentally they allow the exploration of the interesting phase space region close to the proton remnant where differences in the partonic activity of various models are expected to be pronounced. The measurement is not limited as far as high transverse momenta of the π°-meson are concerned. The available reach in $p_{T,\pi}$ is given by the available amount of data only.

1.9 QCD Models

Predictions made by QCD inspired models involve two different aspects: the short range processes which can be calculated perturbatively, and the long range non-perturbative effects of hadronization. The perturbative part of QCD is treated either with fixed order (α_s) calculations of the matrix elements, or with parton showers, or a combination of both. Perturbative QCD makes predictions for the *partonic* final state. Hadronization is the link to the observable world. The string fragmentation model and the cluster fragmentation model are used by the QCD models which will later be compared to the experimental results. A short description of both fragmentation models is given.



Figure 1.13: Parton cascades in deep-inelastic scattering: (a) Leading Log parton shower: initial and final state parton showers are initiated by successive parton branching. (b) The color dipole model: Partons are radiated from color dipoles stretched between outgoing quark and remnant, and between the resulting partons.

Matrix Elements

In fixed order $(\alpha_s)^n$ perturbation theory, all graphs to order n (including the virtual corrections from loop diagrams) are calculated and summed to obtain a prediction for the process under investigation. In Figure 1.12 (b) tree-level diagrams for γP scattering are shown. The calculations are usually done numerically employing Monte Carlo techniques. Up to now order $n = 2$ calculations are available which, for instance, predict dijet production to NLO(α_s).

Leading Log parton cascade

Complete fixed order $(\alpha_s)^n$ calculations are not available for $n > 2$. Higher order effects can be approximated using the parton cascade ansatz [26]. It has the advantage that arbitrarily high orders in α_s can be simulated, but only in the Leading Log approximation. The quark struck by the electroweak boson in deep-inelastic scattering may emit partons both before and after the boson vertex, corresponding to initial and final state parton showers respectively (see Figure 1.13 (a)). A parton (close to mass shell) in the incoming nucleon can initiate an initial state parton cascade. In each branching one parton becomes increasingly space-like ($m^2 < 0$) and the other is on-shell or has a time-like virtuality ($m^2 > 0$). The result of this initial state (space-like) cascade is a space-like quark which interacts with the electroweak boson. The outgoing quark is either time-like or on-shell. If the outgoing quark is time-like, the result is a final state (time-like) cascade. The off-shell mass is reduced by branching into daughter partons until all partons are essentially on-shell. Any time-like parton from the initial state cascade will undergo a similar development. Both the initial and the final state cascade are based on the branching processes



Figure 1.14: (a) The string fragmentation model: New $q\bar{q}$-pairs are created when the string between the two separating partons of a $q\bar{q}$-pair breaks up. Mesons are formed when the string energy is too small to create further $q\bar{q}$-pairs. (b) The cluster fragmentation model: gluons (double lines) are split non-perturbatively into $q\bar{q}$-pairs. Color neutral clusters are formed which decay into mesons.

$q \to qg$, $g \to gg$ and $g \to q\bar{q}$ as described by the DGLAP splitting functions in the LL(Q^2) approximation of QCD (see Section 1.5).

Color Dipole parton cascade

An alternative description of parton showers is provided by the Color Dipole Model (CDM) [17]. Unlike the leading-log parton model the CDM does not divide the QCD cascade into an initial and a final state. Instead all parton radiation is described starting from a color dipole formed between the struck quark and the hadron remnant. Subsequent gluon radiation emanates from color dipoles spanned between the newly created color charges and the original color charges. The process is iterated until all the available energy has been used up. In DIS one of the color charges, the proton remnant, is not point-like. The emission of small wavelengths from an extended antenna is suppressed. The CDM takes this effect into account by introducing an appropriate suppression factor. The boson-gluon fusion (BGF, see Figure 1.12) process is not described by this approach, but has to be inserted by hand. The k_T phase space for gluon emission is bounded by the k_T of the previous emission. The emissions are uniform in rapidity. If one arranges the gluon emissions according to increasing rapidity, the final gluon configuration will therefore be unordered in k_T.

String fragmentation

The string model for fragmentation [24] starts by stretching a color field between the two separating partons of a $q\bar{q}$-pair. The color field is thought to be string-like with a constant energy density per unit length. In DIS strings will also be stretched between the (anti-)quark and the proton remnant. When the energy stored in the string is large enough, the string breaks up by creating a new $q\bar{q}$-pair, forming new string pieces. The process is iterated until all the available energy has been used up. The resulting string fragments are combined into mesons and baryons (see Figure 1.14 (a)). Gluons are realized as kinks in the string, which receive the four-momentum of the gluon.

Cluster fragmentation [27]

In the cluster fragmentation approach color singlets are formed from the partons present after the perturbative evolution. First all gluons are forced to split into $q\bar{q}$-pairs. Neighboring (color connected) $q\bar{q}$-pairs are then combined into low mass color neutral clusters. These clusters are then taken to be superpositions of meson resonances, and they decay into mesons (and baryons) depending on the available phase space (see Figure 1.14 (b)).

Chapter 2

The ep Collider HERA

The **H***adron*-**E***lectron*-**R***ing*-**A***ccelerator* is the first ever storage ring facility to provide colliding beams of electrons and protons. The two main accelerators reside in a tunnel of 6.3 km circumference. They produce counter-rotating beams of 27.5 GeV (electrons) and 820 GeV (protons) resulting in a center of mass energy of 300.3 GeV. The electron ring uses conventional magnet technology, whereas superconducting magnets are needed for the bending fields in the proton ring. Four interaction regions are available to the experimentalists. H1 (north hall) and Zeus (south hall) study electron-positron scattering events. Both experiments have been taking data from ep collisions since 1992. HERA-B (west hall) only utilizes the proton beam in fixed target mode by inserting tungsten wire targets into the beam halo. The HERA-B detector components are expected to be fully operational in 1999. The main focus will be the search for decay modes in the $B^\circ/\bar{B}^\circ$ system indicating a violation of the combined symmetry of charge conjugation and parity (CP). Collisions of the electron beam in polarized mode with a polarized gas target (H_2, D, ^3He,^4He) have been used by the HERMES (east hall) collaboration since 1995 to study phenomena related to the spin of the colliding particles.

A schematic layout of the accelerator complex is shown in Figure 2.1. A chain of accelerators originally built for previous experiments now serves to produce the electron and proton beams at the energy required to inject them into the two main accelerators. Protons are produced by sending negatively charged hydrogen ions through a stripper foil. These protons are then collected into bunches and accelerated to 7.5 GeV in DESY III before they are transfered to PETRA. After the maximum of 70 bunches in PETRA has been reached, the protons are accelerated to 40 GeV and injected into HERA. With a capacity of 210 bunches in HERA 3 iterations of this process are necessary for a complete proton fill.

Electrons are provided by a 500 MeV linear accelerator. The electrons are first accumulated into a single bunch before they are accelerated to 7 GeV in DESY II and transfered to PETRA. Again 70 bunches are collected here before they are



Figure 2.1: Schematic overview of the HERA accelerator complex.

accelerated to 11 GeV and injected into the HERA electron ring. Also here three iterations are needed for a complete fill. Normally the proton beam is produced first and accelerated to its final energy of 820 GeV. Afterwards the electron beam is inserted and brought to 27.5 GeV before both beams are optimized for luminosity production.

During 1994 the electrons were replaced by a positron beam. Positively charged ion-clusters produced by getter pumps which maintain the high vacuum in the electron ring had been found to reduce the lifetime of the beam. These ions are repelled by the positron beam, which therefore has a longer lifetime.

Chapter 3

The H1 Detector

Many aspects of the physics to be studied and the variety of the signatures they produce were not known at the design stage of the detector. To exploit the potential provided by the high energy ep collisions as much as possible, the H1 apparatus was not constructed for a number of specialized measurements but rather as a general purpose detector for ep collisions. This concept is based on a full coverage of the solid angle around the interaction point with the exception of the beam holes where the colliding beams enter the apparatus. Figure 3.1 shows an isometric view of the H1 detector. The asymmetric design of the detector is motivated by the asymmetric beam energies. The center of mass frame of the colliding particles is boosted with respect to the laboratory frame in the direction of the much more energetic proton beam ($\gamma_{\rm cm} = 2.86$). The highest particle and energy densities are expected in this region and the detector is considerably more massive and highly segmented here.

The laboratory frame is described by a right-handed coordinate system such that the proton beam moves along the $+z$ direction. The origin is defined to be the nominal interaction point. The x-axis and y-axis point to the center of the ring and upwards respectively. For many physics parameters, a spherical coordinate system is more appropriate. The azimuthal angle (ϕ) is defined such that $\phi = 0$ points along the x-direction. The polar angle (θ) is measured relative to the z-axis ($\theta = 0$ corresponds to the $+z$-direction). For ultra-relativistic particles the pseudorapidity η is commonly used in place of θ:

$$\eta = -\ln\left[\tan\frac{\theta}{2}\right] = \tanh^{-1}(\cos\theta)\,. \qquad (3.1)$$

For massless particles η is the same as the rapidity y, defined by

$$y = \frac{1}{2}\ln\left[\frac{E+p_Z}{E-p_Z}\right]\,. \qquad (3.2)$$

The dimensions of the H1 detector are approximately 12 m $\times$ 10 m $\times$ 15 m, with a total weight of about 2800 tons. No single device can perform all particle