Tests of Perturbative and Non-Perturbative QCD from Identified Proton, Kaon and Pion Studies in Deep Inelastic Scattering *ep* Interactions at HERA

Glen R. White *MPhys.(Hons.)*

Department of Physics, Lancaster University

This thesis is submitted in part fulfillment

of the requirements for the degree of Ph.D

from Lancaster University.

No part of this thesis has been, or will be,

submitted to any other university.

November 14, 2000

Abstract

Measurements of kaons, pions and protons identified using the dE/dx technique have been made in deep-inelastic scattering ep interactions at HERA in the kinematic range $5 < Q^2 < 70$ and $10^{-5} < x < 10^{-2}$. Tests of QCD were made possible through the production of transverse momenta and pseudo-rapidity spectra with comparisons made to ARIADNE and LEPTO Monte Carlo models, utilising the JETSET hadronisation model with tuning taken from DELPHI experiments, and with HERWIG using the cluster fragmentation model. Further measurements were made of the proton-antiproton asymmetry $A_B = \frac{2.(N(p)-N(\bar{p}))}{N(p)+N(\bar{p})}$ and compared to measurements made elsewhere using photo-production data and recent theoretical results. A search for QCD instanton induced events was also made using the kaon and pion sample. Results from kaon and pion spectra show a preference for the ARIADNE model with the DELPHI tuned JETSET parameters. Models with the DELPHI tuning are preferred over LUND default values, which includes a lower than default value for the strangeness suppression parameter $\lambda_s = 0.23$. None of the Monte Carlo models tested agreed with the proton and antiproton measurements, all over-estimating the measured yields. The proton-antiproton asymmetry was found to be zero within experimental errors, in the kinematical range of the measurements. An excess in strange particle yield over standard DIS background was found in a set of events selected for their instanton-like properties, although not statistically significant enough to draw firm conclusions.

To my wife, Denise, and to my family. Thank you all for your support.

“Those to whom everything is clear are unhappy people.”

- Louis Pasteur

Contents

List of Figures

List of Tables

Acknowledgements

Firstly, I would like to thank the Lancaster University HEP group for giving me the opportunity to study for this PhD in the first place. I would especially like to thank Terry Sloan, my supervisor, and Rob Henderson for their patient help and guidance throughout my years at Lancaster, and for putting up with no end of stupid questions! Also, thanks go to the whole HEP group, most of whom I have had contact with in some form or other. Especially, thanks to Alex and Rob for keeping the computer network operational, and Tim for his help with some statistical questions.

An especially big thank you to Dave Milstead for his great guidance and patience, without which I may well have been still sat at DESY looking bemused in front of a computer screen to this day! I would also like to thank all who helped me during my stays at DESY, especially those in the Energy Flow and Spectra working group for listening to and commenting on my work over the years.

Good luck to Gareth (keep on having it large!), Nick (don't get lost on your way around the World!), Paul 'Raging' Buck (you'll get there eventually!) and the Daves. Thanks for a truely messy few Lancaster years! God Bless, and get Jiggy with it ...

I also better mention the DESY boys, without which I may have come back to the UK a whole lot earlier and with money; 'Pies' Dave, 'Mike' Mike and 'and another thing' Jon.

Many thanks to my wife, Denise for putting up with me over the last few years and my frequent trips everywhere. Also, many thanks for reading this thesis and correcting my terrible English! Thanks also to my Mother and Father for your love and support, and to my Sister- best of luck for your upcoming degree course.

Lastly, but not leastly, I must thank the Particle Physics and Astronomy Research Council (PPARC) for their financial support.

Preface

The purpose of this thesis was to use the newly available (at the time of writing) calibrated dE/dx information to identify charged kaons, pions and protons passing through the central tracking region of the H1 detector in deep-inelastic ep scattering events. Through differentially plotting spectra of these particles, tests of QCD are carried out through the comparison with QCD Monte Carlo models.

The kaon and pion results allow comparisons of different QCD parton evolution models, as well as tests of the fragmentation process; in particular of the strangeness suppression factor as utilised in the JETSET fragmentation model. Through the addition of further cuts developed elsewhere, a search for QCD instanton induced events is also performed using these identified kaon and pion samples.

The proton selection allows the testing of baryon production in these interactions. In particular, it allows the possibility of looking at potential proton-antiproton asymmetries where none is naively expected, many units of rapidity away from the proton fragmentation region.

Chapter one looks at the physics underpinning the analysis. i.e. that of the deep-inelastic lepton-proton scattering process and the theory that describes it. It also outlines the mechanisms proposed to be responsible for the production of kaons, pions and protons in the hadronic final state. Chapter two outlines the workings of the H1 detector within the HERA collider and the various subdetectors necessary for this analysis. Chapter three describes the dE/dx technique and the calibration work that has been done to allow it to be used. Chapter four details the selection procedure for events and tracks used in the analysis, whilst chapter five gives the physics results and chapter six the instanton search results. Conclusions are presented in chapter 7.

Chapter 1

Physics at HERA

The experimental environment at H1 involves the collision of counter-rotating beams of electrons or positrons and protons. This analysis is concerned with the strong interaction of the beam constituents and the hadronic final state of those interactions.

The H1 detector at HERA is described in Chapter 2. In this chapter, the relevant theoretical framework for this analysis is discussed.

1.1 The Theory Of Deep Inelastic Scattering

The dominant process at H1 is that of photo-production [1], where the beam lepton radiates a quasi-real photon which then interacts with the proton. Within QCD framework, the photon couples either directly with a parton in the proton or indirectly via the photon's own parton content. The first type of interaction is known as direct photo-production (figure 1.1a), while the second is referred to as resolved photo-production (figure 1.1b).

The framework for this analysis, however, is provided by the theory of Neutral Current Deep Inelastic Electron Scattering (DIS)[1]. Detailed reviews of the

[1]Unless otherwise stated, the generic term electron is also taken to mean positron



Figure 1.1: *Photo-production in lepton-proton scattering, (a)direct and (b) resolved processes shown.*

theoretical aspects of DIS are available, in for example [2]. Here, aspects of the theory are discussed such that the measurements presented in this analysis can be understood.

1.1.1 DIS Kinematic Variables

The simplest tree-level diagram for electron-proton DIS is shown in figure 1.2. The kinematics of this process can be described by two independent variables for



Figure 1.2: *Basic Deep Inelastic Scattering Picture.*

a fixed centre of mass energy $\sqrt{s}$. These are chosen from; Q^2, the negative four-momentum transfer squared between the incoming and outgoing lepton, which is also the invariant squared mass of the virtual exchanged boson and gives the probing power of the photon on the proton; x_{Bj} (Bjorken x), which is a dimensionless variable associated with the fraction of the incident proton momentum carried off by the struck parton in the quark parton model (see later); y_{Bj} is the fraction of energy transferred from the incident electron to the hadronic system in the proton rest frame; both x_{Bj} and y_{Bj} are confined to the range 0-1.

From the four-vectors of the incoming electron (k) and the proton (P), the centre of mass energy squared can be obtained thus:

$$s = (P + k)^2 \tag{1.1}$$

The kinematic variables are thus defined as:

$$\begin{aligned} Q^2 &= -q^2 = -(k - k')^2 && (1.2) \\ x &= \frac{Q^2}{2P.q} && (1.3) \\ y &= \frac{P.q}{P.k} && (1.4) \end{aligned}$$

where, k is the 4-momentum of the incident lepton, k' is the 4-momentum of the scattered lepton, q is the 4-momentum of the exchanged boson and P is the 4-momentum of the incident proton.

The invariant mass squared of the outgoing hadronic system is given by:

$$W^2 = (q + P)^2 \tag{1.5}$$

Neglecting the lepton and nucleon masses, the following relations hold:

$$Q^2 \approx sxy, \qquad W^2 \approx Q^2 \left(\frac{1-x}{x}\right), \qquad s \approx 4E_e E_p \tag{1.6}$$

Experimentally, the kinematics can be determined by either the electron alone, or from the measured hadronic system alone, or from a combination of both. In this analysis, the electron method is used. It can be shown from the above equations:

$$y_e = 1 - \frac{E'_e}{E_e} sin^2 \frac{\theta_e}{2}, \qquad Q^2_e = 4E'_e E_e cos^2 \frac{\theta_e}{2} \tag{1.7}$$

where, E_e and E_e' are the energies of the incident and scattered lepton respectively and θ_e is the scattering angle with respect to the proton beam direction, x can then be obtained from 1.6.

1.1.2 The Deep Inelastic *ep* Scattering Cross-Section

The fundamental process of DIS as shown in figure 1.2 is:

$$e \ + \ P \rightarrow e' \ + \ X \tag{1.8}$$

where e and e' are the incoming and scattered electron respectively, P is the incoming proton and the hadronic final state resulting from the DIS process is represented by X.

Here, the scattering takes place via the exchange of either a virtual photon or Z^0 boson. This is known as a neutral current (NC) process. Charged current (CC) processes also occur, where the exchanged boson is a $W^{\pm}$, although it is only the NC processes that are considered in this analysis. The NC cross-section is:

$$\sigma_{NC} = \sigma(\gamma) + \sigma(Z^0) + \sigma(\gamma Z^0) \tag{1.9}$$

The Z^0 and γZ^0 interference term only becomes significant at high Q^2 values. At relatively low values of Q^2 (compared to the Z^0 mass squared) studied in this analysis, these terms are highly suppressed.

The differential cross-section can now be written in terms of the two independent structure functions $F_1(x, Q^2)$ and $F_2(x, Q^2)$ [2]:

$$\frac{d^2\sigma_{ep}^{NC}(x, Q^2)}{dxdQ^2} = \frac{4\pi\alpha^2}{xQ^2}\left[xy^2 F_1(x, Q^2) + (1 - y).F_2(x, Q^2)\right] \tag{1.10}$$

here α is the electromagnetic coupling constant.

The two structure functions F_1, F_2 are required for the independent contributions from the absorption of transversely (σ_T) and longitudinally (σ_L) polarised virtual photons. $F_1(x, Q^2)$ describes σ_T while $F_2(x, Q^2)$ describes the sum $\sigma_T + \sigma_L$.



Figure 1.3: *The quark-parton model.*

The following longitudinal structure function $F_L(x, Q^2)$ and the ratio R are also commonly used:

$$F_L = F_2 - 2xF_1, \qquad R = \frac{F_L}{F_2 - F_L} = \frac{\sigma_L}{\sigma_T} \tag{1.11}$$

1.1.3 The Quark-Parton Model

The first experiments investigating DIS found that the structure function $F_2(x, Q^2)$ was independent of Q^2 for fixed x values. This scaling behaviour is known as Bjorken scaling and was interpreted as evidence that the proton consisted of point like 'partons'. These partons were then taken to be equivalent to the quarks introduced by Gell-Mann to explain the structure of hadrons [3].

The quark-parton model (QPM) was introduced by Feynman [4] to interpret this scaling behaviour of the structure functions, and was based on the following assumptions:

1. The hadron involved in the DIS process is constructed of quarks travelling in the direction of the hadron, with the momentum shared amongst them.
2. The interaction of photon-hadron is thus expressed as the sum of incoherent scatterings from the point-like quark constituents (figure 1.3).

Important results that are obtained from the QPM are:

1. The QPM provides a simple relationship between the structure functions and the parton density functions $f_i(x)$ which can be understood as the probability of finding a quark of flavour i in the proton with momentum fraction given by x_{Bj}.

$$\frac{F_2(x)}{x} = 2.F_1(x) = \sum_i e_i^2 f_i(x) \qquad (1.12)$$

where e_i is the charge of quark flavour i.

2. The structure functions do not depend on Q^2, only on x, which was consistent with predictions made by Bjorken and by measurements taken at SLAC [5].

3. The Callen-Gross relation:

$$F_2(x) = 2xF_1(x) \qquad (1.13)$$

which is a consequence of the spin 1/2 nature of quarks and is only valid for the case where quarks are the only constituents of the proton. From 1.13, the results $R = 0$ and $F_L = 0$ are obtained.

More recent measurements in DIS covering larger ranges of Q^2 have shown that the QPM cannot be the whole story however. For example, if quarks and anti-quarks are the only constituents of the proton, then their momenta fractions should add up to unity, where in actuality:

$$\sum_q \int_0^1 x(q(x) + \bar{q}(x))dx \simeq 0.5 \qquad (1.14)$$

i.e. quarks only carry approximately half of the momentum of the proton as seen in DIS. The rest is carried by gluons in the framework of quantum chromodynamics.

1.1.4 DIS in Quantum Chromodynamics

The evidence for existence of other constituents of the proton apart from the quarks, as indicated by the deviation from unity in eq. 1.14 was also backed up by more recent measurements of F_2. With larger ranges of Q^2 available, it was soon seen that F_2 also depends upon Q^2 at smaller and larger values of x than previously reachable. This indicates that the scaling behaviour of the structure functions is violated (see figure 1.4).

The QPM is clearly not the whole truth, then, when dealing with DIS reactions involving the proton. Current models use the theory of quantum chromodynamics (QCD), which doesn't suffer from the above problems. The key aspects of this theory are outlined below:

- In addition to electric charge, quarks carry a colour charge (red, green, blue) which is the charge of the strong interaction.
- The gluon is the mediator of the strong force, of which there are eight bi-coloured varieties having spin 1 and zero electric charge.
- The quark-gluon colour interactions are computed by rules analogous to that of QED, with the substitution of $\sqrt{\alpha}$ for $\sqrt{\alpha_s}$ at each vertex and the introduction of colour factors.
- Unlike QED, the gauge bosons of this theory, the gluons, are able to interact with other gluons as they themselves carry colour charge.

Because of the self-coupling nature of the gluons in QCD, unlike QED, an anti-screening effect is present as a result of vacuum polarisation around a colour charge. This means that the closer one probes this charge, the less strong it appears to be.

When cross-sections are calculated within the QCD framework, integrals over the phase space of all quarks and gluons are found to diverge. A method to leave out these divergent parts of the integrals, called regularisation, is used. The re-calculated cross-sections become dependent on the energy scale μ_r^2 used in the regularisation. Renormalisation compensates for this unphysical effect by using an effective coupling constant and effective masses which absorb the divergent



Figure 1.4: *The Proton structure function $F_2^P(x, Q^2)$ from H1 and from fixed target experiments [6].*

contributions. The coupling constant α_s is then dependent on the scale μ_r^2 known as the renormalisation scale. In DIS, this scale is set by the virtuality of the photon Q^2, and to first order, the coupling constant can be written as:

$$\alpha_s(Q^2) = \frac{12\pi}{(33 - 2N_f)ln(Q^2/\Lambda^2)} \tag{1.15}$$

where, N_f is the number of quark flavours and Λ is a free parameter only determinable experimentally. Λ governs the dependence of α_s on energy and marks the scale that calculable, perturbative QCD breaks down. Or, put another way, it represents the boundary between dealing with quarks and gluons and the world of physically measurable hadrons.

1.1.5 The Proton Structure In QCD

Gluon emission from quarks in QCD extends the QPM picture of the proton which can be viewed as the lowest order QCD process. Unlike before in the QPM picture, with QCD, the fractional momentum sum (equation 1.14) no longer deviates from unity. The missing momentum is carried by the electrically uncharged gluons.

The violation of Bjorken-scaling at high and low x is explained in terms of the ability of gluons to split into $q\bar{q}$ pairs. More and more of these fluctuations can be resolved with increasing Q^2 (shorter wavelength of photonic probe). Hence, with increasing Q^2, a depletion of quarks at large x and a corresponding enhancement at smaller x is seen (figure 1.5). It is actually the gluon content that governs the proton at low x, giving rise to the DIS cross-section via $q\bar{q}$ pair creation.

1.1.6 QCD Evolution Equations

Extending the structure of the proton to include QCD effects does of course mean that the relation in equation 1.12 no longer holds.

A more general formulation of the structure functions using the idea of fac-



Figure 1.5: *Scaling violations: the photon can see more partons in (b) which carry a smaller fraction of the proton's momentum*

torisation to separate long and short range processes is:

$$F_2(x, Q^2) = \sum_{i=q,g} \int_x^1 d\xi C_i^V \left(\frac{x}{\xi}, \frac{Q^2}{\mu_r^2}, \mu_f^2, \alpha_s \right) .q_{i,h}(\xi, \mu_r^2, \alpha_s), \tag{1.16}$$

here, the sum encompasses all quark and gluon densities. C_i^V are the coefficient functions for each type of parton i and exchanged boson V (the virtual photon in this case), it does not depend upon the type of hadron involved. The factorisation scale μ_f^2 defines the point of separation between long and short range phenomena (figure 1.6). If the relevant scale (here, Q^2) is larger than μ_r^2, this process is calculable in perturbative QCD; whereas everything below this is absorbed into the parton densities $q_{i,h}$. These parton densities are specific to the hadron involved, h, but independent of the hard subprocess under consideration and are known as 'universal parton densities'.

Since these equations can, so far, not be solved exactly, several approximations of QCD are commonly used which are valid in different regions of phase-space. In the perturbative expansion, terms of the form $(\alpha_s ln(Q^2/Q_0^2))$, $(\alpha_s ln(1/x))$ and mixed terms $(\alpha_S ln(Q^2/Q_0^2)(ln(1/x)))$ appear. In the DGLAP



Figure 1.6: *(a) The ep cross-section factorised into lepton-parton cross-sections σ_{ei} and parton densities $q_{i,p}$. (b) Lowest order diagram contributing to σ_{ei} in (a)*

scheme, $(\alpha_s ln(1/x))$ terms are neglected, while the BFKL approach takes these terms to be dominant over the $(\alpha_s ln(Q^2/Q_0^2))$ terms used in the DGLAP approximation. The mixed terms start to become important as x decreases where Q^2 remains not too small.

1.1.7 DGLAP Evolution

In the region of large Q^2, the evolution of the quark and gluon densities is approximated by the DGLAP (Dokshitzer, Gribov, Lipatov, Altarelli, Parisi) evolution equations [7].

These are written in terms of quark and gluon splitting functions $P_{i,j}(z)$, which are calculable by perturbative expansion. These give the probability for the parton branchings $q \to qg$, $g \to q\bar{q}$, $g \to gg$, where the daughter parton i carries a fraction $(1 - z)$ of its mothers (j) momentum (see figure 1.7).

The DGLAP equations are obtained using the leading log approximation (LLA) where in leading order α_s, the dominant contributions are $(\alpha_s ln(Q^2/Q_0^2))^n$ terms summed for all orders of n. Dokshitzer [7] showed that this is equivalent to the sum of ladder graphs as shown in figure 1.8. Important consequences of



Figure 1.7: *The Feynman graphs for the splitting functions* (P_{qg}, P_{gq})*,* P_{qq} *and* P_{gg}

this scheme are, that in this gluon ladder:

- The transverse momenta $k_{T,i}^2$ of the emitted partons are strongly ordered such that, $k_{T,i}^2 \ll k_{T,i+1}^2 \ll \ldots \ll Q^2$.
- The longitudinal momenta x_i of the emitted partons satisfy $x_i \gg x_{i+1} \gg \ldots \gg x$.

The applicable region for the LLA is given by the use of perturbation theory ($\alpha_s(Q^2) \ll 1$) and by neglecting $ln(1/x)$ terms.

When moving towards smaller x, where contributions from $\alpha_s ln(1/x)$ and $\alpha_s ln(1/x) ln(Q^2/Q_0^2)$ become important, the LLA ceases to be valid. In this case, the Double LLA (DLLA) can be used. Here, in addition to the strong ordering in $k_{T,i}^2$, the longitudinal momenta of the emitted partons are also required to be strongly ordered: $x_i \gg x_{i+1} \gg \ldots \gg x$.

1.1.8 BFKL Evolution

In the region where x becomes small, but where Q^2 is not large enough for the DLLA to be valid, the DGLAP approximations cannot be used. Here, the BFKL (Balitsky, Fadin, Kuraev, Lipatov) equation [8] is used which sums diagrams leading in $ln(1/x)$ and independent of $ln(Q^2)$. With BFKL, the gluon ladder need not be ordered in k_T. The gluon distribution, then, is not integrated over



Figure 1.8: *Parton evolution in QCD (ladder diagram). x_i and $k_{T,i}$ represent the gluon longitudinal and transverse momenta*

k_T.

The different scope of the BFKL evolution leads to an ordering scheme for the parton emissions in LL(1/x) which is very different from the DGLAP type ordering:

- The transverse momenta $k_{T,i}^2$ of the emitted partons are unordered. The parton emissions perform a random walk in k_T^2 space.

- The longitudinal momenta x_i of the emitted partons are strongly ordered: $x_i \gg x_{i+1} \gg \ldots \gg x$.

The fact that the BFKL evolution is not ordered in k_T leads to a diffusion of k_T^2 which can extend into the infra-red region. This is usually avoided by introducing a cut-off parameter k_0^2, below which gluon emissions are not allowed. A side effect of this is that the results obtained show a dependence on k_0^2.

1.2 QCD Models

Model predictions of QCD DIS events have two parts to them: perturbative calculations of short range processes, and the long range non-perturbative effects of hadronisation. Perturbative QCD makes predictions for the **partonic** final state with fixed order (α_s) calculations of the matrix elements, or with parton showers, or a combination of both. Hadronisation turns the partonic final state into the observable particles we measure in the detector. For this, QCD models implement either the string fragmentation model or the cluster fragmentation model.

1.2.1 Matrix Elements

The tree level diagrams in figure 1.9(b) are the basic, first order α_s matrix elements, modelling the upper part of the DIS process in 1.9(a) and form the cornerstone of the QCD calculations shown in this thesis. Up to now, beyond this only NLO(α_s) (including virtual corrections from loop diagrams) calculations have been made available which, for example, have been able to predict dijet production to NLO(α_s). Although specialised calculations have been performed to NLO with regards to the hadronic final state, for example see [9], [10], such models are still not in a widely available format.

1.2.2 Leading Logarithmic Parton Showers

Higher order effects can be approximated using the leading log parton cascade idea [11]. Here, the quark struck by the virtual photon may emit partons both before and after this vertex corresponding to initial and final state parton showers (see figure 1.10). The initial state showering can be initiated by a parton close to its mass shell in the incoming proton. In each branching, one parton becomes increasingly space-like and the other is on-shell or has a time-like virtuality. The



Figure 1.9: *Tree level diagram for ep scattering (b) contributing to σ_{ei} in (a).*



Figure 1.10: *The (a) Leading log parton shower and (b) colour dipole model in DIS*

initial state cascade results in a space-like quark interacting with the virtual photon. The outgoing quark is then either time-like or on-shell. If the outgoing quark is time-like, the off-shell mass is reduced by subsequent branching into daughter particles until all partons are on-shell.

These cascade processes are based on the branching described by the DGLAP splitting functions in the LL(Q^2) approximation of QCD (see section 1.1.7).

1.2.3 The Colour Dipole Model

The Colour Dipole Model [12] (CDM) gives an alternative description of parton showers by describing the gluon bremsstrahlung in terms of radiation from colour dipoles between partons (figure 1.10(b)).

Initially, a colour dipole is formed between the struck quark and the proton remnant. Subsequent gluon emission is formed between the newly created colour charges and the original ones. This is iterated until the available energy is used up. An additional complication arises from the fact that the proton, unlike its dipole partner, the struck quark, is an extended object. The CDM introduces a suppression factor here due to the fact that emissions of small wavelengths from an extended antenna are suppressed.

In the CDM, the boson-gluon fusion process (BGF, see figure 1.9) is not described at all and has to be inserted by hand.

The final gluon configuration in the CDM is unordered in k_T.

1.2.4 String Fragmentation

In the confinement picture of quarks, as a $q\bar{q}$ pair separates, a colour field dipole is formed between them. As the $q\bar{q}$ separation increases, so does the strength of the colour field. The colour field can be considered to be string-like with a constant energy density [13]. When the energy stored in the string (which in DIS can also form between struck quark and the proton remnant) is large enough, the string breaks up creating a new $q\bar{q}$ pair which forms new string pieces. This process iterates until the available energy is used up, with resulting string fragments combined into mesons and baryons (see figure 1.11). Gluons are realised as kinks in the string which receive the four-momentum of the gluon.



Figure 1.11: *Fragmentation models: (a) the string and (b) the cluster model*

1.2.5 Cluster Fragmentation

Colour singlets are formed from the perturbatively generated partons in the cluster fragmentation model [52]. Initially, all gluons are non-perturbatively split into $q\bar{q}$ pairs and combined into low mass colour neutral clusters. These clusters are taken to be superpositions of meson resonances and decay into mesons or baryons according to the available phase space (see figure 1.11).

1.3 Strangeness Production In DIS

In the DIS picture shown in figure 1.8, strange particles may be produced in different ways. A strange quark can be scattered out of the proton sea, or $s\bar{s}$ pairs can be created; either in BGF processes (shown in top part of figure 1.8) or during parton showering and hadronisation. Strange particles can also arise from heavy quark decays, or as discussed later, from non-standard processes like instantons.

Most of the strange particles are expected from hadronisation with small fractional energy. In the Lund hadronisation model, $q\bar{q}$ pairs are created in the colour field of the string from the vacuum. Here, $s\bar{s}$ production is suppressed with respect to the other light quarks as a consequence of their relative masses ($m_s > m_u \approx m_d$). This is normally expressed as the ratio $\lambda_s : 1 : 1$, where λ_s is known as the strangeness suppression factor. The Lund default value for λ_s is 0.3 which represents the experimental mean from [14] and the typical value seen in electron-positron annihilation and moderate to large x DIS experiments [15], [16].

1.4 Protons in DIS

The production of baryons away from the proton remnant region in the DIS picture shown in figure 1.8 occurs during the fragmentation process. This is a poorly understood area however. The simplest mechanism used in the Lund string model allows the introduction of a diquark in a colour anti-triplet state that can be treated like an ordinary antiquark in the string. The string can break by quark-antiquark or antidiquark-diquark pair production. The diquarks can then combine with a quark to form a baryon. A more complex scenario is the 'popcorn' model [23], in which diquarks are not produced and baryons appear from the successive production of several $q\bar{q}$ pairs.

At HERA, a net baryon number is present due to the beam proton. In collisions, the proton is disrupted and the majority of the momentum is carried by diquarks giving rise to leading baryon production in the forward proton fragmentation region. In this picture, which describes the bulk of the data studied on leading baryon production, baryon number transfer over large rapidity intervals is strongly suppressed.

There are a number of measurements however [24], [25], [26], [27] which are difficult to explain by this diquark picture. There are also some recent results from photo-production studies at H1 [28] in which there is evidence of a baryon asymmetry in events with large charged particle multiplicities several units of rapidity away from the proton remnant.

A gluonic mechanism for the propagation of baryon number proposed some 25 years ago [29], [30] is discussed in a recent paper [31] with reference to the recent H1 preliminary results. The gluonic mechanism discussed in [31] successfully predicts the results shown in [28], and is predicted to be valid in the DIS regime also.

As mentioned, the dominant source of baryon production in the measurable region of phase space arises from the hadronisation of the DIS partonic state. DIS interactions at low values of x_{Bj} ($< 10^{-2}$; as in this analysis) are dominated by the gluonic content of the proton. This leads to an octet-anti-octet colour configuration of the partonic final state. Thus, the standard description in terms of a system of triplet strings is likely to be inadequate. In relation to this, baryon production could be expected to be heavily suppressed in hadronisation of an octet-anti-octet field [33].

1.5 Instantons at HERA

Strangeness production in DIS at HERA at levels above that predicted by the QCD picture discussed up to now is possible through the non-perturbative mechanism introduced by QCD instanton theory.

Instantons [17] introduce into the standard model processes which cannot be described by perturbation theory. In the case of electroweak interactions, baryon and lepton number is violated; and chirality is violated in the case of the strong interaction. As the name suggests, instantons are non-perturbative fluctuations confined to an 'instant' in space-time and have no corresponding free particle solutions for $t \rightarrow \infty$.

Unlike in electroweak theory where high energies ($O \gtrsim 10TeV$) are required, in QCD, instantons are expected to play a role at energies accessible by today's colliders. DIS at HERA is particularly interesting due to the hard scale for the instanton sub-process provided by the virtual photon probe Q^2. This is needed for theoretically sound predictions [18], [19], [20]. While instanton effects have not as yet been experimentally observed, their discovery would be of fundamental significance for particle physics.

1.5.1 Instanton Theory

Instantons originate from the topological structure of the vacuum in non-abelian gauge field theories. Figure 1.12 shows the structure of the vacuum, where neighbouring vacua have the same minimal potential energy and are separated by a potential barrier of height E_B. The vacua are labelled by their Chern Simons winding numbers N_{CS}, defined as an integral over the gauge fields A^a_μ with coupling g,



Figure 1.12: *The structure of the vacuum.*

$$N_{CS} = \frac{g^2}{16\pi^2} \int d^3x \epsilon_{ijk} (A^a_i \delta_j A^a_k - \frac{g}{3} \epsilon_{abc} A^a_i A^b_j A^c_k) \tag{1.17}$$

Transitions between neighbouring minima in figure 1.12 are possible classically when the energy E is large enough $E > E_B$, or by quantum mechanical tunnelling when $E < E_B$. The latter corresponds to the instanton solutions of the classical field equations. This instanton induced tunnelling process has a transition amplitude which is exponentially suppressed $\propto exp(-4\pi/\alpha)$.

The barrier height in electroweak theory is given by $E_B \approx m_w/\alpha_w = \mathcal{O}(10TeV)$. Instanton transitions between vacua ΔN_{CS} apart violate baryon (B) and lepton



Figure 1.13: *instantons in a) the electroweak and b) the strong interactions.*

(L) numbers according to:

$$\Delta(B + L) = -2n_{gener}\Delta N_{CS}, \tag{1.18}$$

with the following conditions:

$$\Delta(B - L) = 0 \quad \Delta L_e = \Delta L_\mu = \Delta L_\tau = \Delta B/3, \tag{1.19}$$

$n_{gener} = 3$ being the number of fermion generations. Figure 1.13(a) shows an electroweak interaction with $\Delta(B + L) = -6$.

This kind of interaction may have played an important part in the early universe when energies were typically this high. Importantly, in this respect, instantons in the electroweak sector may have played a part in the setting of the asymmetry between matter and anti-matter.

Chirality is violated with instanton induced QCD reactions. Where chirality (Q_5) is the difference between the number of left- and right-handed fermions, $Q_5 = \#L - \#R$. For n_f active quark flavours, the selection rule is:

$$\Delta Q_5 = 2n_f.\Delta N_{CS} \tag{1.20}$$

Here, the hard scale of the process, e.g. $E_B = \mathcal{O}(Q)$ for DIS, selects the barrier height. Also, the exponential suppression is less than in the electroweak case because $\alpha_s \gg \alpha_w$.

1.5.2 Instantons in DIS

Events due to instantons I (and anti-instantons $\bar{I}$) in DIS are predominantly produced in a photon-gluon fusion process [19], [20] (figure 1.14):



DIS variables:
$Q^2 := -q^2$
$x := Q^2 \,/(2P \cdot q)$
$W^2 := (q+P)^2 = Q^2\,(1-x)/x$
$\hat{s} := (q+g)^2$
$\xi = x(1+\hat{s}/Q^2)$

Variables of instanton subprocess:
$Q'^2 := -q'^2$
$x' := Q'^2 \,/(2\, g \cdot q')$
$s' := (q'+g)^2 = Q'^2\,(1-x')/x'$

Figure 1.14: *Kinematics of instanton induced processes in DIS*

$$\gamma^* + g \to^I \sum_{n_f}(\bar{q}_R + q_R) + n_g g \quad \gamma^* + g \to^{\bar{I}} \sum_{n_f}(\bar{q}_L + q_L) + n_g g \tag{1.21}$$

Quarks and anti-quarks of all n_f active flavours are found in each event as well as n_g gluons.

The kinematics of a DIS event with an instanton sub-process present is shown in figure 1.14. It can be seen that the instanton subprocess kinematics Q', x', s' are analogous to the standard DIS kinematic variables and are governed by the exchanged quark and gluon 4-momenta (q', g).

The instanton induced cross-section is given by a convolution of the probability to find a gluon in the proton, the probability that the virtual photon splits into a $q\bar{q}$ pair in the instanton background, and the cross-section of the instanton subprocess given by [20]:

$$\sigma^I_{q'g}(x', Q'^2) \approx \frac{\Sigma(x')}{Q'^2}\left(\frac{4\pi}{\alpha_s(\mu(Q'))}\right)^{\frac{21}{2}} .exp\left(\frac{-4\pi}{\alpha_s(\mu(Q'))}F(x')\right) \tag{1.22}$$

The cross-section depends critically on the functions $F(x')$ (called the 'holy grail' function), which modifies the exponent factor, and on $\Sigma(x')$, which depends on



Figure 1.15: *The instanton cross-section as a function of a) x' for $Q' = 5GeV$ and b) Q' for $x' = 0.2$*

$F(x')$. There is also a scale dependence due to the choice of renormalisation scale $\mu(Q')$. $F(x')$ can be estimated reasonably well for x' not too small ($x' \gtrsim 0.2$) [20], the theory contains ambiguities below this. The cross-section grows towards low x' and is peaked at $Q' = 5GeV$ (see figure 1.15). In the more theoretically safe region $x' > 0.2$, the expected fraction of instanton events in all DIS events is of $\mathcal{O}(10^{-3} - 10^{-4})$.

Although the cross-section is subject to some uncertainties, the properties of the final states of instanton induced events are reliably calculable. Whilst the property of non-conservation of chirality is difficult to exploit experimentally, further properties allow for possible discovery signatures. Striking features of the hadronic final state would be present due to the fact that, in the $q'g$ rest frame, $2n_f - 1$ quark and anti-quarks and n_g gluons are emitted isotropically from the instanton subprocess. n_g follows a Poisson distribution, with [21], [22]:

$$< n_g > \approx \frac{2\pi}{\alpha_s} x'(1-x') \frac{dF(x')}{dx'} \tag{1.23}$$

After hadronisation, this gives a high multiplicity spherical hadronic final state with the multiplicity depending on available centre of mass energy $\sqrt{s'}$. For a typical DIS kinematic event ($x' = 0.2$, $Q' = 5\ GeV \ \Rightarrow \ \sqrt{s'} = \ 10\ GeV$, $< n_g >= \mathcal{O}(2)$); about $n_p = 10$ partons and $n = 20$ hadrons would be expected.

Furthermore, with the decay into all kinematically allowed flavours of quark, another strong signature could be the excess of strange final state particles compared with the expected flux from DIS events.

Chapter 2

HERA and the H1 Detector

2.1 The HERA Collider

The **H**adron-**E**lectron-**R**ing-**A**ccelerator [34] at the **DESY** [35] facility in Hamburg is the first storage ring providing electron/positron-proton colliding beams. The two main accelerators providing these beams lie in a tunnel of circumference 6.3 km. They produce counter-rotating beams of 27.5 GeV (electrons) and 820 GeV (protons) giving a centre-of-mass energy of 300.3 GeV. The proton beam has now been upgraded to run at 920 GeV, an energy which it has run at since the start of the 1999 running period. During the 1996 and 1997 running periods considered in this analysis, positrons were used as the lepton beam.

HERA contains four interaction regions. H1 [36], situated in the North Hall and ZEUS [37] in the South Hall have been studying electron/positron-proton scattering events since 1992. HERA-B [38] (West Hall) uses tungsten wire targets inserted into the halo of the proton beam to study decay modes in the $B^0/\overline{B}^0$ system to investigate CP violation (this experiment started in 1999). Collisions of the positron beam operating in a polarised mode with a polarised gas target $(H_2, D,^3 He,^4 He)$ have been used by HERMES [39] (East Hall) since 1995 to study the origin of nucleon spin.



Figure 2.1: *The HERA collider.*

Figure 2.1 shows a schematic layout of the HERA accelerator complex. A chain of pre-accelerators (originally built for previous experiments) produces the electron/positron and proton beams at the required energy to inject into the main accelerators.

Using a 50 MeV linear accelerator, negatively charged hydrogen ions are stripped of their electrons and injected into DESY III. Here they are accelerated to 7.5 GeV before being transferred to PETRA. After a maximum of 70 bunches have been stored in PETRA, the protons are accelerated to 40 GeV and injected into HERA. Three iterations of this process are performed to complete a proton fill, leaving HERA with 210 proton bunches.

Electrons are accumulated and stored in a 60 mA single bunch from a 500 MeV linear accelerator. They are then fed into DESY II and accelerated to 7 GeV and transferred to PETRA (again with 70 bunches). PETRA accelerates the electrons to 11 GeV before injecting them into HERA. With the electrons, there are again 3 iterations of this procedure.

The proton beam is produced first in HERA and then ramped up to 820 GeV, then the electron beam is inserted and ramped to 27.5 GeV. Both beams are then optimised for maximum luminosity.

The electron beam was first replaced with positrons for the 1994 run. Positively charged ion clusters produced by getter pumps[1] had been found to reduce the lifetime of the beam. These ions are repelled by the positron beam, which therefore has a longer lifetime. Problems with the electron beam have been overcome lately, with electron beams included once more in the 1998 running period.

2.2 Overview of the H1 Detector

The H1 apparatus was constructed as a general purpose detector for ep collisions, based on a full coverage of the solid angle around the interaction point (with the exception of the beam holes).

From the isometric view of the H1 detector in figure 2.2, it can be seen that H1 is asymmetric in design. This is motivated by the asymmetric beam energies. The centre-of-mass frame is boosted with respect to the laboratory frame in the direction of the much more energetic proton beam (right to left in figure 2.2). As can be seen, the detector is considerably more massive and segmented in this region, where the highest particle and energy densities are expected.

H1 has the dimensions $12m \times 10m \times 15m$ approximately, and has a total weight of about 2800 tonnes.

The H1 detector is a group of devices providing complementary information on the various reaction products created at the interaction point. In the following sections, an overview is given of these components. A more detailed description is given in reference [36].

[1] Getter pumps maintain the high vacuum in the electron ring.