# A Search for Anomalous Top Production at H1

**Ronald Weber**

Diploma thesis

supervised by
Prof. Dr. R. A. Eichler

Tutor: Dr. C. Grab

Institute for Particle Physics
ETH - Zürich

Switzerland

September 13th 2001

*Diese Arbeit ist*
*meiner Grossmutter gewidmet*

# Contents


**1 Introduction** **3**

**2 An Introduction to the Theory** **5**
2.1 General Kinematics of *ep* Reactions . . . . . 5
2.2 Deep Inelastic *ep*-scattering in the Parton Model . . . . . 7
2.3 Photoproduction . . . . . 8
2.4 Anomalous Top Production and its decay modes . . . . . 9
2.5 *ep* collisions at HERA . . . . . 11

**3 Experimental Setup** **13**
3.1 The HERA Collider . . . . . 13
3.2 H1 Detector . . . . . 14
3.2.1 Tracking Chamber . . . . . 16
3.2.2 Liquid Argon Calorimeter (LAr) . . . . . 18
3.2.3 Superconducting Coil and Iron Yoke . . . . . 20
3.2.4 Trigger System and Reconstruction . . . . . 20
3.3 Luminosity System . . . . . 22

**4 Monte Carlo and Background** **23**
4.1 Anomalous Top Production Monte Carlo . . . . . 23
4.2 Background . . . . . 25
4.2.1 Non *ep* Background . . . . . 25
4.2.2 SM $W$ Production ($\gamma p$) . . . . . 25
4.2.3 Lepton Pair Production ($\gamma\gamma$) . . . . . 26
4.2.4 Standard Model Jets ($\gamma p$) and Single $\pi$-Jets . . . . . 27
4.2.5 Anomolous top decay in muons . . . . . 27

**5 Event Selection** **29**
5.1 Principle of the analysis . . . . . 29
5.2 Reconstruction of the Event . . . . . 30
5.3 The Jet Algorithm *QJCONE* . . . . . 31
5.3.1 Definition of the Jet . . . . . 31
5.3.2 Reconstruction of the $\tau$ Jet . . . . . 33
5.4 Sensitive kinematical variables for Anomalous Top Production . . . . . 33

5.5 The Preselection . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . 36
5.6 Final Analysis . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . 37
5.7 Calculating the Efficiency . . . . . . . . . . . . . . . . . . . . . . . . . . . . . 45
5.8 Result . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . 46

**6 The Cross Section for Anomalous Top Production 47**
6.1 Production Cross Section . . . . . . . . . . . . . . . . . . . . . . . . . . . . . 47
6.2 Upper Limit . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . 48
6.2.1 Background Estimation . . . . . . . . . . . . . . . . . . . . . . . . . . 49

**7 Conclusion and Outlook 51**

**8 Acknowledgments 53**

**A Event Display 55**

**Abstract**

A search for Anomalous Top Production mediated by a flavour changing neutral current was performed in $ep$ collisions at HERA. The search examined data collected with the H1 detector from 1999 and 2000 with an integrated luminosity of 62.76 pb$^{-1}$. The analysis was motivated by the previous observation of outstanding events with isolated leptons, missing transverse momentum and a jet with large transverse momentum. The top decay is considered for the first time in the decay chain $t \to Wb$, $W \to \tau\nu_\tau$ and a hadronic final state. Within this decay channel no events were found to be compatible with the hypothesis of Anomalous Top Production. The resulting upper limit for the production cross section of Anomalous Top Production was calculated to be

$$\sigma(ep \to e'tX, \sqrt{s} = 320 \text{ GeV}) \leq 6.8 \ pb \qquad \text{at } 95 \ \% \text{ CL}$$

and supports the upper limit derived from the previous Anomalous Top Production searches at HERA in the electron and muon channel.

# Chapter 1

# Introduction

In todays High Energy Physics, the *Standard Model* (SM) is well accepted and has shown its validity up to $\sim$1000 GeV. But what happens beyond this energy, is subject to many new experiments and collaborations, such as LHC[1] at CERN[2] or TESLA[3] at DESY[4]. Even though the SM has helped to understand many things in modern particle physics, the fundamental question about the origin of matter and especially of the masses is still unanswered. The SM is not able to answer this question satisfyingly, therefore the origin of the masses is to be searched in physics *Beyond the Standard Model* (BSM).

One field of research is the *single top quark physics*. It started more than 11 years ago with the discovery of the heaviest quark, the top quark ($t$), at the Fermilab Tevatron. The fact that the mass of the $t$ quark ($\sim$175 GeV) is of the same order of magnitude as the Electroweak Symmetry Breaking scale (EWSB, $\sim$250 GeV) makes the top a very interesting object of studies. With the EWSB one expects also new physics to show up, and the top is certainly the most promising candidate for searches toward that direction. The variety of these possible physics Beyond the Standard Model is widely spread: anomalous quark couplings [1], anomalous $Wtb$ couplings [2], new strong dynamics [3], flavour changing neutral currents (FCNC) [4], R-Parity violating SUSY[5] effects [5], CP-violation effects [6], and so on. But all models agree, that the effects should be first visible in the top sector.

The FCNC is the origin for the *Anomalous Top Production* studied here. The FCNC allows a light quark to interact with a neutral gauge boson to form a top quark. Since the SM preserves all flavours in neutral current interactions in leading order, this process is not allowed within the SM.

[1]**L**arge **H**adron **C**ollider
[2]**C**onseil **E**uropéen pour la **R**echerche **N**ucléaire
[3]**T**era electron volt **E**nergy **S**uperconducting **L**inear **A**ccelerator
[4]**D**eutsches **E**lektronen-**SY**nchrotron
[5]**SU**per**SY**mmetry

At HERA[6], an exciting experimental fact is the observation of isolated electrons and muons with high transverse momentum along with missing energy and a large hadronic final state [7]. The topology and the kinematics of these events are of a kind that cannot be explained within the SM, but they would fit quite well to the Anomalous Top Production . The present analysis was looking for Anomalous Top Production in the $\tau$-channel, using the data from 1999 - 2000 of H1 with an integrated luminosity of $\mathcal{L} = 62.76\ \mathrm{pb}^{-1}$.

[6]**H**adron **E**lektron **R**ing**A**nlage

# Chapter 2

# An Introduction to the Theory

This chapter will give a short introduction to the theoretical aspects encountered in this analysis. The kinematics, the parton model as well as the Anomalous Top Production will be briefly explained. The SM predicts quark flavour conservation in all strong and electromagnetic interactions, therefore the $t$ can only be produced in pairs ($t\bar{t}$). The FCNC 'converts' a lighter quark to a single, or anomalous, $t$ and must be a non-SM process. The decay of the anomalous $t$ is considered to be SM like.

## 2.1 General Kinematics of $ep$ Reactions

The interaction between electrons and protons ($ep$)

$$e + p \quad \rightarrow \quad l + X \tag{2.1}$$

is mediated by a gauge boson. The interaction forces are either electromagnetic ($\rightarrow \gamma$) or weak ($\rightarrow W$, $Z^0$). The gauge boson transfers the (momentum)$^2$

$$-Q^2 = q^2 = (k - k')^2, \tag{2.2}$$

where $k$ and $k'$ represent the four-vectors $(E, \vec{p})$ of the incoming and outgoing lepton, respectively. Fig. (2.1) shows the corresponding Feynman graph. The outgoing lepton is either an electron (Neutral Current, NC) or a neutrino (Charged Current, CC), and $X$ represents the *Hadronic Final State* (everything else but the lepton).

For $Q^2 \approx 0$ GeV$^2$ the photons are real, this process is called *Photoproduction* opposed to the *Deep Inelastic Scattering* (DIS) with virtual photons for large $Q^2$.
In this analysis the entire kinematical $Q^2$-range was covered.
The two incoming 4-vectors define also the *center of mass energy*

$$\sqrt{s} = \sqrt{(P + k)^2} \overset{*}{\sim} \sqrt{4 E_p E_e} \tag{2.3}$$

The last relation ($*$) is valid for storage rings neglecting the masses of the beam particles, electron and proton at HERA. The kinematics of the scattering can be expressed by $s$ and two further independent variables. One possible choice is




Figure 2.1: The general Feynman graph for *ep* interactions. The values in brackets stand for the 4-vectors, $l$ is the scattered lepton and $X$ the hadronic final state.

$$x = -\frac{q^2}{2(q \cdot P)} \quad \text{and} \quad y = \frac{q \cdot P}{k \cdot P} \tag{2.4}$$

These variables have the advantage of being dimensionless. $x$ is the *Bjorken variable* and $y$ the momentum transfer by the photon (in proton rest frame). The relation to describe the kinematics is now

$$Q^2 = x \cdot y \cdot s \tag{2.5}$$

Besides $x$ and $y$, one can also choose $Q^2$ as independent variable. Historically the second parameter would then be

$$\nu = \frac{P \cdot q}{m_p} \tag{2.6}$$

$Q^2$ is also a measure for the virtuality of the process ('Is the photon real or virtual?') and $\nu$ is the energy of the gauge boson in the proton rest fame.

In the early 1970's, Richard Feynman proposed the *Parton Model* [8], describing the basic constituents of the proton. It explained the phenomenon of the *scaling* [9] described by J. D. Bjorken in 1967.

## 2.2 Deep Inelastic *ep*-scattering in the Parton Model

Gell-Mann [10] and Zweig [11] introduced the *Quarks* as constituents of the proton in analogy to the nucleons in the nucleus. The quarks were later recognized as one kind of Feynman's partons, the other kind were the *Gluons*. Each kind of particles carries about 50 % of the proton momentum. The quarks inside the proton are two $u$ 's and one $d$ (valence quarks), and virtual $q\bar{q}$-pairs (sea quarks).
The parton model interprets the deep inelastic *ep*-scattering as an elastic scattering of the electron off the partons [12]. The electron has to interact by electromagnetic or weak forces with the partons, exchanging a photon or a $Z^0$-boson (NC) or a $W$-boson (CC).
The Bjorken variable gets a very descriptive meaning in the infinite momentum frame[1]: It represents the fraction of momentum carried by the parton (see Fig. (2.2)).




Figure 2.2: Deep inelastic $ep-$scattering as described by the parton model, in brackets the 4-vectors. $X$ is the hadronic final state. The momentum carried by the scattering parton is $xP$, where $x$ is the Bjorken variable.

Most partons coming from the DIS will hadronize and finally end up as hadrons, a process called *fragmentation*. This fragmentation is not yet fully modeled. It may be described by fragmentation functions of heavy quarks. The most used functional form was suggested by Peterson *et al.* [13]. A heavy parton may either hadronize (e.g.[2] $b$) or decay in a $W$ (e.g. $t$) and decay further to leptons and quarks, where the quarks will undergo fragmentation to form hadrons. These hadrons may decay as well (e.g. $\pi^0 \to \gamma\gamma$). All undecayed hadrons together build the *Hadronic Final State X*.
These hadrons may either be isolated or collimated in a narrow group. The collimated hadrons form a so called *Jet*. Jets are mostly the remnants of a single parton (quark or gluon). The reconstruction from the hadrons to the parton is due to the open question during fragmentation very challenging.

[1]A reference frame where the proton carries a large momentum and the transverse momentum of the partons inside the proton is neglectable. In this frame these partons are not interacting with themselves. The laboratory frame at HERA is a valid approximation of the infinite momentum frame.

[2]exempli gratia

## 2.3 Photoproduction

For low $Q^2$ the electron scatters off the proton at small angles, exchanging a (quasi)-real-photon. The photoproduction describes a photon-proton scattering process with the electron as photon source.

The hadronic structure of the photon is the origin for several subprocesses contributing to the total photoproduction cross section. These subprocesses can be classified in various ways ([14] and references therein). A photon may split in a $q\bar{q}$-pair (anomalous component) or transform into a vector meson (VDM). These photons are also called *resolved photons*, whereas the photons interacting as point-like gauge bosons (BGF, QCD-Compton) are called *direct photons*. The Feynman graphs for leading order are given in (Fig. (2.3)). This distinction is lost in next to leading order, since higher order effects mix the different processes up.




(a) VDM-type resolved




(b) anomalous resolved




(c) photon-gluon fusion (BGF)




(d) QCD-Compton process

Figure 2.3: Leading order Feynman graph for photoproduction processes. The corresponding process is given below the figure. The V denotes a vector meson ($\rho, \omega, \phi, J/\psi, \Upsilon$) and $\mathbb{P}$ the pomeron.

## 2.4 Anomalous Top Production and its decay modes

The Anomalous Top Production arises from a *Flavour Changing Neutral Current*. The possible gauge bosons are the $\gamma$ and the $Z^0$. For $ep$ collisions the t-channel exchange of a $\gamma$ is much more likely than the $Z^0$-exchange [15] since the massive $Z^0$ is highly suppressed in the propagator compared to the massless $\gamma$. Due to the larger parton density in the proton, the coupling to the $u$-quark is in favor compared to the $c$-quark coupling (c-quarks are only expected as sea quarks in the proton[3]). For the coupling of the top to the light quarks ($u$,$c$) only upper limits from radiative top decays set by the CDF Collaboration [17] are available: $\kappa_{\gamma,u} < 0.28$ [18].

Fig. (2.4) shows the Feynman graphs for the Anomalous Top Production for $ep$ collisions.




(a) Anomalous Top Production in $ep$ collisions with $\gamma$-exchange ...

(b) ... and $Z^0$-exchange

Figure 2.4: The Feynman graphs for the Anomalous Top Production in leading order for $ep$ collisions. Only the scattering quark from the proton is drawn. Fig. (a) is the relevant process at HERA.

Considering only SM-decays, the top is assumed to decay only into $W$ and $b$. The b-quark will cause a high energy hadronic jet, whereas the $W$ has several decay channels. The leptonic decays into electron and muon were previously used for searches for Anomalous Top Production at HERA [15]. In this new approach, the third leptonic decay ($W \to \tau\nu_\tau$) was examined. The corresponding Feynman graph is drawn in Fig.(2.5). Since the electromagnetic charge is to be conserved in all interactions, the decay of the top $t \to Wb$ produces only positively charged $W$ bosons and therefore only positively charged $\tau$ and $\pi$ are produced in the top decay chain.

[3] There are hints for an 'intrinsic charm distribution' [16]




Figure 2.5: The top decay in the $\tau$-channel with missing energy ($\nu_\tau$) and a high energy jet ($b$). The decay $t \to Wb$ separates the space in two hemispheres.

To isolate the Anomalous Top Production, the decay products of the top[4] have to be identified in the detector. The tau lepton decays not only leptonically but also hadronically. The dominant decay is a so-called *hadronic one prong decay*: $\tau^+ \to h^+ + n$ neutrals with a *Branching Ratio* (BR) of 46.7 % [19]. The $h^+$ stands for a pion or a kaon, where as the pion is much more likely than the kaon. Possible neutral particles are $\pi^0$ and neutrinos. Within the one-prong decay the process $\tau \to \pi^+ + n\,\pi^0 + \nu_\tau$ is dominating with a BR of 36 % for $1 \leq n \leq 3$. The main task was to identify the charged pion in the detector. Each neutral pion will decay almost immediately into two photons. The characteristics of the $\tau$ decay are the missing energy from the neutrino and one charged track with additional energy in the calorimeter from the $\gamma$'s. The $\tau$-decay is drawn in Fig.(2.6).




Figure 2.6: The $\tau$-decay: $\tau \to \pi^+\pi^0\nu_\tau$ and $\pi^0 \to \gamma\gamma$.

[4]The expression 'top' refers to the anomalous top ($t$) as produced in Fig. (2.4(a))

Table (2.1) gives a short summary on the most important particles associated with the decay of the top.

| **Particles associated with the Top in this Analysis** | | | | |
|---|---|---|---|---|
| Particle | Particle Type | mass [GeV] | Mean Life Time [s] | Considered Decay Mode (BR) |
| top ($t$) | quark | 174.3 | — | $Wb$ (100 %) |
| $W$ | gauge boson | 80.4 | $\Gamma = 2.12 \pm 0.05$ GeV | $\tau\nu_\tau$ (10.4 %) |
| Tau ($\tau$) | lepton | 1.78 | $290.6 \times 10^{-15}$ | hadronic one-prong (49.5 %) |
| Pion $\pi^+$ | meson | 0.139 | $2.6 \times 10^{-8}$ | no decay |
| Pion $\pi^0$ | meson | 0.135 | $8.4 \times 10^{-17}$ | $2\gamma$ (98.8 %) |

Table 2.1: Summary on all particles associated with the top encountered in this analysis

## 2.5 *ep* collisions at HERA

This section will give a summary of the possibilities at HERA. A more detailed description of the experimental setup is given in chapter 3.
The dominating process at HERA is the photoproduction. With a center of mass energy of roughly 300 GeV (see chapter 3.1) no $t\bar{t}$-pairs can be produced. So any reported top quarks at HERA would imply a single top physics process.

H. Fritzsch examined the theoretical possibilities of detecting single tops at HERA [20]. With an expected integrated luminosity of about 100 pb$^{-1}$and favorable circumstances, he predicted the possibilities to find 'a few' events of this type. The isolated leptons reported from H1 Collaboration strongly support the existence of the single top's.

# Chapter 3

# Experimental Setup

## 3.1 The HERA Collider




Figure 3.1: The HERA collider with the storage ring and the preaccelerators. The experiment H1 is situated in the northern hall.

At the DESY laboratories in Hamburg, Germany, is the HERA (**H**adron **E**lectron **R**ing**A**nlage). HERA accelerates and stores electrons (or positrons[1]) and protons and provides the unique opportunity to study lepton - quark interactions. The two particle types are gathered in bunches with up to $10^{11}$ single particles per bunch. During normal operation roughly 220 bunches in 96 $ns$ time intervals circulate in the storage rings representing the electron and proton beams. The two storage rings are 6.3 km long and are roughly 20 m below the surface. For the proton ring superconducting magnets have to be used to bend the proton beam to a stable trajectory.

[1]most runs at H1 were made using positrons ($e^+$) instead of electrons ($e^-$), from now on 'electron' refers to both particles.

Since 1992 the electrons and protons are preaccelerated with various linear accelerators (LINAC) and two former storage rings (DESY, PETRA) and then fed to the large HERA rings where the particles will reach their final energies of 27.6 GeV and 820 GeV, respectively, leading to a center of mass energy of $\sqrt{s} = 301$ GeV in the collision (see equation 2.3). Since 1999, the protons are even accelerated to 920 GeV pushing the center of mass energy to $\sqrt{s} = 319$ GeV. An overview of the preaccelerators and the HERA rings is given in Fig. (3.1).

Along the storage rings of HERA are 4 large experimental halls. In the northern hall is the experiment H1, an international collaboration of about 400 scientists from 39 institutes of 12 countries throughout the world. Here the electron and proton beams are collided head-on. The resulting particles of the collisions are detected with the general purpose H1 detector. The main interest of research of the H1 collaboration is to measure the structure of the proton, to study the fundamental interactions between particles, and to search for physics Beyond the Standard Model of the elementary particles. The three other halls are occupied by the experiments HERMES, ZEUS and HERA-B.

## 3.2 H1 Detector

Unlike most other High Energy Physics Detectors, the H1 Detector is not symmetric with respect to the nominal interaction point[2]; this is due to the different energies of the two beam particles. The proton has more energy (momentum) than the electron and the greater momentum along the beam axis causes the main event to be shifted in flight direction of the proton beam. This also defines the *reference frame* at H1: The positive z axis is along the proton beam and is called the forward direction, perpendicular to it is the $xy$ plane (where the x-axis points toward the center of the ring and the y-axis points upward). It is also referred to as the *transverse plane*. The origin of the coordinate system is defined as the nominal interaction point.
There are further two angles of importance: $\theta$ and $\phi$: $\theta$ is the *polar angle* between the trajectory and the z-axis (scattering angle), $\phi$ the *azimuthal angle* in the transverse plane with $\phi = 0$ corresponding to the positive x-axis. For lorentz invariance one introduced the variable *rapidity* $\hat{y}$. In most cases the relevant parameter is the *pseudo-rapidity* $\eta$ which is an approximation of $\hat{y}$, neglecting the masses of the particles ($m = 0$). The pseudo-rapidity is correlated with the polar angle $\theta$

$$\eta = -\ln\left(\tan\frac{\theta}{2}\right) \tag{3.1}$$

$\eta$ and $\phi$ assure a set of coordinates, where the difference in each coordinate is lorentz invariant.

Fig. (3.2) shows the H1 detector with the reference frame and the asymmetric structure.

[2]The nominal interaction point is the point where the electron and proton beams collide.