Departement Natuurkunde
Faculteit van de wetenschappen
Vrije Universiteit Brussel

Triple Regge analysis of single diffractive photoproduction at HERA

Roel Heremans
december 2002

Triple Regge analysis of single diffractive photoproduction at HERA

PROEFSCHRIFT

ingediend met het oog op
het behalen van de wettelijke graad
van doctor in de wetenschappen
aan de Vrije Universiteit Brussel

door

Roel Heremans

Promotiecommissie:

Promotor:	Prof. Dr. R. Roosen, Vrije Universiteit Brussel
Voorzitter:	Prof. Dr. J. Van Craen, Vrije Universiteit Brussel
Secretaris:	Prof. Dr. R. Roosen, Vrije Universiteit Brussel
Overige leden:	Prof. Dr. E. De Wolf, Universiteit Antwerpen
	Prof. Dr. J. Lemonne, Vrije Universiteit Brussel
	Prof. Dr. P. Marage, Université Libre de Bruxelles
	Prof. Dr. A. Sevrin, Vrije Universiteit Brussel
	Prof. Dr. T. Sloan, Lancaster University

Cover: HERA, the Hadron Electron Ring Accelerator at the DESY laboratory in Hamburg. (Photo: David Parker/SPL taken from *The Particle Odyssey: A Journey to the Heart of Matter*, F. Close, M. Marten, C. Sutton, *Oxford University Press* (2002))

Dedicated to the memory of my father.

Abstract

A new measurement is presented, using the H1 detector, of the differential cross section $M_X^2\ d\sigma/dM_X^2$ for the process $\gamma p \to XY$, with a large rapidity gap between the systems X and Y and with Y a proton or a low mass proton excitation, at three centre of mass energies $\langle W \rangle = 91$, 187 and 231 GeV. The slope parameter $\alpha_{eff}(0)$ obtained from a single Regge type parameterization with a single effective trajectory is presented as a function of M_X^2. A combined triple Regge fit is performed over these data together with leading proton data from the same experiment and lower energy fixed target data. The pomeron intercept $\alpha_{I\!P}(0)$ is extracted from this fit and was found to be $\alpha_{I\!P}(0) = 1.127 \pm 0.004(\mathrm{stat}) \pm 0.025(\mathrm{syst}) \pm 0.046(\mathrm{mod})$. The extracted triple Regge couplings are compared with those obtained from pp-data and found to be in reasonable agreement. Diffractive data like $F_2^{D(3)}$, F_2^{LP} and F_2^{LN} were compared in the deep inelastic region with a triple Regge model, using either the triple Regge couplings extracted from the combined triple Regge fit or using the triple Regge couplings which were determined from pp data. A reasonable description of the data at low Q^2 and high M_X is found for both methods.

Contents

List of Figures

List of Tables

Chapter 1

Introduction

The data analysed in this thesis has been collected at HERA, which is the world's first electron-proton collider situated near to Hamburg, Germany. The large centre of mass energy ($\sqrt{s} \approx 300$ GeV) obtained by colliding electrons of 27.5 GeV on 820 GeV protons, allows to explore new regions, where particles with masses up to 300 GeV can be produced, where the structure of the proton can be studied with a resolving power varying over 5 orders of magnitude down to dimensions of 10^{-18} m and where partons with very small fractional proton momenta (Bjorken-x down to 10^{-6}) become experimentally accessible.

The H1 and ZEUS experiments, both consisting of multipurpose detectors, installed respectively around the northern and southern interaction point of HERA, observed in 1993 a large number of events in which in a large interval in rapidity no final state particles were observed. These events, nowadays known as rapidity gap events, are interpreted as diffractive events resulting from the exchange of a colorless object called the *pomeron*. Since their observation, rapidity gap events have been extensively studied. In particular, because the large Q^2 photons available at HERA allow to investigate the internal structure of the pomeron.

In the present analysis, rapidity gap events are studied using the process $ep \to Xp$, where X represents the photon dissociation system. The process will be analysed in the photoproduction limit where the photon's momentum transfer $Q^2 \to 0$, i.e. the photon is quasi-real. In particular the differential cross section $M_X^2\ d\sigma/dM_X^2$ is measured and subjected to a triple Regge analysis in which also data from fixed target experiments and leading proton data are considered. The results of this analysis are then also extrapolated to the deep inelastic region to investigate where the transition between the soft and hard processes takes place.

The structure of the thesis is as follows: Chapter 2 introduces the main feature of lepton-nucleon scattering going over the deep inelastic diffractive scattering, towards the topic of this thesis, diffractive scattering in photoproduction. Here the triple Regge theory is introduced and a summary of existing analysis in this domain is reviewed stressing those measurements which will be used in either the triple Regge analysis or in comparisons made in chapter 6. Chapter 3 reviews the main layout of the HERA accelerator and introduces the pilot bunch concept. Also the main H1 detector components are briefly described, with emphasis on those which are of particular importance in this analysis. Chapter 4 treats the subject of background identification, data selection and mass reconstruction. Chapter 5 describes the extraction of the differential cross section, $M_X^2\, d\sigma(\gamma p \to XY)/dM_X^2$. In chapter 6, the results obtained in chapter 5 together with earlier diffractive data are subjected to a triple Regge analysis. These results are then extrapolated to the deep inelastic regime.

Chapter 2

Theoretical introduction

The HERA physics program to date has primarily focused on testing our understanding of the strong force. Measurements have been performed in kinematic regions where perturbative QCD calculations are accurate, as well as in regions where no hard scale is present and non-perturbative processes dominate. In the following section, the structure of the proton and the measurement of the proton structure function will be briefly reviewed. Subsequently the concept of structure functions in deep inelastic diffractive scattering will be schematically outlined. Then the framework in which soft diffractive processes are considered, i.e. the triple Regge formalism, constituting the theoretical framework in the analysis of this thesis, will be treated in detail. As the roots of this theory date back to the late sixties, the triple Regge formalism has been applied to several diffractive final states in γp- and pp-experiments in the past. These results will be reviewed at the end of this section with emphasis on those obtained at the HERA experiment.

2.1 Lepton-nucleon scattering

In the most general case the lepton-nucleon interaction proceeds via the exchange of a virtual vector boson as depicted in figure 2.1. Since the lepton number has to be conserved, we expect the presence of a scattered lepton in the final state, while the nucleon fragments into a hadronic final state X. The interaction is called inelastic when a quark is knocked out of the proton, and the proton is broken up. It is called deep when the proton is probed with a gauge boson which resolves small distance scales.

The following four Lorentz invariant variables are most commonly used to describe the lepton nucleon scattering:

$$\begin{aligned}
s &= (k+P)^2, \\
Q^2 &= -q^2 = -(k-k')^2, \\
x &= \frac{Q^2}{2P.q}, \\
y &= \frac{q.P}{k.P}.
\end{aligned}$$



Figure 2.1: *Schematic diagram describing inelastic lepton-nucleon scattering. The four-vectors of the particles, or system of particles, are given in parentheses.*

where the four-vectors of the initial and final lepton, of the incoming nucleon, and of the outgoing hadronic system, are represented by k, k', P and P' respectively. The variable s is the centre of mass energy squared with $\sqrt{s} \approx 300$ GeV at HERA[1]. Because of the large value of s at HERA, photons with a virtuality $Q^2 \approx 10^4$ GeV^2 can be produced, resolving sizes as small as $\sim 10^{-18}$ m. In the proton rest frame the expression for y reduces to

$$y = 1 - \frac{E_{l'}}{E_l}, \tag{2.1}$$

where $E_{l'}$, E_l are the energies of the scattered lepton and incoming lepton, respectively. In this frame the dimensionless variable y is a measure of the fraction of the incident electron energy which enters the interaction. In the parton model the Bjorken variable x can be interpreted as the fraction of the proton momentum carried by the massless struck quark. At fixed centre of mass energy $\sqrt{s}$, only two of these four variables (x, y, s, Q^2) are independent due to energy momentum conservation. When the electron and proton are assumed to be massless the relation between them is:

$$Q^2 = xys. \tag{2.2}$$

In addition W is defined as the centre of mass energy of the exchanged boson and the incoming proton

$$\begin{aligned} W^2 = (q+P)^2 &\simeq ys - Q^2 \\ &\simeq Q^2\left(\frac{1}{x} - 1\right) \end{aligned} \tag{2.3}$$

[1]This is for 27.5 GeV incident electrons or protons and 820 GeV protons.

2.2 Proton structure function

Most generally, the cross section of unpolarized ep-scattering is written as a tensor product made up of a leptonic and a hadronic part, $\sigma \sim L_{\mu\nu}W^{\mu\nu}$, where $W^{\mu\nu}$ represents the hadronic tensor, which parameterizes the interaction occurring between the exchanged boson and the proton. Consideration of the requirements of Lorentz invariance and current conservation lead to the following expression for the deep inelastic neutral current cross section for not too high Q^2 [1, 2]:

$$\frac{d^2\sigma(ep \to eX)}{dxdQ^2} = \frac{4\pi\alpha_{em}^2}{xQ^4}\left[(1-y+\frac{y^2}{2})F_2(x,Q^2) - \frac{y^2}{2}F_L(x,Q^2)\right] \tag{2.4}$$

$$= \frac{4\pi\alpha_{em}^2}{xQ^4}\left[(1-y+\frac{y^2}{2})\ \sigma_r\right] \tag{2.5}$$

where α_{em} is the electromagnetic coupling constant and $F_2(x,Q^2)$ and $F_L(x,Q^2)$ are structure functions which provides information on the quark and gluon content of the proton and $\sigma_r = F_2 - \frac{y^2}{1+(1-y)^2}F_L$. In order to describe fully the neutral current DIS at high Q^2, a third structure function, $F_3(x,Q^2)$ is required, which in addition to the photon also takes into account the parity violating Z^0 exchange. In the region of low Q^2 ($Q^2 \ll M_{Z^0}^2$) only photon exchange contributes. The Z^0 exchange and the interference between the Z^0 and the photon exchange can be neglected safely since the gauge boson masses enter the matrix element of the cross section by the propagator terms $1/Q^2$ and $1/(Q^2+M_{Z^0}^2)$ for the photon and Z^0 respectively.

The two structure functions $F_2(x,Q^2)$ and $F_L(x,Q^2)$ can also be expressed as a function of two independent contributions, σ_T and σ_L of the photo-absorption cross section [3]:

$$F_2(x,Q^2) = \frac{Q^2}{4\pi^2\alpha_{em}}\left[\sigma_T(x,Q^2) + \sigma_L(x,Q^2)\right], \tag{2.6}$$

and

$$F_L(x,Q^2) = \frac{Q^2}{4\pi^2\alpha_{em}}\sigma_L(x,Q^2). \tag{2.7}$$

The quantity $R(x,Q^2)$ can be defined as the ratio of longitudinally to transversely polarized photon cross sections and is given by,

$$R(x,Q^2) = \frac{\sigma_L(x,Q^2)}{\sigma_T(x,Q^2)} = \frac{F_L(x,Q^2)}{F_2(x,Q^2) - F_L(x,Q^2)}. \tag{2.8}$$

It can be seen from equation (2.8) that the condition that the transverse and longitudinal photon induced cross sections must be positive, leads to

$$0 \le F_L \le F_2. \tag{2.9}$$

From equation (2.9) and the fact that the F_L term appears with a proportionality factor y^2 in equation (2.5), it can be seen that the F_2 term is dominant in regions of low y, with the F_L contribution increasing at high y. From the difference between the measured σ_r and the (from low y) extrapolated F_2, $F_L(x, Q^2)$ is determined as described in [4], [5].

Another way of looking at the total cross section, $\sigma_{tot} = \sigma_T + \sigma_L$, is by describing it as a convolution of a flux of quasi-real or virtual gauge bosons from the electron with the absorption cross section of that boson by the proton as depicted in figure 2.2. At small enough Q^2, the Z^0 and $W^\pm$ exchange can be neglected and the boson can be interpreted as a virtual photon. This massive spin 1 particle acquires three polarization vectors corresponding to the three helicity states of the photon, $\lambda = \pm 1, 0$. The absorption cross section may than depend on the helicity. We will thus have two independent cross sections, one for absorbing a transversely polarized photon σ_T ($\lambda = \pm 1$) and one for a longitudinally polarized photon σ_L ($\lambda = 0$). The differential cross section can then be written in the form:

$$\frac{d^2\sigma(ep)}{d\,W\,d\,Q^2} = f_{\gamma/e}(\sigma_T^{\gamma p} + \epsilon\sigma_L^{\gamma p}) \tag{2.10}$$

where W represents the γp centre of mass energy, $f_{\gamma/e}$ the flux factor of virtual photons and $\epsilon = \frac{2(1-y)}{1+(1-y)^2}$ the ratio of the flux for longitudinal to transverse photons.



Figure 2.2: *At small Q^2, ep-scattering can be thought of as a radiation of a virtual photon γ^* from the electron, followed by $\gamma^* p$-scattering.*

2.2.1 The Quark Parton Model (QPM) and the scaling violation

The first DIS experiments at SLAC [9, 10] in the early sixties showed that the structure function $F_2(x, Q^2)$ exhibited a scaling behavior in the limit $Q^2 \to \infty$ predicted by Bjorken [6]:

$$F_2(x, Q^2) \to F_2(x). \tag{2.11}$$

In the quark parton model (QPM) [6, 7, 8], in which the proton is considered to be made of point-like charged quarks, the structure function F_2 is expected to be independent of Q^2 and is related to the parton momentum distributions $f_i(x)$ of flavor i by

$$F_2(x, Q^2) = \sum_i e_i^2\, x f_i(x) \tag{2.12}$$

where e_i is the electric charge of parton i. In the QPM, for spin-1/2 partons, $R = 0$ [3], implying that $\sigma_L(x) = F_L(x) = 0$. The parton distributions $f_i(x)$ are constrained by the sum-rule:

$$\sum_i \int x f_i(x) dx = 1, \tag{2.13}$$

where the summation runs over all species of charged partons. However measurements showed that this sum-rule is not fulfilled and that quarks and anti-quarks account for only half of the proton's momentum [15],[16]. Observation at fixed target experiments [11, 12, 13] later verified by HERA [14], showed that the scaling behavior is applicable for values of $x \approx 0.13$, but outside this x-region a weak logarithmic dependence on Q^2 is observed, implying a violation of the scaling hypothesis. The QPM is only a zeroth order description for the deep inelastic scattering process. The partons in the proton or more generally in the hadron, are not free objects and they interact with each other via a color field mediated by gluons. The theory that describes the interactions of quarks and gluons is called *Quantum ChromoDynamics* (QCD).

The scaling violations, observed in figure 2.4, which is an F_2-compilation of the most up to date measurements [17, 18, 5, 19, 20, 12, 21, 11], can be explained by considering processes of the type depicted in figure 2.3. The result of this is that for instance the quark distribution becomes scale dependent and satisfies the integro differential equation

$$\frac{dq(x,Q^2)}{d\ln Q^2} = \frac{\alpha_s(Q^2)}{2\pi} \int_x^1 \frac{dy}{y} q(y,Q^2) P_{qq}\left(\frac{x}{y}\right) \tag{2.14}$$

which is one of the DGLAP (Dokshitzer, Gribov, Lipatov, Altarelli and Parisi) [22] evolution equations. The splitting function $P_{qq}(x/y)$ represents the probability for a quark to emit another quark with momentum fraction x and can be expanded as a power series in α_s:

$$P_{qq}(x/y) = \sum_n \alpha_s^n \, p^{(n)}(x/y). \tag{2.15}$$



Figure 2.3: *The three different splitting functions $P_{ij}(x/y)$ of first order in α_s are shown.*

Not only the quarks but also the gluon distributions evolves, such that the full DGLAP equation becomes:

$$\frac{dq(x,Q^2)}{d\ln Q^2} = \frac{\alpha_s(Q^2)}{2\pi} \int_x^1 \frac{dy}{y} \left(q(y,Q^2) P_{qq}\left(\frac{x}{y}\right) + g(y,Q^2) P_{qg}\left(\frac{x}{y}\right) \right) \tag{2.16}$$



Figure 2.4: F_2 *is plotted versus* Q^2 *for fixed* x *values for the HERA data, as well as for the fixed target data.* F_2 *shows strong scaling violations particularly at small* x*.*

$$\frac{dg(x,Q^2)}{d\ \ln Q^2} = \frac{\alpha_s(Q^2)}{2\pi}\int_x^1 \frac{dy}{y}\left(\sum_q q(y,Q^2)P_{gq}\left(\frac{x}{y}\right) + g(y,Q^2)P_{gg}\left(\frac{x}{y}\right)\right) \tag{2.17}$$

where y is the momentum of the emitting parton and x is the fractional momentum of the resulting parton as illustrated in figure 2.3.

In leading order (LO), the DGLAP equation resums all contributions of the type $(\alpha_s\ \ln Q^2)^n$ and the actual gauge corresponds to the emission of a chain of n gluons with strongly transverse ordering:

$$Q^2 \gg k_{T,n}^2 \gg k_{T,n-1}^2 \gg ... \gg k_{T,1}^2 \gg Q_0^2, \tag{2.18}$$

and also weaker ordering of longitudinal momenta,

$$x < x_n < x_{n-1} < ... < x_1. \tag{2.19}$$

In the next-to-leading order (NLO) also terms of the type $\alpha_s^n\ (\ln Q^2)^{(n-1)}$ are resummed. In fits to structure function measurements, the parton density functions $q_i(x,Q^2)$ and $g(x,Q^2)$, which are inherently non-perturbative, are parameterized at a starting scale Q_0^2. The equations (2.16) and (2.17) can be used to predict the parton distributions at any other value of Q^2. An example of the gluon distribution obtained at NLO is shown for three different values of Q^2 in figure 2.5 [5]. The gluon contribution is dominant at the lowest values of x.

The large centre of mass energy at HERA, not only may result in a large Q^2 but also in a large $1/x \propto ys/Q^2$-value. This means that not only leading log in Q^2 may be important but also those in $\alpha_s\ \ln(1/x)$. A resummation of those terms not coupled to large $\ln(Q^2)$ leads to the BFKL (Balitsky, Fadin, Kuraev, Lipatov) [23] formalism. In contrast to the DGLAP approximation there is a strong ordering of longitudinal momentum fractions,

$$x \ll x_n \ll x_{n-1} \ll ... \ll x_1, \tag{2.20}$$

and no ordering of k_T.

To date the DGLAP evolution scheme alone can describe structure function data throughout the entire HERA phase space. Hence, so far there is no unambiguous evidence for the need for BFKL contributions. Figure 2.6 shows the kinematical plane in x and Q^2, illustrating the regions of validity for both formalisms.

2.3 Diffractive scattering

One of the surprises that came along with the first results from HERA was the observation of DIS events with a large rapidity gap in the hadronic final state, located between the photon and proton fragmentation regions [31, 32]. The observed fraction of events with large rapidity gaps is of the order of 10%, fairly independent of the γp centre of mass energy, W, and the photon virtuality squared, Q^2. In QCD, the fragmentation process driven by parton radiation leads to an exponential suppression of large rapidity gaps between hadrons [33]. Large rapidity gap formation, with little dependence on W, is typical of diffractive



Figure 2.5: *The gluon distribution from next-to-leading order (NLO) DGLAP fits with starting scale* $Q_0^2 = 4.0\ GeV^2$.

scattering and invokes the notion of the pomeron. The fact that the diffractive exchange can be probed with large virtual photons means that its structure can be studied in much the same way as the partonic structure of the proton. The large range of the photon virtuality available at HERA opened the possibility to measure precisely the partonic structure of the diffractive exchange (see section 2.4).

At HERA, one usually identifies three types of interactions that differ in the characteristics of the final states. In elastic scattering, both initial particles emerge relatively unscathed and no other particles are produced. A diagram of the reaction $\gamma p \to Vp$ where V is any vector meson, may be considered in the framework of the VDM model [34], in which the photon is considered to consist of a two component state (i.e a part interacting only electro-magnetically and a hadronic part undergoing conventional strong interactions) to be elastic. Such an elastic (EL) γp reaction is shown in figure 2.7.a. In diffractive scattering, the energy transfer between the two interacting particles remains small, but one (single dissociation, i.e. gamma (GD) or proton dissociation (PD)) or both (double dissociation



Figure 2.6: *The kinematic plane in x and Q^2 showing where the DGLAP and BFKL approximations are theoretically valid.*

(DD)) particles dissociate into multiparticle final states preserving the quantum numbers of the initial particle. Figures 2.7.b-d show the three distinct classes of dissociative diffraction that can occur with single pomeron exchange in γp-collisions. The remaining configurations correspond to so called inelastic interactions which in contrast with the elastic and diffractive scatterings show no large rapidity gaps in the final state.

Characteristic of high energy elastic and diffractive scattering is the exponential behavior of the cross section as a function of the square of the momentum transfer t. As this property is reminiscent of the diffraction of light by a circular aperture, diffractive scattering in strong interactions derives its name from the optical analogy [35]. The slope of the t-distribution will be linked with the radius of the interacting object in the following section.

Defining in an unique way what diffraction is, is very hard. But to get an idea how a single diffractive scattering can come true, two constraints have to be valid:



Figure 2.7: *Diffractive interactions proceeding by single pomeron exchange, occurring in γp-collisions. Figure a) represents the EL process, b) the GD process, c) the PD process and d) the DD process. The symbol $I\!P$ is used to denote the exchange of the leading pomeron trajectory.*

- **conservation of intrinsic quantum numbers of the initial state**
  The diffractive state preserves the intrinsic quantum numbers of the initial particles. However spin and parity can be changed since the momentum transfer of the interaction may change the internal motion of the interacting hadrons.

- **coherence constraint**
  The absorbing hadron should preserve its identity, which limits the invariant mass of the diffracted state. The minimum momentum transfer squared for a particle of mass m and momentum p to create a mass M_X via the interaction with a stationary target is
  $$| t_{min} | = \left(\frac{M_X^2 - m^2}{2p} \right)^2 . \qquad (2.21)$$
  Using the uncertainty principle, the coherence of the absorbing hadron of radius R will be preserved if
  $$(M_X^2 - m^2) \leq \frac{2p}{R}. \qquad (2.22)$$

Hence, when studying diffractive events, the study of low momentum transfer is especially of great interest [36, 39, 37]. In the pre QCD era these events have been successfully described in the framework of Regge-theory [38] which is based on the S-matrix formalism, in which scattering processes are described in terms of the exchange of reggeons. This Regge framework will be described below.

2.3.1 Diffractive scattering linking with optical diffraction

In figure 2.8 the cross section for elastic proton-proton scattering is plotted for different centre of mass energies as a function of $\mid t \mid$ where t is the squared momentum transfer between the two hadrons. For low centre of mass energies a falling cross section is observed where the slope gets steeper towards higher centre of mass energies. In addition a minimum and a maximum appear. This shape of the cross section down to the first minimum can be approximated by an exponential function of the form:

$$\frac{d\sigma/dt}{(d\sigma/dt)_{t=0}} = e^{bt} \approx 1 - b(p\theta)^2 \tag{2.23}$$

with $b \sim 10$ GeV^{-2}.

The data can be interpreted in the spirit of the similarity between optical and hadronic diffraction [39]. In this picture one discusses the scattering of a fast hadron represented by a plane wave in the field of a hadron at rest. In optical diffraction the following formula holds for $kR \ll 1$:

$$\frac{I}{I_0} \approx 1 - \frac{R^2}{4}(k\theta)^2 \tag{2.24}$$

which describes the fraction of observed intensity with respect to the initial intensity as a function of the diffraction angle θ, where R is the radius of the disc and k denotes the wave number of the incident light wave. If one relates the scattering angle θ to the momentum transfer t in hadron-hadron interactions, the wave number k to the momentum of the hadrons, and the radius of the disc R to the mass of the pion one can deduce a t-slope of $b = 12.5$ GeV^{-2} resulting in the right order of magnitude as the one measured.

2.3.2 The S-matrix and aspects of Regge theory

In the study of hadron-hadron interactions at low momentum transfer t, where no hard scale is present and perturbative QCD is not applicable, the phenomenological models very often still give the best descriptions available. A consistent description of such strong interactions is offered by Regge theory.

Regge theory was introduced in the framework of non-relativistic quantum mechanics and generalized the ideas of Yukawa explaining particle interactions by the exchange of virtual particles. Later it was generalized to describe also relativistic particle interactions.

Since Regge theory is based on the S-matrix formalism we will first briefly introduce the basic assumptions:

- **Unitarity of S**
  The scattering matrix is unitary, which means that the probability that there will be some final state starting from any given state $\mid i\rangle$ must be unity and is expressed as:

$$S^\dagger S = 1 = SS^\dagger \tag{2.25}$$



Figure 2.8: *Proton-proton elastic differential cross section versus four-momentum transfer squared at various incident proton momenta. As the energy increases, a classical diffraction pattern as in optical diffraction appears.*

- **Maximal analyticity**
  If one considers the reaction $A + B \to C + D$, the Mandelstam variables are real numbers defined by relations of the form $s_{ij} = (\pm P_i \pm P_j)^2$ with $i, j \in \{A, B, C, D\}$. Maximal analyticity of first order means that these variables can be extended into the complex plane which allows the use of complex analysis. The S-matrix can now be written as a function of these complex numbers. In the limit of real numbers one gets back the S-matrix for particle reactions in the physical region.

- **Crossing symmetry**
  A consequence of the maximal analyticity assumption is the crossing symmetry which states that if the reaction $A+B \to C+D$ is described by the variable $s = (P_A + P_B)^2$ and $t = (P_A - P_C)^2$ by the S-matrix $S(s,t)$ one can calculate the amplitude for the reaction $A + \overline{C} \to \overline{B} + D$ by exchange of s and t which leads to the amplitude $S(t,s)$ i.e.
  $$S_{AB \to CD}(s,t) = S_{A\overline{C} \to \overline{B}D}(t,s). \tag{2.26}$$

- **Generalization of angular momentum in the complex plane**
  In the s-channel, all the possible resonances which can be produced by the interaction of A and B and which conserve quantum numbers (parity, charge etc.) are linked by a linear trajectory in a two-dimensional space defined by mass squared and angular momentum. In Regge theory, angular momentum is treated as a continuous complex variable, $\alpha(t)$, with physically observable t-channel resonances occurring only at physical values of spin, such that $\mathcal{R}e(\alpha(t))$ is an integer or half integer, known as Regge poles. Stable hadrons have pure real values of $\alpha(t)$, whilst unstable hadrons have an imaginary component that is related to their decay width. Bound states therefore lie near to the real axis of $\alpha(t)$. The linear trajectories can be parameterized as:
  $$\alpha(t) = \alpha(0) + \alpha' t. \tag{2.27}$$
  As an example, the trajectory for which the ρ meson is the lowest lying state is shown in figure 2.9.

Amplitudes in the t-channel may be computed by decomposition into partial waves of different angular momenta

$$A(s,t) = \sum_{l=0}^{\infty} (2l+1)\ A_l(t)\ P_l(\cos\theta) \tag{2.28}$$

where $A_l(t)$ is the amplitude of the l^{th} partial wave, and $P_l(\cos\theta)$ is the Legendre polynomial for angular momentum l. In continuing the amplitude to describe the s-channel, where $\cos\theta > 1$, equation (2.28) may be re-expressed as a sum over poles. When expressed in terms of contributing Regge trajectories, the pole in the l^{th} partial wave may be written as [38]

$$A_l(t) = \frac{\beta(t)}{l - \alpha_l(t)}. \tag{2.29}$$

At $l = \alpha_l(t)$ a Regge pole occurs with a residue function described by $\beta(t)$.



Figure 2.9: *The ρ trajectory, $\alpha_\rho(t)$. The data shown with triangles at negative t are measured in the charge exchange reaction, $\pi^- p \rightarrow \pi^0 n$, and are taken from [40]. In the t-channel physical region the bound states (ρ, ρ_3) are shown. A straight line through these states is continued into the s-channel region. The comparison there with the measured hadron-hadron data is remarkable.*

In the asymptotic limit where $s \rightarrow \infty$ and $t/s \rightarrow 0$, only the leading trajectory contributes and equation (2.28) becomes:

$$A(s,t) = f(t) \left(\frac{s}{s_0} \right)^{\alpha(t)} \tag{2.30}$$

where $f(t)$ includes $\beta(t)$ and the part of the amplitude not depending on s. s_0 defines the energy scale on which s must be large, typically taken to be ~ 1 GeV^2. Hence, at high energies the s-dependence of the s-channel scattering amplitude is a simple power behavior.

Using the optical theorem, relating the forward scattering amplitude to the total cross section

$$\sigma_{tot} = \frac{1}{s} \mathcal{I}m(A(s, t=0)) \tag{2.31}$$

one obtains, combining equations (2.30) and (2.31), the following expression for $\sigma_{tot}(s)$

$$\sigma_{tot}(s) \propto \left(\frac{s}{s_0} \right)^{\alpha(0)-1} . \tag{2.32}$$

The measurement of the energy dependence of the cross section therefore provides access to the intercept $\alpha(0)$ of the leading trajectory. When measuring also the t-dependence (see section 2.3.3) of cross sections, the determination of the slope α' becomes possible.

2.3.3 t-dependence and shrinkage

For a process that is dominated by the exchange of a single trajectory, the expression for the differential cross section follows from equations (2.27) and (2.30) and is given by,

$$\frac{d\sigma}{dt} \propto \frac{1}{s^2} \mid A(s,t) \mid^2 \tag{2.33}$$

$$\propto F(t) \left(\frac{s}{s_0}\right)^{2(\alpha(0)-1)} \left(\frac{s}{s_0}\right)^{2\alpha' t}. \tag{2.34}$$

If the dependence of the coupling, $F(t)$, is small in t, then at large s, the t-dependence must be dominated by the final factor of equation (2.34). Since t is negative, and α' positive, $d\sigma/dt$ decreases faster as a function of t as s increases. Regge theory thus predicts that if the centre of mass energy increases the forward peak shrinks as seen experimentally in figure 2.8.

Since the fast variation of the differential cross section with t at fixed s is found experimentally to be well described at low $\mid t \mid$ by an exponential parameterization, equation (2.34) may be re-expressed as

$$\frac{d\sigma}{dt} = \left(\frac{d\sigma}{dt}\right)_{t=0} e^{bt} \tag{2.35}$$

where b, known as the slope parameter, is given by;

$$b \sim b_0 + 2\alpha' \ln\left(\frac{s}{s_0}\right) \tag{2.36}$$

and describes a logarithmic shrinkage of the forward peak with increasing centre of mass energy s.

It was observed (see figure 2.10) that the total cross section in $p\overline{p}$- and pp-interactions and later in γp-interactions, decreases with energy at low energies which is explained by Regge trajectories, whereas the rise of the cross section for high energy was not expected. To account for this rise the **pomeron** was introduced which is characterized by the quantum numbers of the vacuum. Donnachie and Landshoff parameterized this energy behavior of the total cross section by the sum of two reggeon amplitudes ($I\!\!P$ and $I\!\!R$):

$$\sigma_{tot}(s) = A_{I\!\!P} s^{\alpha_{I\!\!P}(0)-1} + A_{I\!\!R} s^{\alpha_{I\!\!R}(0)-1} \tag{2.37}$$

with $A_{I\!\!P}$ and $A_{I\!\!R}$ the normalization parameters and $\alpha_{I\!\!P}(0)$ and $\alpha_{I\!\!R}(0)$ the trajectory intercepts also left as free parameters. To account for the rising cross section at high energies the pomeron intercept in equation (2.37) had to be larger than one. The fit of equation (2.37) to the $p\overline{p}$- and pp-data (see figure 2.10.a) results in,

$$\alpha_{I\!\!P}(0) = 1 + \epsilon = 1.08 \tag{2.38}$$

$$\alpha_{I\!\!R}(0) = 0.55. \tag{2.39}$$

Assuming $I\!\!P/I\!\!R$-universality, the intercepts were fixed to the values obtained from the fit on the $p\overline{p}$- and pp-data, and a new fit was performed on π^-p-, and π^+p-data (see figure 2.10.b))



Figure 2.10: *Fits performed by Donnachie and Landshoff to the centre of mass energy dependence of the total cross section for a) pp-and $p\overline{p}$-scattering [41], b) π^-p and π^+p scattering [41], and c) γp-scattering [42]. The high s points in the $p\overline{p}$ case are from the Tevatron. Those in the γp case are from HERA. In c) the solid line is the Donnachie Landshoff fit with $\alpha_{I\!P}(0) = 1.08$, the dotted line with $\alpha_{I\!P}(0) = 1.11$, whilst the broad dashed curve represents the ALLM parameterization [43].*

and γp-data (see figure 2.10.c)) leaving only the normalizations free in the fit. The reggeon has the larger coupling to hadrons, and consequently dominates at low centre of mass energies. However, the reggeon contribution falls approximately as $s^{-1/2}$, such that at large centre of mass energies, the pomeron term ($\propto s^{0.08}$) dominates, and the cross section starts to rise slowly with s. At large centre of mass energies, the coupling strength to particle and anti-particle are observed to become the same leading to similar cross sections for particle-particle interactions with respect to particle-anti-particle interactions, in accordance with the Pomerancuk theorem.

Assuming an s-dependence of the slope b as given by equation (2.36) in the $\overline{p}p$ elastic scattering data at the Tevatron Collider and the CERN ISR data, yields a value of the pomeron slope of $\alpha'_{I\!P} = 0.26 \pm 0.02$ GeV^{-2} [44]. The pomeron trajectory takes thus the form:

$$\alpha_{I\!P}(t) = 1.08 + 0.26\,t. \tag{2.40}$$

2.4 Diffraction in DIS

Where, in the region of inclusive deep inelastic scattering at high Q^2 a hard scale is present, suitable to resolve the partonic content of the proton, here in diffractive scatterings, this hard scale can be used in resolving the partonic content of the pomeron. This analogy leads to the Ingelmann-Schlein approach [45], that the diffractive deep inelastic scattering in γp-interactions proceeds in two steps as depicted in figure 2.11: first a pomeron is emitted from the proton and then the virtual photon is absorbed by a constituent of the pomeron. In order to describe this process new variables are introduced, in addition to the inclusive



Figure 2.11: *Pictorial representation of the common diffractive variables.*

DIS variables from section 2.1. Defining the total γp centre of mass energy, the invariant mass of system X, and momentum transfer to the scattered proton respectively as:

$$W^2 = (q+P)^2 \tag{2.41}$$

$$\begin{aligned} M_X^2 &= (q + P - P')^2 \quad (2.42) \\ t &= (P - P')^2. \quad (2.43) \end{aligned}$$

The variable $x_{I\!P}$ which represents the momentum fraction of the proton carried by the pomeron defined as

$$x_{I\!P} = \frac{q.(P - P')}{q.P} \quad (2.44)$$

can be rewritten as

$$x_{I\!P} = \frac{M_X^2 + Q^2 - t}{W^2 + Q^2 - m_p^2} \simeq \frac{M_X^2 + Q^2}{W^2 + Q^2} \quad (2.45)$$

where P and P' denote respectively the initial and final four-momenta of the proton and where m_p stands for the mass of the proton. The last relation on the right-hand side holds for $m_p^2 \ll Q^2$, W^2 and $\mid t \mid \ll Q^2$, M_X^2 which is a good approximation in the kinematical region of high energy diffraction at HERA.

Since the deep inelastic scattering can now be interpreted to take place on the pomeron, one can define an equivalent variable as Bjorken x but now relative to the pomeron momentum instead of the proton momentum. This variable called β, representing the fraction of the four-momentum of the pomeron carried by the struck quark, is given by,

$$\beta = \frac{Q^2}{2q.(P - P')} = \frac{x}{x_{I\!P}} = \frac{Q^2}{Q^2 + M_X^2 - t} \simeq \frac{Q^2}{Q^2 + M_X^2}. \quad (2.46)$$

2.4.1 Diffractive structure function

Analogous to the inclusive DIS cross section (see section 2.2), the diffractive DIS cross section can be written, after integration over t, as a three fold differential cross section depending on $x_{I\!P}$, β, andQ^2 [48],

$$\frac{d\sigma^D(ep \to epX)}{d\beta\, dQ^2\, dx_{I\!P}\, dt} = \frac{4\pi\alpha_{em}^2}{\beta Q^4}\left(1 - y + \frac{y^2}{2}\right)\sigma_r^{D(3)}(\beta, Q^2, x_{I\!P}, t). \quad (2.47)$$

In the region, $Q^2 \ll M_{Z^0}^0$, where contributions from Z^0 exchange can be neglected, $\sigma_r^{D(3)}$ is related to the diffractive structure functions $F_2^{D(3)}$ and $F_L^{D(3)}$ by,

$$\sigma_r^{D(3)} = F_2^{D(3)} - \frac{y^2}{1 + (1 - y)^2} F_L^{D(3)} \quad (2.48)$$

where F_L^D is the longitudinal diffractive structure function. Also here the positivity constraint on the cross sections for longitudinally and transversely polarized photon scattering leads to the relation $0 \le F_L^D \le F_2^D$. The contribution of F_L^D to $\sigma_r^{D(3)}$ can be sizeable only at large values of y, and in most of the kinematic range the relation $\sigma_r^{D(3)} \approx F_2^D$ holds to good approximation. In contrast to the inclusive deep inelastic scattering were F_L is measured in the range of large y, F_L^D is not yet measured in diffractive scattering interactions.

QCD hard scattering factorization in diffraction

The proof that QCD hard scattering factorization can be applied to diffractive DIS [46] implies that in the leading $\log(Q^2)$ approximation, the cross section for the diffractive process $\gamma^\star p \to XY$ can be written in terms of convolutions of universal partonic cross sections $\sigma^{\gamma^\star i}$ with diffractive parton distributions F_i^D [47], representing probability distributions for a parton i in the proton under the constraint that the proton is scattered with particular values of t and $x_{I\!P}$. Thus, at leading twist,

$$\frac{d^2\sigma(x, Q^2, x_{I\!P}, t)^{\gamma^\star p \to p' X}}{dx_{I\!P}\ dt} = \sum_i \int_x^{x_{I\!P}} d\xi\ \sigma^{\gamma^\star i}(x, Q^2, \xi)\ f_i^D(\xi, Q^2, x_{I\!P}, t). \tag{2.49}$$

The factorization formula is valid for large enough Q^2 and fixed $x_{I\!P}$ and t. The partonic cross sections $\sigma^{\gamma^\star i}$ are the same as those for inclusive DIS and the diffractive parton distributions f_i^D, which are not known from first principle, should obey the DGLAP evolution equations.

Regge factorization

An additional assumption is often made in the analysis of diffractive cross sections, saying that the shape of the diffractive parton distribution is independent of $x_{I\!P}$ and t and that their normalization is controlled by Regge asymptotics [45]. The diffractive parton distributions can then be factored into a term depending only on $x_{I\!P}$ and t and a second term depending only on β and Q^2 (see figure 2.12):

$$f_i^D(x_{I\!P}, t, \beta, Q^2) = f_{I\!P/p}(x_{I\!P}, t).f_i^{I\!P}(\beta, Q^2). \tag{2.50}$$

Under this Regge factorization assumption, often refered to in the literature as the Ingelman-Schlein or resolved pomeron model, the diffractive exchange can be treated as a quasi-real object with a partonic structure, given by parton distributions $f_i^{I\!P}(\beta, Q^2)$. The variable β then corresponds to the longitudinal momentum fraction of the diffractive exchange carried by the struck parton in the pomeron. The pomeron flux factor $f_{I\!P/p}(x_{I\!P}, t)$ represents the probability that a pomeron with particular values of $x_{I\!P}$ and t couples to the proton. Although equation (2.50) has no firm basis in QCD, at the present level of precision it appears to be supported by data [48, 49, 50].

Figure 2.13 show the $x_{I\!P}$-dependence of the measured diffractive reduced cross section $\sigma_r^{D(3)}(\beta, Q^2, x_{I\!P})$ [51] for fixed β and Q^2 respectively, using a parameterization of the form

$$\sigma_r^{D(3)}(x_{I\!P}, \beta, Q^2) = f_{I\!P}(x_{I\!P})A_{I\!P}(\beta, Q^2) + f_{I\!R}(x_{I\!P})A_{I\!R}(\beta, Q^2) \tag{2.51}$$

where the pomeron and reggeon flux factors are defined as

$$f_{\{I\!P, I\!R\}}(x_{I\!P}) = \int_{t_{cut}}^{t_{min}} \frac{e^{B_{\{I\!P,I\!R\}}t}}{x_{I\!P}^{2\alpha_{\{I\!P,I\!R\}}(t)-1}}\ dt \tag{2.52}$$

where the pomeron and reggeon trajectories are assumed to be linear. The free parameters in the fit are $\alpha_{I\!P}(0)$, $A_{I\!P}(\beta, Q^2)$ and $A_{I\!R}(\beta, Q^2)$. A striking feature of the data presented



Figure 2.12: *Representation of the Regge factorization assumption.*

in [51], is the scaling violations with positive $\partial\sigma_r^{D(3)}/\partial \ln Q^2$ throughout most of the phase space, becoming negative only at the highest measured values of β. This behavior is different from that observed for the scaling violation of $F_2(x, Q^2)$ at fixed x, which is negative for $x \geq 0.1$. In a QCD interpretation, this means that the measured Q^2-dependences are strongly suggestive of a large gluonic component of the diffractive exchange.

In order to investigate whether $\alpha_{I\!P}(0)$ has any dependence on Q^2, the fits were performed in different Q^2-intervals. Those results of the pomeron intercepts are shown in figure 6.7 at $Q^2 > 0$ GeV^2 with some values of the pomeron intercept extracted in this thesis at $Q^2 = 0$ GeV^2.

2.5 Diffractive dissociation in photoproduction

Returning back to the low Q^2-region of soft physics, where no QCD predictions exist, Regge theory offers predictions in terms of trajectories for the dynamics of dissociative diffraction, and of inclusive processes in general. Feynman [53] predicted that high energy inclusive reactions of the type

$$A + B \to C + X$$

where C is the observed particle and X represents everything else produced, should scale in analogy to Bjorken scaling in DIS, such that cross sections should be function of the transverse and longitudinal momenta of the particle C, but not of the centre of mass energy. An extended version of the optical theorem for such inclusive processes, depicted in figure 2.14, was proposed by Mueller [54], relating the inclusive reaction $AB \to CX$ to the theoretically much simpler three-body elastic scattering process, $AB\overline{C} \to AB\overline{C}$,

$$f \equiv E_C \frac{d\sigma}{d^3p_C}(A + B \to C + X) \simeq \frac{1}{s}\mathrm{Disc}_{M^2} A(AB\overline{C} \to AB\overline{C}) \tag{2.53}$$

where the discontinuity is to be taken only across the M_X^2 cut of the elastic amplitude. Here M_X^2 is defined as $(P_A + P_B + P_{\overline{C}})^2$. The importance of Mueller's optical theorem is that the behavior of the inclusive reaction can be obtained from that of the theoretically much



Figure 2.13: *β-dependence (left) and Q^2-dependence (right) of the reduced cross section (left) scaled at each $x_{I\!P}$ by the values assumed for the t-integrated pomeron flux in the QCD fits. Only data with $y < 0.6$ are shown to minimize the influence of the longitudinal structure function F_L^D. The data are compared with the prediction of the NLO QCD fits under the assumption that $F_L^D = 0$.*



Figure 2.14: *Pictorial representation of Mueller's Optical theorem.*

simpler, elastic three-body amplitude. Much of the knowledge gained from Regge type analyses of two-body scattering was immediately carried over into the inclusive domain.

To calculate the inclusive cross section for $A+B \to C+X$ we shall assume Regge behavior for the three-body amplitude. It is necessary to distinguish between three kinematical regions for particle C, i.e. the beam and target fragmentation region and the central region. The beam and target fragmentation region corresponds to finite values of $t \equiv (p_A - p_C)^2$ and $u \equiv (p_B - p_C)^2$ respectively. Intuitively C can be regarded as a fragment of A (or B) if the momentum difference between them remains finite at large s. In the third region, i.e. the central region, where t and u are both large, C is not closely associated with either incoming particle. A general criterion for a Regge expansion of a multi particle amplitude, such as $AB\overline{C} \to AB\overline{C}$, is that the kinematic invariant spanning the Regge exchange should be large (a typical value $\geq$ 5 GeV^2). In the central region where both t and u are large, we have the double Regge limit. On the other hand, in the beam fragmentation region ($s \to \infty$, $t/s \to 0$) there are three different Regge limits when different combinations of the invariants M_X^2 and s/M_X^2 become large.

2.5.1 The fragmentation region

Typical Regge contributions for the three different limits [38] are shown in figure 2.15.



Figure 2.15: *The single, normal and triple Regge limit is depicted in the beam fragmentation region.*

1. **The single Regge limit ($s_{AB} \to \infty$ and $x_{I\!P} = M_X^2/s_{AB}$ fixed $\Rightarrow M_X^2 \to \infty$).** Since M_X^2 is the total centre of mass energy for the $AB\overline{C}$ elastic scattering process, and since large centre of mass energy (here M_X^2) and small $|\, t \,|$ suggest a single Regge pole exchange picture as in figure 2.15.a, this leads to an inclusive cross section of the form;

$$s \frac{d^2\sigma}{dt\, dM_X^2} = \sum_i \beta_i \left(\frac{M_X^2}{s}, t\right) \left(\frac{M_X^2}{s_0}\right)^{\alpha_i(0)-1} \tag{2.54}$$

where i runs over all possible reggeons that can be exchanged between particles A and B. Notice also that the argument of α_i is taken at $t_{BB'}$ which is for this forward three-body process always equal to 0, and should not be confused with $t \equiv t_{AC}$ which gives the invariant mass of the quasi-particle ($A\overline{C}$). Assuming the leading singularity, the pomeron, has $\alpha_{I\!P}(0) = 1$ and the leading meson exchanges have $\alpha_{I\!R}(0) = 0.5$, the inclusive cross section will scale for $M_X^2 \to \infty$. At low invariant mass reggeon contribution will go down as $(M_X^2/s_0)^{-1/2}$. The same behavior was also seen in the total cross sections as a function of the centre of mass energy s (see figure 2.10).

2. **The normal Regge limit (M_X^2 fixed , $s/M_X^2 \to \infty$).**
   In this case equation (2.53) yields an inclusive cross section:

$$s\frac{d^2\sigma}{dt\, dM_X^2} = \frac{1}{s}\sum_{ij} \beta^i_{AC}(t)\beta^j_{AC}(t) \left(\frac{s}{M_X^2}\right)^{\alpha_i(t)+\alpha_j(t)} \quad \text{Im}\, A_{(iB\to jB)}(\nu, t_{AC}, t_{BB}=0) \tag{2.55}$$

   where the sum is over all contributing trajectories. β^i_{AC} are the reggeon couplings. $A_{(iB\to jB)}(\nu, t_{AC})$ is usually refered to as the forward reggeon-particle scattering amplitude with $\nu \equiv M_X^2 - t - m_p^2$ and the mass of the state X constrained to be M_X^2 as indicated by the dashed line in the middle diagram of figure 2.15.

3. **The triple Regge limit ($M_X^2 \to \infty$, $s/M_X^2 \to \infty$ or $s_{AB} \gg M_X^2 = s_{\alpha_i B} \gg t_{AC}$).**
   This kinematic region is the overlap of the two preceding regions. For large M_X^2 the Reggeon-particle forward scattering amplitude of equation (2.55), which is related by Mueller's optical theorem via an intermediate state with mass M_X^2 to the Reggeon-particle cross section, can be expanded in terms of Regge exchanges in the $B\overline{B}$-channel which leads to the following triple Regge expansion:

$$\text{Im}\, A_{(iB\to jB)}(M_X^2, t) = \sum_k g_{ijk}(t)\beta^k_{BB}(0)\left(\frac{M_X^2}{s_0}\right)^{\alpha_k(0)} \tag{2.56}$$

   where g_{ijk} is the triple reggeon coupling at the centre of the last diagram of figure 2.15. The overall differential cross section in the triple Regge limit is then given, from equations (2.55) and (2.56), by

$$s\frac{d^2\sigma}{dt\, dM_X^2} = \frac{1}{s}\sum_{ijk} \beta^i_{AC}(t)\beta^j_{AC}(t)\beta^k_{BB}(0)g_{ijk}(t) \left(\frac{s}{M_X^2}\right)^{\alpha_i(t)+\alpha_j(t)} \left(\frac{M_X^2}{s_0}\right)^{\alpha_k(0)} \tag{2.57}$$

2.5.2 The triple pomeron amplitude

In the asymptotic limit in which $M_X^2 \to \infty$ and $s/M_X^2 \to \infty$ only the $I\!PI\!PI\!P$ term survives and equation (2.57) reduces to the triple pomeron amplitude given by:

$$\frac{d^2\sigma}{dtdM_X^2} \simeq \frac{\beta^2_{AI\!P}(t)\beta_{BI\!P}(0)g_{I\!PI\!PI\!P}(t)}{s^2}\left(\frac{s}{M_X^2}\right)^{2\alpha_{I\!P}(t)}\left(\frac{M_X^2}{s_0}\right)^{\alpha_{I\!P}(0)}. \tag{2.58}$$

When rewriting equation (2.58) as:

$$\frac{d^2\sigma}{dtdx_{I\!P}} \simeq \beta^2_{AI\!P}(t)\beta_{BI\!P}(0)g_{I\!PI\!PI\!P}(t)\,(x_{I\!P}s)^{\alpha_{I\!P}(0)-1}\, x_{I\!P}^{(1-2\alpha_{I\!P}(t))} \tag{2.59}$$

$$\simeq \mathcal{F}_{I\!P/P}(x_{I\!P}, t)\sigma^{tot}_{I\!PB}(x_{I\!P}s, t) \tag{2.60}$$

where $x_{I\!P} \simeq M_X^2/s$ (see equation (2.44)), the pomeron flux is separated from the pomeron-particle cross section. The pomeron flux and the pomeron-particle cross section are respectively equal to:

$$\mathcal{F}_{I\!P/P}(x_{I\!P}, t) \propto \beta^2_{AI\!P}(t)x_{I\!P}^{1-2\alpha_{I\!P}(t)} \tag{2.61}$$

$$\sigma^{tot}_{I\!PB}(x_{I\!P}s, t) \propto \beta_{BI\!P}(0)g_{I\!PI\!PI\!P}(t)\,(x_{I\!P}s)^{\alpha_{I\!P}(0)-1} \tag{2.62}$$

Writing the pomeron trajectory as:

$$\alpha_{I\!P}(t) = 1 + \epsilon + \alpha'_{I\!P} t \tag{2.63}$$

the equation (2.58) becomes:

$$\frac{d^2\sigma}{dt dM_X^2} \simeq \kappa(t) s^{2\epsilon + 2\alpha'_{I\!P} t} \left(\frac{1}{M_X^2} \right)^{1+\epsilon+2\alpha'_{I\!P} t} \tag{2.64}$$

where $\kappa(t)$ contains all the couplings of equation (2.58).



Figure 2.16: *The differential cross section for the reaction $pp \to pX$ at a centre of mass energy $s = 930$ GeV^2 and momentum transfer $t = -0.75$ GeV^2 [56]. The line drawn is proportional to $1/M_X^2$, which fits the data well in the triple Regge region. The dashed line is the continuation of the triple Regge region fit.*

Figure 2.16 shows a measurement of the differential cross section $d^2\sigma/dtdM_X^2$ for diffractive *pp*-collisions at the CERN ISR. In the triple Regge region defined by the large dissociation masses beyond the resonance region, the $1/M_X^2$-dependence is clearly observed.

2.5.3 Interference terms

Since the pomeron and the f reggeon exchange identical quantum numbers (see table 2.1) they may interfere [57, 58, 59, 60]. Such a contribution is expressed in terms of the 'non-diagonal' amplitudes $\{I\!P I\!R\}I\!P$ and $\{I\!P I\!R\}I\!R$ where,

$$\{I\!P I\!R\}k \equiv I\!P I\!R k + I\!R I\!P k. \tag{2.65}$$

Restrictions are imposed on the magnitudes of the interference relative to the non-interference terms through the Schwartz inequality [61]

$$\left(\sum_k G_{\{I\!PI\!R\}k}(t)(M_X^2)^{\alpha_k(0)}\right)^2 \leq 4\sum_k G_{I\!PI\!Pk}(t)(M_X^2)^{\alpha_k(0)} \sum_k G_{I\!RI\!Rk}(t)(M_X^2)^{\alpha_k(0)}. \tag{2.66}$$

Following [62, 60] two component duality can be assumed in the reggeon-photon scattering amplitude, such that the diagonal and non-diagonal terms are separately related for $k = I\!P$ and $k = I\!R$. When maximal coherence is assumed and through equation (2.66), the interference couplings at $t = 0$ GeV^2 can be parameterized as:

$$G_{\{I\!PI\!R\}I\!P}(0) = 2\sqrt{|\ G_{I\!PI\!PI\!P}(0)G_{I\!RI\!RI\!P}\ |}\,, \tag{2.67}$$

$$G_{\{I\!PI\!R\}I\!R}(0) = 2\sqrt{|\ G_{I\!PI\!PI\!R}(0)G_{I\!RI\!RI\!R}\ |}. \tag{2.68}$$

2.5.4 Pion exchange

At high centre of mass energies and low $x_{I\!P}$ (i.e. $x_{I\!P} < 0.03$) the diffractive cross section is dominated by pomeron exchange [50] but at $x_{I\!P} > 0.03$ reggeon exchange contribution to the diffractive structure function becomes important [63]. And especially at even higher $x_{I\!P}$ (i.e. $x_{I\!P} \geq 0.1$) pion exchange becomes important [63]. Data taken at HERA where the scattered proton is measured in the Forward Proton Spectrometers lie in this high $x_{I\!P}$-region. Since real pion-hadron scattering data exist, the pion-flux factor $\mathcal{F}_{\pi/P}$, in contrast with pomeron-hadron data, can be normalized absolutely to:

$$\mathcal{F}_{\pi/P} \propto x_\pi^{1-2\alpha_\pi(t)}\,\frac{-t\,\exp\left(b_\pi(t-m_\pi^2)\right)}{(m_\pi^2-t)^2}, \tag{2.69}$$

so that the differential cross section for pion exchange has the following form:

$$s\frac{d^2\sigma}{dtdM_X^2} = \sum_{k=I\!P,I\!R}\frac{G_{\pi\pi k}(t)}{s}\left(\frac{s}{M_X^2}\right)^{2\alpha_\pi(t)}(M_X^2)^{\alpha_k(0)}\,\frac{-t\,\exp\left(b_\pi(t-m_\pi^2)\right)}{(m_\pi^2-t)^2}. \tag{2.70}$$

2.6 Review of diffractive dissociation analysis.

Since all ingredients of the triple Regge model are introduced, we can come to the review of existing diffractive dissociation analysis. First the results from the reaction $pp \to Xp$ and $\overline{p}p \to Xp$ at respective pp- and $\overline{p}p$-collider experiments will be discussed, followed by a summary of the existing HERA results on this topic.

2.6.1 Fixed target and ISR results

The application of triple Regge phenomenology to inclusive diffractive final states dates back to the early seventies [64]. These first analyses used data obtained with 24 GeV/c

PID	isospin	signature	C parity	G parity
$I\!\!P$	0	+	+	+
f	0	+	+	+
ρ	1	-	-	+
ω	0	-	-	-
a	1	+	+	-
π	1	-	+	-
γ	0,1	-	-	/

Table 2.1: *The most relevant quantum numbers of the considered reggeons.*

incident protons and concluded that they were still far from having verified the Regge predictions in these experiments. It is only with the arrival of FNAL fixed target and ISR data at $\sqrt{s} > 20$ GeV [65] that quantitative conclusions could be drawn. Two global Regge fits by Field and Fox [57] and Roy and Roberts [61], have been performed which have been comprehensively reviewed and re-analysed by Ganguli and Roy in [66]. Various combinations of the triple Regge couplings, represented in equation (2.72), were considered, being $G_{I\!\!PI\!\!PI\!\!P}$, $G_{I\!\!PI\!\!PI\!\!R}$, $G_{I\!\!RI\!\!RI\!\!P}$, $G_{I\!\!RI\!\!RI\!\!R}$, $G_{\pi\pi I\!\!P}$ and $G_{\pi\pi I\!\!R}$. The latter two couplings were added as it was pointed out [67] that the π contribution could be sizeable (50% at $x_{I\!\!P} \simeq 0.83$). In order to cover a wide M^2-range very low masses had to be included. This was achieved by using the finite mass sum rules (FMSR). The triple Regge couplings resulting from the fits as a function of t are shown in table 2.2. They will be used to compare the results on the pp-data with those from γp-data, see section 2.6.2 and 6.3. The trajectories in these fits were assumed to be $\alpha_{I\!\!P}(t) = 1 + 0.25t$, $\alpha_{I\!\!R}(t) = 0.5 + t$ and $\alpha_\pi(t) = -0.02 + t$.

-t	$G_{I\!\!PI\!\!PI\!\!P}(t)$	$G_{I\!\!PI\!\!PI\!\!R}(t)$	$G_{I\!\!RI\!\!RI\!\!P}(t)$	$G_{I\!\!RI\!\!RI\!\!R}(t)$	$G_{\pi\pi I\!\!P}(t)$	$G_{\pi\pi I\!\!R}(t)$
0.05	3.80 ± 0.13	0.73 ± 0.15	44 ± 5	1 ± 10	167	259 ± 200
(0.05	4.24 ± 0.24	0.51 ± 0.18	15 ± 5	27 ± 18	619 ± 112	21 ± 150)
0.1	2.74 ± 0.11	1.16 ± 0.12	38 ± 5	0 ± 9	124 ± 60	0 ± 116
0.15	1.89 ± 0.08	1.13 ± 0.10	35 ± 5	15 ± 11	104 ± 62	4 ± 153
0.2	1.43 ± 0.07	0.92 ± 0.10	35 ± 4	22 ± 11	47 ± 44	0 ± 147
0.25	1.11 ± 0.24	0.94 ± 0.12	21 ± 5	41 ± 12	49 ± 45	0 ± 139
0.35	0.80 ± 0.06	0.70 ± 0.07	23 ± 5	20 ± 16	133 ± 98	0 ± 88
0.45	0.54 ± 0.03	0.47 ± 0.05	22 ± 4	46 ± 15	34 ± 56	0 ± 105
	$4.95e^{4.95t}$	0.83	$47.7e^{1.89t}$	24.4	$200.0e^{4.38t}$	30.0

Table 2.2: *Triple Regge couplings (in mb/GeV^2) for the six terms considered in the fit from Ganguli and Roy [66]. A simple exponential fit parameterizes the t-dependence.*

A fixed target experiment of importance performed at Fermilab, where incident photons were scattered on a hydrogen target, measured the diffractive dissociation cross section,

$d^2\sigma/(dtdM_X^2)$, at fixed t and at two centre of mass energies W=12.9 and 15.3 GeV [74]. In the analysis, Chapin et al., tested the FMSR and the factorization hypothesis by comparing γp-data with $\pi^- p$-data which were obtained in the same experiment. In addition an exponential behavior in t and a dominant $1/M_X^2$-behavior for $M_X^2 > 4$ GeV^2 was observed.

Those data will be combined in a triple Regge fit to the main rapidity gap data presented in this analysis and leading proton data [69] which will be introduced in section 2.6.2. By doing so the covered range in W is extended by more than one order in magnitude towards lower centre of mass energies, providing us a much larger lever-arm in W to disentangle the W-dependence.

2.6.2 HERA results

Both Collaborations at HERA (ZEUS and H1) have used two different methods to select diffractive dissociative processes. The most straightforward selection consists in tagging the scattered proton by means of a forward or leading proton spectrometer, called FPS in the case of the H1 experiment or LPS for the ZEUS experiment. The other method relies on the presence of a rapidity gap in the forward part of the detector. Both methods have their advantages and disadvantages. Generally the leading proton spectrometers measure well the proton momentum without much background resulting from dissociative processes. However because of their limited acceptance, the accumulated statistics is rather limited. Almost the reverse is true for the rapidity gap selection where the collected statistics is much larger, but because the leading proton is not tagged, background resulting from low mass proton excitation processes can be substantial. They provide complementary information in that the mass range covered by both methods differ.

Tagged leading proton data

In the case where the leading proton is tagged in the forward proton spectrometers the photon dissociative mass M_X can be measured by measuring the centre of mass energy of the photon-proton system W, by $M_X = W\sqrt{1-z}$. Here z is the energy fraction of the proton leaving the interaction, $z = E'_p/E_p$, where E'_p and E_p represent the energy of the outgoing and incoming protons respectively. The process $\gamma p \to Xp$ is interpreted in the framework of triple Regge theory depicted in figure 2.17. The similarity between figure 2.15 and 2.17 is seen, but here the picture is presented for γp-scattering. The possibility of contributions of interference terms was also neglected in figure 2.17, such that the two reggeon trajectories coupling to the proton are identical. The cross section can than be expressed as a sum over contributing Regge trajectories such that equation (2.57) becomes:

$$W^2 \frac{d^2\sigma}{dtdM_X^2} = \frac{d^2\sigma}{dtdz} = \frac{s_0}{W^2} \sum_{i,k} G_{iik}(t) \left(\frac{W^2}{M_X^2}\right)^{2\alpha_i(t)} \left(\frac{M_X^2}{s_0}\right)^{\alpha_k(0)}. \tag{2.71}$$

Here α_i refers to the trajectory exchanged between the photon and the proton and α_k refers to the additional trajectory describing the total cross section between α_i and the photon. The triple Regge coupling $G_{iik}(t)$ may be factored into products of couplings $\beta_{pi}(t)$



Figure 2.17: *Representation of the Mueller-Regge approach for the inclusive photon dissociative cross section.*

of trajectory i to the proton, of couplings $\beta_{\gamma k}(0)$ of trajectory k to the photon and of a triple Regge coupling $g_{iik}(t)$ as:

$$G_{iik}(t) = \frac{1}{16\pi}\beta_{pi}^2(t)\beta_{\gamma k}(0)g_{iik}(t). \tag{2.72}$$

The t-dependence of the reggeon-proton and three-reggeon couplings are parameterized here as

$$\begin{aligned} \beta_{pi}(t) &= \beta_{pi}(0)e^{b_{pi}t} && (2.73)\\ g_{iik}(t) &= g_{iik}(0)e^{b_{iik}t}. && (2.74) \end{aligned}$$

With the three trajectories, $\alpha_{I\!P}$, $\alpha_{I\!R}$ and α_π, assuming no interference between the $I\!P$ and $I\!R$, equation (2.71) leads to a total of six terms with distinct W^2- and M_X^2-dependences. Those terms, all with their individual W^2- and M_X^2-dependences on the differential cross section $M_X^2 d\sigma/dM_X^2$ are listed in table 6.1.

In the triple Regge limit in which both $W^2/M_X^2 \to \infty$ and $M_X^2 \to \infty$ only the $iik = I\!PI\!PI\!P$ term survives and equation (2.71) reduces to

$$\frac{d^2\sigma}{dtdM_X^2} = \frac{G_{I\!PI\!PI\!P}(0)}{s_0^{\alpha_{I\!P}(0)-1}}\left(W^2\right)^{2\alpha_{I\!P}(0)-2}\left(\frac{1}{M_X^2}\right)^{\alpha_{I\!P}(0)} e^{B(W^2,M_X^2)t}, \tag{2.75}$$

where $B(W^2, M_X^2) = 2b_{pI\!P} + b_{I\!PI\!PI\!P} + 2\alpha'_{I\!P}\ln(W^2/M_X^2)$.

In this kinematical domain where $x_{I\!P} = M_X^2/W^2 = 1 - z \geq 0.12$, the complete mixture of $I\!P$, $I\!R$ and π contributing trajectories are important as explained by Field and Fox in [57]. But given the complexity of a triple Regge analysis and given the limited kinematical region to make a complete decomposition, the sum over all possible exchanges was replaced by single effective trajectories α_i and α_k. A triple Regge fit of the following form,

$$\frac{d\sigma}{dz} = W^2\frac{d\sigma}{dM_X^2} = \frac{A}{b}s_0^{1-\alpha_k(0)}\left(e^{bt_{min}} - e^{bt_{max}}\right)W^{4\alpha_i(0)-2}M_X^{2\alpha_k(0)-4\alpha_i(0)}, \tag{2.76}$$

was performed on the data, with t_{min} the minimum kinematical value of t and $t_{max} \sim p_T^2 \leq 0.04$ GeV^2. The t-dependent terms are assumed to be absorbed into a single exponential e^{bt}, where $b = b_0 - 2\alpha_i' \ln(1-z)$. The cross section shown in figure 2.18 shows an approximate independence of $d\sigma/dz$ of M_X^2 and W which leads to the expectation that $4\alpha_i(0) - 2 \sim 0$ and $2\alpha_k(0) - 4\alpha_i(0) \sim 0$, or $\alpha_i(0) \sim 0.5$ and $\alpha_k(0) \sim 1$. From the fit one indeed finds:

$$\begin{aligned} \alpha_i(0) &= 0.33 \pm 0.04 \text{ (stat) } \pm 0.04 \text{ (syst)}, \\ \alpha_k(0) &= 0.99 \pm 0.01 \text{ (stat) } \pm 0.05 \text{ (syst)}, \\ b_0 &= (3.6 \pm 0.7 \text{ (stat) } \pm 1.4 \text{ (syst)}) \text{ GeV}^{-2}. \end{aligned} \tag{2.77}$$

The result of the fit is shown by the full line in figure 2.18. The exchange between the photon and the proton can be understood as a mixture of trajectories dominated by π and $I\!R$ exchange, since the extracted value of $\alpha_i(0)$ was found between zero and one, the values of the intercept for respectively the pion and reggeon trajectory [37]. This result is also consistent with the previous H1 measurement at higher Q^2 [70]. In addition since $\alpha_k(0)$ was found to be unit, the total cross section for the scattering between the mixture of Regge trajectories α_i and the photon seemed to be dominated by pomeron exchange, as would be expected for total cross sections at these energies ($M_X^2 \in [1010, 16915]$). The value b_0 extracted from the fit implied that $b = (6.6 \pm 0.7 \text{ (stat) } \pm 1.5 \text{ (syst)}) \text{ GeV}^{-2}$ for a mean value of $z = 0.78$ which was found to be compatible with the direct measured value of ZEUS [71] for $p_T^2 < 0.5$ GeV^2.

A comparison was made with $pp \to Xp$ data from Ganguli and Roy [66], assuming vertex factorization and shown by the dashed curve in figure 2.18.a-e. The prediction at $W = 187$ GeV and its decomposition obtained from a triple Regge analysis on $pp \to Xp$ data into the three dominant triple Regge terms are shown in figure 2.18.f. The predicted photoproduction cross sections lie 25 to 65% above the measured values, which can be considered as a reasonable agreement given the simplicity of the model, the uncertainties on the couplings and the basic difference between γp- and pp-data very often described by the gap survival probability. This agreement contrasts with the discrepancies of an order of magnitude or more observed when predictions were made for diffractive hard scattering cross sections at large z in $\overline{p}p$ collisions at the Tevatron [72] using parton densities extracted from diffractive DIS at HERA [48, 49].

However very recently the ZEUS collaboration has measured the differential cross section using leading proton data [73]. Instead of measuring $d\sigma(\gamma p \to Xp)/dz$ like in the case of the H1 collaboration [69], the ZEUS collaboration measured $1/\sigma_{tot} d\sigma(\gamma p \to Xp)/dz$ leading to the advantage that most acceptance correction factors and systematic uncertainties cancel in the ratio. Figure 2.19 shows the comparison between the H1 and ZEUS measurement,therefore the differential cross section from H1 is divided by the total cross section $\sigma_{tot} = 160$ μb [42] at $W = 187$ GeV. Notice that the ZEUS measurement is performed at $\langle W \rangle = 200$ GeV, while the γp centre of mass energy from the H1 measurement is at $\langle W \rangle = 187$ GeV. The mean difference factor between both experiments is of the order of 1.9.

Non tagged proton data

Non tagged $\gamma p \to XY$ large rapidity gap data, with Y a proton or a low mass proton excitation are used in [68] to measure the differential cross section $d\sigma/dM_X^2$ at two γp



Figure 2.18: *The differential cross sections $d\sigma/dz$ as a function of W for each measured z value in the leading proton spectrometer. The data are compared with the results of a triple Regge fit and with the predictions from a triple Regge analysis on $pp \to Xp$ data. f) Shows the measured cross section as a function of z at W=187 GeV compared with the prediction derived from triple Regge analysis of $pp \to Xp$ data together with the decomposition into the three dominant terms.*



Figure 2.19: *The normalized cross section $(1/\sigma_{tot})\; d\sigma/dz$ for photoproduction data in the region $p_T^2 < 0.04$ GeV^2 is compared between the H1 and ZEUS experiment. The inner error bars indicate the statistical uncertainties, and the outer error bars are the statistical and systematical uncertainties added in quadrature.*

centre of mass energies $\langle W\rangle$=231 and 187 GeV. Those data were subjected to a triple Regge fit, taking interference terms into account, of the form:

$$\frac{d^2\sigma}{dtdM_X^2} = \frac{s_0}{W^4}\sum_{i,j,k} G_{ijk}(t)\left(\frac{W^2}{M_X^2}\right)^{\alpha_i(t)+\alpha_j(t)}\left(\frac{M_X^2}{s_0}\right)^{\alpha_k(0)}\cos[\phi_i(t)-\phi_j(t)], \qquad (2.78)$$

where $\phi_i(t)$ represents the phase of reggeon i which is determined by the signature factor. When no interference between the diffractive and non-diffractive exchanges is considered, which means $i = j$, then equation (2.78) is identical to previously mentioned triple Regge formula given in equation (2.71). The sum in the triple Regge formula is performed over only six terms given in table 6.1. Here no π-exchange was considered since they give contributions which are negligible at low M_X and become comparable to diffractive pomeron exchange

Parameter	Value		
$\alpha_{I\!R}$	0.55	± 0.10	
α_{π}	0.	$\pm 0.$	
$\alpha'_{I\!P}$	0.26	± 0.02	GeV^{-2}
$\alpha'_{I\!R}$	0.90	± 0.10	GeV^{-2}
α'_{π}	1.	± 0.50	GeV^{-2}
$b_{pI\!P}$	2.3	± 0.30	GeV^{-2}
$b_{pI\!R}$	1.0	± 1.0	GeV^{-2}
$b_{p\pi}$	1.0	± 1.0	GeV^{-2}
b_{ijk}	0.0	± 1.0	GeV^{-2}

Table 2.3: The fixed parameters in the triple Regge fit and their assumed values.

only at $M_X^2 \approx 0.03W^2$ [57].

In order to investigate the importance of the six terms a large range in W^2 and M_X^2 is required. Therefore the fixed target data from [74] with $\langle W \rangle$=12.9 and 15.3 GeV and $4 \leq M_X^2 \leq 18$ GeV^2 are added to these rapidity gap data [68] with $\langle W \rangle$=231 and 187 GeV and $0.16 \leq M_X^2 \leq 1860$, on which simultaneous fits were performed. The parameters listed in table 2.3 were fixed to the measured values from [36, 75, 44, 76], $\alpha_{I\!P}(0)$ and the four couplings, $G_{I\!PI\!PI\!P}(0)$, $G_{I\!PI\!PI\!R}(0)$,$G_{I\!RI\!RI\!P}(0)$ and $G_{I\!RI\!RI\!R}(0)$ were free fit parameters.

Three scenarios were examined in this analysis. Based on the χ^2/ndf values from each fit, a preference for either a large interference or a large isovector contribution to the subleading terms (see section 6.2.1), or else a mixture of both could be concluded. A single value of the pomeron intercept was obtained by taking the average of the three fits, giving $\alpha_{I\!P}(0) = 1.068 \pm 0.016 \pm 0.022 \pm 0.041$. The result of the fit where interference was taken into account based on the assumption of maximal coherence (see section 2.5.3) is shown in figure 2.20. No contribution from isovector exchange is taken into account by the fit. The fit represents the scenario in which the f is the dominant subleading exchange and the f and the pomeron couple similarly to the proton. The ZEUS Collaboration has also measured the photon dissociation process at $\langle W \rangle$=200 GeV using rapidity gap data [77]. As they only used their own data in the fit, a less detailed analysis was performed. The photon dissociation mass spectrum was analysed in the range $3 \leq M_X \leq 24$ GeV, assuming the presence of only $I\!PI\!PI\!P$ and $I\!PI\!PI\!R$ terms in the triple Regge sum. It was shown that the $I\!PI\!PI\!P$ term could describe the large mass spectrum, $8 \leq M_X \leq 24$ GeV with a value of the pomeron intercept, $\alpha_{I\!P}(0) = 1.12 \pm 0.04 \pm 0.08$. They also verified that a two component contribution of $I\!PI\!PI\!P$ and $I\!PI\!PI\!R$ with $\alpha_{I\!P}(0) = 1.08$ and $\alpha_{I\!R}(0)$=0.45 gave a good description of the data over the whole M_X-range. The contribution of the $I\!PI\!PI\!R$ term to the cross section over the full M_X-range was estimated to be $f_{I\!PI\!PI\!R} = 23 \pm 5$(stat) %. It should be emphasized that in the ZEUS Collaboration, the diffractive sample is obtained by subtracting from the rapidity gap sample the non-diffractive component which is modeled by the Monte Carlos EPSOFT and PYTHIA and which is in contrast with the definition of diffraction in H1.



Figure 2.20: *A triple Regge fit taking maximal constructive interference between the diffractive and non-diffractive exchange into account on the diffractive differential cross sections from [68] and [74].*

Chapter 3

The H1 experiment at HERA

H1 is one of the two general purpose detectors installed at HERA (Hadron Electron Ring Anlage) and designed to study electron-proton or positron-proton interactions. The present chapter, after a brief introduction on HERA, describes the H1 detector components with emphasis on those which are of primary importance to this analysis. An exhaustive detector description can be found in [78].

3.1 The HERA collider

The HERA machine [79] is the world's first lepton-nucleon collider. It is located in Hamburg, Germany, and has been providing luminosity to the experiments H1 and Zeus since the summer of 1992. It is schematically shown in figure 3.1, along with the pre-HERA accelerator elements.

The HERA collider is composed of two rings one for the electron (positron) and one for the proton beam storage, having a circumference of 6.3 km. Electrons are accelerated to 27.5 GeV and protons to 820 GeV [1]. Conventional magnets are used to steer the electron beam, but fields well in excess of the magnetic saturation of iron are required to constrain the 820 GeV protons. Therefore a field of 4.6 Tesla is used with a system of superconducting coils. The beams are steered in a way that they cross at zero angle at two points on the ring where collisions may take place.

The beams are bunched in small buckets (called bunches) separated 28.8 m (or 96 ns). Some bunches are left empty so that a filled electron or proton bunch might arrive at the interaction point without a corresponding colliding partner. These so called *pilot bunches* are useful for studying background events due to the collision of an electron or proton with residual gas in the beam-pipe (*beam-gas interactions*) or with the beam-pipe-wall (*beam-wall interactions*).

Two detectors, H1 located at the north and Zeus located at the south side of HERA, have been built around the interaction points. Both multi-purpose detectors provide nearly

[1] Since 1998 the HERA accelerator operates with 920 GeV protons.



Figure 3.1: *The layout of HERA is shown on the right, along with the location of the different detector halls. The pre-accelerators are shown in the blowup on the left.*

full angular coverage and are asymmetric in the beam direction, optimizing the acceptance for high multiplicity final states in the proton direction.

The electron (positron) beam can be polarized up to 70 %, by making use of the fact that over a period of time, synchrotron radiation leads to a natural transverse polarized beam. A third experiment, HERMES [80] located on the east side of HERA, makes use of this polarized beam by colliding it with a polarized proton gas jet to study the spin structure of the proton. A fourth experiment, HERA-B [81] located on the west side of HERA, was approved in February 1995, primarily to search for CP violation in decays of B mesons.

The most important HERA parameters are given in table 3.1, showing those used in 1997, the year in which the data for the present analysis were collected.

3.2 General description of the H1 detector

The H1 detector depicted in figure 3.2 is designed to study many aspects of ep scattering and therefore consists of a number of subsystems providing complementary information, from which the properties of the collision processes can be reconstructed. In order to exploit fully the physics potential, the detector has to fulfil a number of major roles, which are:

- **Electron detection:** identification and energy measurement of scattered electrons is vital for the study of neutral current DIS events as well as for photoproduction events, where the scattered electrons are at very low Q^2 values.

Parameter	*Unity*	*1997* e^+/p
Circumference	m	6336
Interaction points		4
Max. number of bunch slots		220/220
Number of colliding bunches		174
Number of e-pilot bunches		15
Number of p-pilot bunches		10
Number of empty bunches		21
Nominal energies	GeV	27.55/820
Mean Currents	mA	40/100
Package separation	ns	96
σ_x *at IP*	μm	179
σ_y *at IP*	μm	48
σ_z^{proton} *at IP*	mm	200
$\mathcal{L}_{inst}$	cm^{-2}s^{-1}	1.4 x 10^{31}

Table 3.1: The main HERA parameters.

- **Hadron detection:** calorimetry with good resolution and fine granularity is needed for characterizing the hadronic final state in many processes. The calorimeter must be hermetic in order to measure the transverse energy balance while for charged current DIS events, the undetected final state neutrino exhibits an imbalance.

- **Charged particle tracking:** charged particle momentum measurement and identification is required for the investigation of many processes at HERA. This is provided by tracking detectors covering a large solid angle. A magnetic field curves the tracks and allows the momentum to be determined, while energy loss measurements enable particle identification.

- **Muon identification:** heavy flavor, rare Standard Model and exotic physics processes are detected through the production of muons in the final state.

- **Luminosity measurement:** accurate determination of the luminosity delivered to the experiment is vital for calculation of cross sections.

- **Triggering and readout:** on the basis of the fast information coming from the different detector subsystems, a selection of *ep* interactions of physics interest has to be done by the trigger system.

In these aspects the H1 detector does not differ strongly from the detectors at e^+e^--colliders, or *pp*-colliders built in the past. The principal H1 detector parameters are given in table 3.2. Specific at HERA are however the imbalance in the energy of the two colliding beams, which requires an asymmetric detector. Different also is the micro-structure of the