Prompt Photon Production at HERA

DISSERTATION

zur Erlangung des Doktorgrades
des Fachbereichs Physik
der Universität Hamburg

vorgelegt von

Rachid Lemrani-Alaoui

aus Frankreich

Hamburg
2003

Gutachter der Dissertation:	Prof. Dr. F.-W. Büsser Prof. Dr. B. Naroska
Gutachter der Disputation:	Prof. Dr. F.-W. Büsser Prof. Dr. G. Heinzelmann
Datum der Disputation:	25.02.2003
Vorsitzender des Prüfungsausschusses:	Dr. H.D. Rüter
Vorsitzender des Promotionsausschusses:	Prof. Dr. G. Huber
Dekan des Fachbereichs Physik	Prof. Dr. F.-W. Büsser

Abstract

Inclusive prompt photons and prompt photons together with a jet are measured in photoproduction at HERA. Cross sections are presented as a function of transverse energy E_T^{γ} and pseudorapidity η^{γ} for $E_T^{\gamma} > 5$ GeV and $-1 < \eta^{\gamma} < 0.9$ in the inelasticity range $0.2 < y < 0.7$ corresponding to γp center of mass energies $140 < W < 270$ GeV. For the prompt photons + jet analysis a jet was required in addition with $E_T^{jet} > 4.5$ GeV and $-1.5 < \eta^{jet} < 2.3$. The data were taken with the H1 detector in the years 1996-2000 using an integrated luminosity of 105 pb^{-1}. After selection of prompt photon candidates, a sizable amount of background from π^0 mesons remains and the γ signal is extracted by a likelihood technique using calorimetric shower shape variables. The results are compared to the PYTHIA and HERWIG event generators and to a NLO-QCD calculation. PYTHIA and HERWIG predictions describes the data well in shape, are however low in normalisation. The NLO-QCD calculation describes the data quite well in the presented η^{γ} and E_T^{γ} ranges. It has however a tendency to overshoot the data especially at large η^{γ} which could be explained by multiple interactions.

Kurzfassung

Es wird eine Messung zur Photoproduktion von prompten Photonen bei HERA vorgestellt. Die Photonen wurden dabei sowohl inklusiv als auch zusammen mit einem Jet nachgewiesen. Erzeugungsquerschnitte werden als Funktion der transversalen Energie E_T^{γ} und der Pseudorapidität η^{γ} für $E_T^{\gamma} > 5$ GeV und $-1 < \eta^{\gamma} < 0.9$ im Unelastizitätsbereich $0.2 < y < 0.7$ presentiert. Dieser Bereich entspricht Schwerpunktsenergien W im Photon-Proton System von $140 < W < 270$ GeV. Im Fall der nichtinklusiven Produktion prompter Photonen wurde zusätzlich ein Jet mit $E_T^{jet} > 4.5$ GeV und $-1.5 < \eta^{jet} < 2.3$ gefordert. Die Daten wurden mit der H1 Detektor in den Jahren 1996-2000 registriert, wobei eine integrierte Luminosität von 105 pb^{-1} benutzt wurde. Die Auswahl der Ereignisse mit Kandidaten prompter Photonen enthält noch einen erheblichen durch π^0-Mesonen verursachten Untergrundanteil. Der Anteil prompter Photonen wird durch eine *Maximum-Likelihood*-Technik unter Verwendung von Variablen bestimmt, welche die im Kalorimeter gemessenen Schauerformen beschreiben. Die Ergebnisse werden mit den Ereignisgeneratoren PYTHIA und HERWIG und einer NLO-QCD-Rechnung verglichen. Die Voraussagen von PYTHIA und HERWIG beschreiben zwar gut die relativen Abhängikeiten der Querschnitte, sie sind aber absolut zu niedrig. Die NLO-QCD-Rechnung beschreibt die Ergebnisse in den untersuchten η^{γ} und E_T^{γ} Bereichen recht gut, allerdings mit einer Tendenz, vor allem bei großen η^{γ}, etwas über den Messungen zu liegen. Dies kann durch partonische Mehrfachwechselwirkungen erklärt werden.

To my brother Lahcen

Contents

Introduction

The standard model, a theory completed in the beginning of the seventies, has been very successful in describing all the collected data in high energy physics. However it is expected that a new theory is needed at the higher energies which will be reached during the next decade. A few aspects of the standard model based on a quantum gauge field theory are described here. The constituents of all the existing matter are elementary particles called fermions. They interact with each other through the electromagnetic, the weak, the strong and the gravitational force. The electromagnetic and weak forces have been unified into the electroweak force. These interactions take place by exchange between fermions of so-called bosons: photon, weak bosons (W^+,W^- and Z) for the electroweak force, and eight gluons for the strong force. The strength of a force is given by the coupling constants, the fine structure constant α for the electroweak and α_s for the strong force. The gravitational force can be neglected in high energy physics and is not part of the standard model. The fermions are classified in two categories: the six quarks which are sensitive to electroweak and strong forces and the six leptons which are only sensitive to electroweak forces. The fermions are classified in three generations of two quarks ($up, down$), ($charm, strange$), ($top, bottom$) and of two leptons ($electron, electron\ neutrino\ \nu_e$), ($muon, muon\ neutrino\ \nu_\mu$,) ($tau, tau\ neutrino\ \nu_\tau$). The mass of the particles is due to a coupling to the Higgs boson, the only still undiscovered particle of the standard model. A few parameters (e.g coupling constants, mixing angles, masses) are needed for the theory and are being measured by the experiments.

Physicists use high energy machines to probe matter at small scales and to produce new particles. In scattering experiments either beams of particles are scattered against fixed targets (fixed target experiments) or two high energetic particle beams are brought to collide with eachother. The proton appears to be formed of quarks and gluons at high energies, and many other particles (the hadrons) composed of quarks and gluons have been observed. Nowdays the proton is being probed at the HERA collider at scale a thousand times smaller than its own size by using beams of electrons and protons. Higher energies allow also to produce new particles and all the fermions of the standard model have indeed been observed starting with the electron in 1897 up to the tau neutrino hundred years later.

After a collision the properties (observables) of the produced particles are measured in a detector. To a first approximation a pair of elementary particles from the colliding beams interact with each other. However a lot of complications arise due to the composite structure of the colliding particles and many other interactions and processes take place. For instance produced quarks and gluons are not observed directly. Instead, quarks and gluons produce jets of hadrons following a process called hadronisation. The hadronisation of quarks and gluons is extremely difficult to describe using the standard model and is not fully understood theoretically. However models have been developed and they are useful to reconstruct the properties of the quarks and gluons themselves from the observation of the jets. The observation of a photon in the final state is a cleaner measurement because it is less sensitive to the hadronisation models. These so-called prompt photons are produced through the interaction of the elementary particles of the beams.

In chapter 1 *ep* physics a HERA is shortly discussed, in particular deep inelastic scattering and photo-production. Chapter 2 introduces theoretical aspects of the prompt photon production and gives some previous experimental results. In chapter 3 the employed Monte-Carlo event generators are presented and the produced Monte-Carlo samples are listed. Chapter 4 describes briefly the HERA machine and the H1 detector, in particular the components relevant for this analysis. In chapter 5 the photon and jet reconstruction are explained. Chapter 6 describes the event selection. In chapter 7 the signal and background separation is presented. In chapter 8 extraction of cross-sections and the systematic errors are discussed. In chapter 9 a comparison of the inclusive prompt photon analysis to the theory is presented and further results from the prompt-photon + jet analysis are discussed. In chapter 10 the presented analysis and the results are summarized.

Chapter 1

Theoretical framework

Prompt photon production as discussed in this report is a special case of ep scattering at HERA. In this chapter the basic concepts of ep scattering are presented, in particular deep inelastic scattering (DIS) and photo-production.

1.1 ep scattering

The main process studied at HERA is deep inelastic scattering (DIS) of electrons and protons. It is shortly described here to define the relevant kinematics and the concept of structure functions.

1.1.1 Kinematics



Figure 1.1: *Diagrams of electron and proton scattering via photon and Z exchange for neutral current interactions and W exchange for charged current interactions. The four-momentum vectors of the particles are given in parentheses.*

Figure 1.1 shows the DIS processes where an electron (e) interacts with a proton (p). In neutral current (NC) interactions ($e^{\pm}p \rightarrow e^{\pm}X$) a neutral boson ($\gamma, Z$) is exchanged leading in the final state to a scattered electron and the hadronic final state X (break-up of the proton). In charged current (CC) interactions ($e^- p \rightarrow \nu_e X$, $e^+ p \rightarrow \overline{\nu}_e X$) a charged boson (W^-, W^+) is exchanged with a neutrino ν in the final state instead of the scattered electron.

We define k and k' as the incident and scattered lepton four-momenta, p as the proton four-momentum and $q = k - k'$ as the momentum transfer. The process is determined by the 3 Lorentz invariant variables:

$$\text{Four-momentum transfer squared:} \quad Q^2 = -q^2 = -(k-k')^2 \;, \quad Q^2 \in [0, s] \tag{1.1}$$

$$\text{Björken scaling variable:} \quad x = \frac{Q^2}{2p \cdot q} \;, \quad x \in [0, 1] \tag{1.2}$$

$$\text{Inelasticity:} \quad y = \frac{p \cdot q}{p \cdot k} \;, \quad y \in [0, 1] \tag{1.3}$$

Neglecting the masses of the electron and of the proton the 3 variables are related by:

$$Q^2 = s \cdot x \cdot y \tag{1.4}$$

where $s = (k + p)^2$ is the total invariant energy squared.
The invariant mass W of the hadronic final state is given by:

$$W^2 = (p + q)^2 \,. \tag{1.5}$$

1.1.2 Structure functions

The DIS cross-section for the photon exchange is given by:

$$\frac{d^2\sigma^{ep}(x, Q^2)}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4} \left[\frac{y^2}{2} 2xF_1(x, Q^2) + (1-y)F_2(x, Q^2) \right] \tag{1.6}$$

where α is fine structure constant and F_1 and F_2 are the structure functions of the proton as a function of x and Q^2. The contributions of Z^0 and $W^{\pm}$ exchanges is small at low Q^2: $Q^2 << M^2_{Z^0}, M^2_W$. This is due to the propagator terms of the form $1/(Q^2 + M^2_{Z^0})$ respectively $1/(Q^2 + M^2_W)$ for the weak exchange. Using the longitudinal structure function $F_L = F_2 - 2xF_1$ the cross-section can be written as:

$$\frac{d^2\sigma^{ep}(x, Q^2)}{dx dQ^2} = \frac{2\pi\alpha^2}{xQ^4} \left[(1 + (1-y)^2)F_2(x, Q^2) - y^2 F_L(x, Q^2) \right] \tag{1.7}$$

In most of the phase space at HERA the differential cross-section is dominated by F_2. Only at large y there is a sizable effect of F_L on the cross-section.

1.2 The naive quark parton model

The structure functions defined in the previous section are related in the quark-parton model (QPM) to the quark distributions in the proton [1]. It is assumed that the proton is a stream of parallel partons i with electric charges e_i which can be described by densities $q_i(x)$. The structure function is then given by: $F_2(x,Q^2) = F_2(x) = \Sigma e_i^2\, x\, q_i(x)$.

For spin $\frac{1}{2}\hbar$ quarks the cross-section for longitudinally polarized photons is zero (Callan-Gross relation [2]): $2xF_1(x) = F_2(x)$

The structure functions do not depend on Q^2. This so-called scaling behavior, predicted by Björken for large Q^2 and large $1/x$, is fulfilled in the QPM.

The QPM approximation holds for protons with very high momenta (infinite momentum frame). The proton is then almost flat due to Lorentz contraction and the interaction time is very small due to time dilatation. In this approximation the quarks of the proton are far from each-other and have no time to interact which each-other. The Björken variable x is in this model equal to the proton momentum fraction carried by struck quark.



Figure 1.2: *Picture of a Lorentz contracted proton with momentum P. Inside the proton the partons i with momentum fraction x_iP are almost free during the interaction. The contraction factor is given by the inverse Lorentz factor* $1/\gamma$.

1.2.1 The QCD Improved Parton Model

That the QPM is insufficient got first apparent by the observation that the momentum sum : $\Sigma \int_0^1 dx q_i(x) x < 1$ (about one half of the momentum is missing) [3].

The theory of strong interaction, Quantum Chromodynamique (QCD), introduces also neutral partons in the proton: the gluons. In QCD, interactions of partons take place through gluons by gluon radiation from quarks and gluon splitting in $q\overline{q}$ pairs. This leads to a sea of quarks and gluons in addition to the valence quarks (see Figure 1.3). The Björken scaling is now broken leading to a decrease of $F_2(x,Q^2)$ at large Q^2 and large x and an increase at small x as indicated in Figure 1.4. This can simply be explained by a depletion of high x quarks due to gluon radiation at large Q^2, where smaller structures can be resolved and a corresponding accumulation of sea quarks at small x. Another QCD effect is that the quarks acquire transverse momenta from gluon radiation leading to a non-zero longitudinal structure function F_L.



Figure 1.3: *The different models for the proton. At low energies the proton behaves as a point-like particle (a). (b) shows the three free valence quarks in the proton. In (c) the valence quarks are interacting via gluons. In (d) a sea of quarks and gluons appears due to QCD effects like gluon radiation from quarks and gluon splitting in $q\overline{q}$ pairs.*



Figure 1.4: *Sketch of the variation of F_2 as a function of x and Q^2.*

1.2.2 The evolution of Parton Distributions

The partons distributions cannot yet be calculated perturbatively. The Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) equations [4–7] give though the evolution of the parton distributions $q_i(x)$ and $g(x)$ with Q^2 (eqs. 1.8,1.9). Thus for known parton distributions at scale Q_0^2 one can derive the parton distributions at scale Q^2.



Figure 1.5: *Picture of a proton probed by photons of different virtualities Q_0 and $Q > Q_0$. As the virtuality increases smaller structure are resolved. A quark appearing as point-like at virtuality Q_0, may appear to have radiated gluons at higher virtuality Q.*

$$\frac{dq_i(x,Q^2)}{dlnQ^2} = \frac{\alpha(Q^2)}{2\pi}\int_x^1 \frac{dy}{y}\left[\sum_j q_j(y,Q^2)P_{q_j\to q_i}\left(\frac{x}{y}\right) + g(y,Q^2)P_{g\to q_i}\left(\frac{x}{y}\right)\right] \tag{1.8}$$

$$\frac{dg(x,Q^2)}{dlnQ^2} = \frac{\alpha(Q^2)}{2\pi}\int_x^1 \frac{dy}{y}\left[\sum_j q_j(y,Q^2)P_{q_j\to g}\left(\frac{x}{y}\right) + g(y,Q^2)P_{g\to g}\left(\frac{x}{y}\right)\right] \tag{1.9}$$

The splitting functions $P_{j\to i}$ reflect for instance that a quark of momentum x can be the product of a quark of momentum $y > x$ radiating a gluon of momentum $y - x$ or of the splitting of a gluon. These functions give the probabilities of the leading-order diagrams of figure 1.6 . They have also been computed at next-to-leading order [8].



Figure 1.6: *The probability of the shown leading order diagrams is given by the splitting functions $P_{q\to q}$, $P_{q\to g}$, $P_{g\to q}$, $P_{g\to g}$.*

1.3 The structure of the Photon

In the standard model the photon is the gauge boson for the electromagnetic force. The photon is an elementary particle. However it can fluctuate into virtual pairs of charged leptons or quarks for a short time according to the quantum uncertainty principle.

For a photon of high energy the quark-antiquark pair can develop into a quark-gluon cascade leading to a complicated partonic object with the same quantum numbers as the photon. This development is not calculable in perturbative QCD and models using parametrisations constrained by measurements have been developed.

At high Q^2 the photon has no time to fluctuate: it behaves as a point-like particle. But when the time of interaction is much smaller than the fluctuation time, the photon behaves as a hadronic object and there are indeed similarities in photon-proton and hadron-proton interactions.

The concept of hadronic structure for the photon was originally introduced in the Vector-Dominance-Model (VDM) [9]. It considers the photon as a superposition of vector-mesons with the same quantum numbers i.e. in terms of vector-meson wave-functions:

$$|\gamma>= \sum_{V=\rho,\omega,\phi,...} c_V|V> . \tag{1.10}$$

The VDM works well at low γp center-of-mass energies but is insufficient at high energies. In general three components are then considered as illustrated in Fig 1.7: the bare component where the photon doesn't fluctuate, the anomalous or point-like component were pQCD can be applied, and the non-perturbative VDM component:

$$\sigma^{\gamma p} = \sigma^{\gamma p}_{dir} + \sigma^{\gamma p}_{anom} + \sigma^{\gamma p}_{VDM} \tag{1.11}$$



Figure 1.7: *The three components of the photon structure: the bare component where the photon interacts directly, the anomalous component where the photon undergoes perturbative QCD processes and the vector-meson component where the photon fluctuates into a complex hadronic object that cannot be calculated by perturbative QCD.*

It is possible to define a structure function F_2 for the photon based on its parton content in analogy to the proton:

$$F_2^\gamma(x,Q^2) = \Sigma e_i^2 (xq_i^\gamma(x,Q^2) + x\overline{q}_i^\gamma(x,Q^2)) \quad (1.12)$$

It describes the distribution of electric charge of the photon structure and can be constrained at e^+e^- experiments in DIS-like $\gamma^*\gamma$ reactions: the almost real γ, the probed photon, is radiated by the electron (or positron) scattered at small angle and the highly virtual photon γ^*, the probe, is radiated by the electron scattered at large angle. DORIS, PETRA, PEP, TRISTAN, LEP experiments contributed to the measurement of F_2^γ.

Similarly to the proton case, the parton density functions (pdfs) in the photon and DGLAP evolution equations are introduced. An additional term $a(x) = 3e_i^2 \frac{\alpha}{2\pi}\left[x^2 + (1-x)^2\right]$ enters the equation 1.8 to take into account the point-like component $\gamma \to q\overline{q}$. It is referred to as the 'inhomogeneous' term and makes F_2^γ rise at large x. The available models for the structure of the photon introduce pdfs at a low scale Q_0 depending on parameters. Fits of the pdfs to the measured F_2^γ are performed using DGLAP equations at LO or NLO to evolve the structure function in Q^2.

The existing data on F_2^γ do not constrain much the gluon density in the photon. As for the proton, F_2^γ is directly related to the quark content and only indirectly to the gluon content. The data on F_2^γ are less precise than in the proton case and there is no momentum sum rule constraining the photon pdfs.

In jet production in $\gamma\gamma$ and and γp scattering there is sensitivity to the gluon density of the photon and AMY, TOPAZ, TRISTAN $\gamma\gamma$ experiments already provided evidence for a non-zero gluon component ($qg \to qg$). At HERA more precise results were achieved.

1.4 Photo-production at HERA

The ep cross-section decreases strongly with Q^2 ($1/Q^2$ propagator term). For $Q^2 \approx 0$ the exchanged photons are almost real. We can then consider these events as γp scattering and speak of photo-production events. The electron behaves as a source of photons with energies in the range from zero up to the HERA electron energy of 27.5 GeV. The spectrum of the photon energies integrated to a maximal Q^2, Q^2_{max}, is given in the Weisz*ä*cker-Williams approximation (WWA) [10] by:

$$f_{\gamma/e}(y) = \frac{\alpha}{2\pi}\left(\frac{1+(1-y)^2}{y}\right) ln\left(\frac{Q^2_{max}(1-y)}{m_e^2 y^2}\right) \quad (1.13)$$

Hard photo-production events contain particles with high transverse momentum P_T: a so-called hard interaction took place. The variable P_T can provide the renormalization scale for perturbative QCD calculations, replacing Q^2 which is used in DIS but is too small here. The factorization scale provides the frontier between the regions where perturbative QCD can be applied and the non-perturbative regions where the evolution ladder is described by the parton distributions.

There are two classes of photo-production. In the direct process the exchanged photon couples directly to the quarks of the proton (Fig. 1.8). In the resolved process the photon fluctuates into a hadronic state as discussed in section 1.3 and a parton of the photon enters the hard interaction (Fig. 1.9). In this case there is a photon remnant which is not participating in the hard interaction, similar as in the case of the proton.



Figure 1.8: *Feynman graphs of direct di-jets photo-production events*



Figure 1.9: *Feynman graphs of resolved di-jets photo-production events*

Chapter 2

Prompt photon production

2.1 Prompt Photon Production in γp reactions

Prompt photon events have a photon in the final state produced in a hard interaction (see section 1.4), i.e. the experimental signature is a photon with substantial P_T. These photons have to be distinguished from photons produced via particle decays and, in *ep* reactions, also from photons emitted by the incident or scattered electrons. Figure 2.1 and 2.2 show examples of prompt photon diagrams at leading order with the direct component (Born diagram)

$$q\gamma \to q\gamma$$

and the resolved components

$$q^{\gamma}g^{p} \to \gamma q$$
$$g^{\gamma}q^{p} \to \gamma q$$
$$q\overline{q} \to \gamma g \ .$$

Parton fragmentation can also lead to a photon in the final state as shown in Figure 2.3(b)(d). For example the following di-jets processes have to be included when a prompt photon is appearing in the final state:

$$\gamma g \to q\overline{q}$$
$$\gamma q \to qg$$

The contributions of these different processes to the total prompt photon cross-section are presented in section 9.2.

Prompt photon measurements are free from some of the complications encountered in jet measurements such as the non perturbative hadronisation process, the jet identification and the jet energy measurement. The photon is directly measured in the detector. On the other hand the cross-section for prompt photon production is much smaller. The electromagnetic coupling constant α enters in the diagrams instead of the strong coupling constant α_s as for the jets.

The main experimental difficulty in the prompt photon measurement is a large background from neutral mesons.

Prompt photon production is sensitive to the parton structure of the proton and due to the resolved processes (see Fig. 2.2) also to the parton content of the photon. In particular it is directly sensitive to the gluon content of the photon, however with the presently available integrated luminosities this cannot seriously be constrained.

At HERA the gluon content of the photon can also be studied in the jet production analyses (see Fig. 1.8, 1.9) and in J/ψ production analyses in the reactions $\gamma p \to$ jets X and $\gamma p \to J/\psi X$.



Figure 2.1: *Example of direct prompt photon processes at leading order. The photon couples directly to a parton from the proton.*



Figure 2.2: *Examples of resolved prompt photon processes at leading order. The photon is resolved and a parton from the photon couples to a parton from the proton.*



Figure 2.3: *Diagrams of the prompt photon production. The direct-direct (a), direct-fragmentation (b), resolved-direct (c) and resolved-fragmentation (d) processes are distinguished. The functions in the circles are defined in the text.*

The prompt photon cross-sections are given in leading order following [11] (see Fig. 2.3) by:

$$d\sigma_{dir}^{nonfrag} = \Sigma f_{a/p}(x_p, Q^2) \otimes f_{\gamma/e}(y) \otimes \sigma_{a\gamma\to\gamma+X} \tag{2.1}$$

$$d\sigma_{dir}^{frag} = \Sigma f_{a/p}(x_p, Q^2) \otimes f_{\gamma/e}(y) \otimes \sigma_{a\gamma\to d+X} \otimes D_{\gamma/d}(z, Q^2) \tag{2.2}$$

$$d\sigma_{res}^{nonfrag} = \Sigma f_{a/p}(x_p, Q^2) \otimes f_{b/e}(y) \otimes \sigma_{ab\to\gamma+X} \tag{2.3}$$

$$d\sigma_{res}^{frag} = \Sigma f_{a/p}(x_p, Q^2) \otimes f_{b/e}(x_e, Q^2) \otimes \sigma_{ab\to d+X} \otimes D_{\gamma/d}(z, Q^2) \tag{2.4}$$

Here $f_{a/p}(x_p, Q^2)$ is the probability that the proton p will produce a parton a carrying a fraction x_p of the proton's momentum.

The parton fragmentation function $D_{\gamma/d}(z, Q^2)$ gives the probability that the parton d will produce a photon γ carrying a fraction z of the parton's momentum.

The flux $f_{\gamma/e}(y)$ of the photon radiated by the electron e is given in the Weisz*ä*cker-Williams approximation (see section 1.4) by

$$f_{\gamma/e}(y) = \frac{\alpha}{2\pi}\left(\frac{1+(1-y)^2}{y}\right) ln\left(\frac{Q^2_{max}(1-y)}{m_e^2 y^2}\right) . \tag{2.5}$$

The probability $f_{b/e}(x_e, Q^2)$ that an electron e will produce a parton b carrying a fraction x_γ of the photon's momentum is:

$$f_{b/e}(x_e, Q^2) = \int_{x_e}^{1} \frac{dy}{y} f_{\gamma/e}(x_\gamma, Q^2) f_{b/\gamma}(x_\gamma, Q^2) \; . \tag{2.6}$$

Several NLO calculations for prompt photon production at HERA have been performed [12–14]. The most recent calculation [14] includes NLO corrections to all the leading order diagrams of figure 2.4: direct non-fragmentation (a), resolved non-fragmentation (b), direct fragmentation (c), resolved fragmentation (d). The box term also shown in figure 2.4 (e) is of order $\alpha^2\alpha_s^2$. The direct processes are of order $O(\alpha^2)$ (two electromagnetic vertices). The resolved processes are of the same order $O(\alpha^2)$ the hard subprocess being of order $O(\alpha\alpha_s)$ and the photon structure function of order $O(\alpha/\alpha_s)$ The fragmentation processes are also of the same order $O(\alpha^2)$ because the fragmentation function are of order $O(\alpha/\alpha_s)$. In the analysis discussed below the contribution of fragmentation is strongly suppressed by an isolation requirement for the photon (see section 6.4)



Figure 2.4: *Diagrams of the prompt photon production. The direct-direct (a), resolved-direct (b), direct-fragmentation (c) and resolved-fragmentation (d) processes are distinguished. The box diagram (e) of order* $\alpha^2\alpha_s^2$ *is also shown*

2.2 Previous experimental results

2.2.1 Prompt Photon at HERA

The latest published results from the ZEUS experiment were based on an integrated luminosity of 38.4 pb^{-1} using 1996 and 1997 data [15–18]. The cross-section is given as a function of the transverse energy E_T^γ (Fig. 2.5) and the pseudo-rapidity[1] η^γ (Fig. 2.6) for $E_T^\gamma > 5$ GeV and $-0.7 < \eta^\gamma < 0.9$ in the γp center-of-mass energy range 134-285 GeV. Comparisons were made with Monte-Carlo models based on leading order matrix elements and leading logarithm parton showers and also with next-to-leading order QCD calculations. Good agreement is observed for the E_T^γ distribution but for negative η^γ all predictions are below the data.



Figure 2.5: *ZEUS E_T^γ distribution of the prompt photons in the phase space discussed in the text. The data points have been compared to the Monte-Carlo programs PYTHIA and HERWIG (see chapter 3) and next-to-leading order calculations.*

The H1 experiment presented preliminary results [19] based on 7.9 pb^{-1} taken in 1996. A cross-section for prompt photon photo-production with $E_T^\gamma > 5$ GeV and $-1.2 < \eta^\gamma < 1.6$ of 104.8±5.9±15.7 pb was measured in agreement with QCD prediction of 84 pb.

[1]The pseudo-rapidity η is defined as $-ln\left(tan\left(\frac{\theta}{2}\right)\right)$, where θ is the polar angle with respect to the proton beam direction.



Figure 2.6: *ZEUS* η^γ *distribution of the prompt photons in the phase space discussed in the text. The data points have been compared to the Monte-Carlo programs PYTHIA and HERWIG (see chapter 3) and next-to-leading order calculations.*

2.2.2 Prompt Photons at other experiments

Table 2.1 shows a list of publications on prompt photon analyses. They have contributed to a better understanding of QCD.

In 1983 the E629 experiment at Fermilab provided an observation of prompt photons [23]. In 1985, NA14 disfavoured the gauge-integer-charge-quark model [24]. In 1988 NA24 [26] and WA70 [28] found evidence for the valence-quark-antiquark annihilation process. In 1998, OPAL [55] measured the quark-to-photon fragmentation function. Many prompt photon measurements helped constraining the gluon content of the proton in the early nineties, but recent results in many hadronic colliders cannot be explained by the present perturbative NLO QCD calculations (see e.g. [50, 51]).

Figure 2.7 shows a compilation [51] of results from hadronic experiments (i.e. using hadron beams). The results are presented as a ratio data/theory as a function of the reduced variable $x_T = 2E_T^\gamma/\sqrt{s}$. The fixed target experiments at CERN are WA70 analyzing the reactions $\pi^+ p$ and $\pi^- p$ and UA6 analyzing the reactions pp and $\overline{p}p$. E706 is a $\pi^- Be$ fixed target experiment at Fermilab. The pp collider experiments at CERN are R806, R807, R110 and CDF and DØ are $\overline{p}p$ experiments at the Tevatron. All experiments show the same feature. The x_T distribution predicted by the NLO calculation is less steep than the measurements.

This discrepancy may be explained by a large intrinsic transverse momentum k_t of the partons in the proton. The partons acquire this transverse momentum through radiation of soft gluons.

Experiments	Year	Reaction	Energy (GeV)	Location
R412	'76 [20]	pp	$\sqrt{s}$=45,53	CERN ISR
R107	'78 [21]	pp	$\sqrt{=}s53$	CERN ISR
R806	'82 [22]	pp	$\sqrt{s}$=31,45,53,63	CERN ISR
E629	'83 [23]	pC, π^+C	E_{beam}=200	Fermilab FNAL
NA14	'85 [24]	γLi	E_{beam}=50-150	CERN SPS
NA3	'86 [25]	pC, π^+C, π^-C	$\sqrt{s}$=19.4	CERN
NA24	'87 [26]	π^+C, π^-C	E_{beam}=300	CERN
WA70	'88 [27,28]	pH, π^+C, π^-C	E_{beam}=280	CERN SPS
UA1	'88 [29]	$p\overline{p}$	$\sqrt{s}$=546,630	CERN $Sp\overline{p}S$
UA2	'91,'92,'93 [30–32]	$p\overline{p}$	$\sqrt{s}$=546, 630	CERN $Sp\overline{p}S$
UA6	'93,'98 [33,34]	pp $p\overline{p}$	$\sqrt{s}$=24	CERN
CDF	'92,'93,'94,'02 [35–39]	$p\overline{p}$	$\sqrt{s}$= 630, 1800	Fermilab Tevatron
D0	'96,'00 [40,41]	$p\overline{p}$	$\sqrt{s}$=630, 1800	Fermilab Tevatron
E706	'99 [42]	pBe	$\sqrt{s}$=32,39	Fermilab
ALEPH	'91,'96 [43,44]	e^+e^-	$\sqrt{s}$=100	CERN LEP
DELPHI	'92 [45]	e^+e^-	$\sqrt{s}$=100	CERN LEP
L3	92 [46]	e^+e^-	$\sqrt{s}$=100	CERN LEP
OPAL	'90,'91,'98 [47,48,55]	e^+e^-	$\sqrt{s}$=100	CERN LEP
H1	'97 [19]	ep	$\sqrt{s}$=300	DESY HERA
ZEUS	'97,'00,'01 [15,16,49]	ep	$\sqrt{s}$=300	DESY HERA

Table 2.1: *List of experiments on prompt photon production, year of publication, the reaction studied, the center-of-mass energy $\sqrt{s}$ or the beam energy for fixed target experiments, and the location.*

Figure 2.8 shows how agreement with the data is achieved by introducing an ad-hoc higher $< k_T >$. The value of $< k_T >$ is fitted to give the best possible description.

The so-called resummed calculations [52, 53] try to compensate for the lack of real gluon radiation with respect to virtual gluon radiation arising from the energy cut-offs used for the numerical calculations. These resummed calculations are less scale dependent than NLO theory, but fail to describe the data. A more recent resummed calculation [54] taking into account the recoil of the parton after soft gluon radiation gives better results (Fig. 2.9). This subject is still under investigation.

The production of prompt photons was also studied at LEP by all four experiments. OPAL gave the first measurement of the quark-to-photon fragmentation function through the inclusive production of prompt photons in hadronic Z^0 decay [55].



Figure 2.7: *The data cross-section divided by the NLO prediction is shown as a function of* $X_T = \frac{2E_T^\gamma}{\sqrt{s}}$ *for different prompt photon experiments.*



Figure 2.8: *E706 prompt photon and π_0 inclusive cross-sections as a function of E_T for* 530 *GeV/c proton-nucleon interaction. It is compared to a NLO prediction without $< k_T >$ (dotted) and with $< k_T >$ enhancement (dashed). A value of $< k_T >=$* 1.2 *GeV for the prompt photon and* 1.3 *GeV for π^0 data applied to the NLO predictions improves the data description.*



Figure 2.9: *NLO calculation compared to E706 prompt photon results. The dotted lines represents the bare NLO calculation, while the dashed line incorporates resummation calculation and the full line both resummation calculation and recoil of the partons.*

Chapter 3

Monte-Carlo simulation

3.1 Event Generators

The Monte-Carlo event generators used in this analysis simulate leading order hard sub-processes calculated by perturbative QCD and treat higher order effects by leading-logarithmic parton-showers as sketched in figure 3.1. They also include non-perturbative physics by the use of hadronisation models discussed below and parton density functions.

The events provided by the Monte-Carlo generators are put through the full H1-detector simulation. They are used to take account of the detector effects like acceptance and kinematic smearings and to determine the correction factors to be applied to the data in order to obtain the cross-sections at the hadron level.



Figure 3.1: *The different components of a Monte-Carlo event generator: the hard subprocess, the partons showers and the hadronisation.*

3.1.1 Treatment of hadronisation

Because of colour confinement partons are not detected directly. Thus only colourless hadrons are observed in the final state and enter the detectors. In a first stage a shower of partons is produced using the Leading-Log approximation of QCD. The resulting partons are in a second stage converted into hadrons using non-perturbative models.

Lund string model

A simplified description of the Lund string model [56] is given. The coloured partons from parton showers are connected by a colour flux along a thin tube or a string of energy density $\sim 1\ GeV\ / fm$. As the partons move apart the string is stretched and its potential energy is increased. When the potential energy of the string is high enough to produce a $q\overline{q}$ pair it breaks up into two strings. If the energy of the fragments is high enough the process can be repeated and new $q\overline{q}$ pairs are formed. The process is repeated until only on-shell colourless hadrons remain. The figure 3.2 shows a simplified sketch of these different steps. The production of baryons is still poorly understood. In the simplest approach a string is also allowed to break into a antidiquark-diquark pair.

The PYTHIA event generator [58] (see section 3.2) uses the Lund string model implemented in the JETSET program [57].



Figure 3.2: *Simplified sketch of the breaking of a string into mesons used in the lund hadronisation model.*

Cluster model

In the cluster fragmentation model [64] the final state gluons obtained after parton showering are split into antiquark-quark pairs or antidiquark-diquark pairs. Close enough quarks, diquarks, antiquarks or antidiquarks are grouped to form colourless clusters . Heavy clusters are split into lighter clusters before all clusters decay into colourless hadrons. The cluster model is used in HERWIG [64] (see section 3.2) .

3.1.2 Multiple interactions

Multiple interactions are implemented into the Monte-Carlo generators to take into account events where several parton-pairs (from the proton and the photon) undergo hard interactions. Experiments have shown that multiple interactions give rise to some energy flow that cannot be neglected e.g. [60]. In PYTHIA independent parton-parton scattering below the transverse momentum of the main partonic scatter are allowed down to some limit (1.45 GeV here). In HERWIG, multiple interactions are simulated by adding low P_T hadrons (the so-called soft underlying event SUE) to a fraction of the events. It has been tuned to the UA5 experimental results [64].

3.2 PYTHIA and HERWIG Monte Carlo Samples

The two Monte-Carlo models PYTHIA [58] and HERWIG [64] are used in this analysis. The generated events were put through the H1 detector simulation.

For PYTHIA, version 6.1 [63] was used in the *ep* interactions mode. It includes leading-order hard interactions, leading-logarithmic parton showers, parton distributions, initial and final state radiation, the Lund string fragmentation model (implemented in JETSET), multiple interactions and decay of the formed instable hadrons.

HERWIG (Hadron Emission Reactions With Interfering Gluons) version 5.1 [65] has been used. HERWIG has a detailed simulation of the QCD parton showers using a coherent showering algorithm applied to initial and final state radiation. The main difference to PYTHIA lies in the evolution of the parton showers, the hadronisation model used, and the treatment of multiple interactions (see also sections 3.1.1 and. 3.1.2).

The GRV (LO) [61, 62] photon and proton structure functions were used for both event generators. The minimum P_T of the hard scatter was set to 3 GeV and the maximum Q^2 to 4 GeV. The intrinsic k_T of the partons in the proton was set to 1 GeV for the PYTHIA Monte-Carlo files.

Prompt photon processes

Figure 3.3 shows the processes generated for the prompt photon events. No fragmentation functions of partons-to-photons (see section 2.1) were used. However contributions from di-jet events where a quark radiates a high E_T photon are also included in the PYTHIA samples.

Samples of PYTHIA events without multiple interaction were also simulated for comparison.

The amounts of simulated prompt photon events with PYTHIA and HERWIG are summarized in tables 3.1 and 3.2.



Figure 3.3: *Example of graphs for prompt photon production as handled by PYTHIA and HERWIG: direct-direct (a) resolved-direct (b) (see section 2.1), and in addition the smaller contribution from di-jet graphs where a quark radiates a photon are included (c),(d) for PYTHIA only.*

	Direct	Resolved	Direct radiative	Resolved Radiative
96	54.	70.	53.	56.
97	135	140.	127.	134.
98-99 e-	104.	94.	100.	83.
99 e+	130.	128.	120.	112.
2000 e+	312.	279.	299.	249.

Table 3.1: *Integrated luminosities in pb^{-1} simulated with PYTHIA for the conditions of the different years of data taking for graphs contributing to the prompt photon signal.*

	Direct	Resolved
96	57.	60.
97	132	137.
98-99 e-	102.	100.
99 e+	123.	133.
2000 e+	308.	303.

Table 3.2: *Integrated luminosities in pb^{-1} simulated with HERWIG for the conditions of the different years of data taking for graphs contributing to the prompt photon signal.*

Photo-production background processes

Di-jet events were also simulated to study the background from high P_T neutral mesons which may fake a photon signal in the detector due to the decay into photons according to:
$\pi^0 \to \gamma\gamma (B.R. = 99\%)$, $\eta \to \gamma\gamma (B.R. = 39\%)$, $\eta^0 \to \pi^0\pi^0\pi^0 (B.R. = 32\%)$.
At high energies the granularity of the detector does not allow a separation of the produced photons, and a single cluster is observed in the detector faking a single photon.

Background calculations were performed using PYTHIA, but they have little influence on the final prompt photon results (see chapter 7). They are only used for comparisons and for the estimate of contamination due to η's in proportion to π^0's.

Due to the very large cross section of di-jets in photo-production, preselection cuts were applied for this background simulation already at the generator level. Events without a high P_T neutral particle (fake prompt photon candidate) fulfilling some isolation requirements are rejected. The isolation requirement is performed around the candidate in cones of radii R in the plane of pseudo-rapidity η and azimuth ϕ.

The following quantities are defined:

$E_T^{cand} = transverse\ energy\ of\ the\ fake\ prompt\ photon\ candidate$
$E_{Tall}(R) = \sum_{r_i<R} E_T(all\ particles\ i)$
$E_{Tneutrals}(R) = \sum_{r_i<R} E_T(neutral\ particles\ i)$
$d_{charged} = distance\ in\ (\eta,\phi)\ plane\ of\ the\ closest\ charged\ particle\ to\ the\ candidate$

The following estimator defining the isolation of the neutral particle is used:

$For\ d_{charged} > 0.7:\ estimator = E_{Tneutrals}(0.1)/E_{Tall}(0.5)$
$For\ d_{charged} < 0.7:\ estimator = (D_{charged}/0.7)(E_{Tneutrals}(0.1)/ET_{all}(0.5) - 0.3)$

The events are accepted if:

$E_T^{cand} > 15$ GeV
or $E_T^{cand} > 5$ GeV and $estimator > 0.2$
or $4 > E_T^{cand} > 5$ GeV and $estimator > 0.3$

PYTHIA Monte-Carlo di-jet samples of smaller integrated luminosity (132 pb^{-1}) without preselection at the generator level are also used to control the above selection. An inefficiency of less than 10% is obtained.

The background calculations showed that after the experimental cuts (see chapter 6) the background is composed of π_0's (94%) and η's (5%). The other sources of background are negligible (below 2% in total e.g. 0.4 % antineutrons).

	Direct	Resolved
96	52.	42.
97	126.	121.
98-99 e-	100.	98.
99 e+	120.	114.
2000 e+	299.	338.

Table 3.3: *Integrated luminosities in pb^{-1} simulated with PYTHIA for the conditions of the different years of data taking for graphs contributing to the hadronic jet background.*

Deep inelastic scattering background processes

To study the background contribution from DIS, events corresponding to 65 pb^{-1} of events were generated using the RAPGAP [66] event generator.

Single particles

Single particles (γ's, π^0's, η^0's), in the liquid Argon calorimeter, with $4 < E_T < 15$ GeV, were simulated. These are used in the prompt photon extraction procedure based on a shower shape analysis (see chapter 7).

Chapter 4

The H1 experiment at HERA

The H1 experiment is located in Hamburg at DESY (the Deutsches Elektronen Synchrotron laboratory) and is one of the two collider experiments at HERA (the Hadron Electron Ring Accelerator). Since 1992 it has collected data on electron-proton and positron-proton scattering over a wide kinematic range. In this chapter the HERA accelerator is shortly presented and the H1 detector [67, 68] is briefly described, in particular the components most relevant for this analysis.

4.1 The HERA accelerator



Figure 4.1: *The storage rings of HERA and the pre-accelerators.*

HERA is composed of two underground storage rings of 6.3 km circumference, one for electrons or positrons, the other for protons. Figure 4.1 shows an overview of the HERA accelerator and the pre-accelerators LINAC, DESY II/III and PETRA. The electrons are accelerated to 27.5 GeV and the protons to 920 GeV (820 GeV before 1998). The center-of-mass energy of 320 GeV is one order of magnitude higher that reached at the previous electron-nucleon fixed target experiments. The other colliding experiment ZEUS is located diametrically opposite to H1. There are in addition two other experiments HERMES and HERA-B making use of only one beam (fixed target experiments) . HERMES studies the collisions of polarized electrons on polarized nucleons of a gaseous target and HERA-B uses proton-nucleon scattering mainly to study B- physics.

Each beam is composed of a maximum of 210 particle bunches separated by 96 nanoseconds. The colliding proton and electron bunches are synchronized such that they collide within the H1 and ZEUS detectors. Pilot bunches are electron or proton bunches without corresponding partners in the opposite beam. Typically 10 pilot bunches are used to study the rate of interactions of the beams with the residual gas in the beam-pipe (beam gas interactions) or with the beam-pipe itself (beam wall interactions). The distribution of the interaction point or vertex is determined by the bunch lengths and is approximately Gaussian with a width of about 11 cm.

4.2 The H1 detector

The H1 detector is composed of many detector elements with different purposes situated around the interaction point of the colliding beams. It offers an almost 4π solid angle coverage, leaving out the regions of the entering and outgoing beams. The detector shows a backward forward asymmetry reflecting the different beam energies. A right handed set of Cartesian coordinates (x, y, z) is defined. The positive z direction is given by the direction of the proton beam. The x axis is directed toward the center of the HERA rings and the y axis points vertically upward. The polar angle θ is the angle with respect to the z axis so that $\theta = 180°$ for an unscattered electron (see Fig. 4.2)



Figure 4.2: *The definition of the Cartesian coordinate system (x, y, z) with respect to the e and p beam directions.The corresponding spherical coordinate system (r, θ, ϕ) is also shown.*