

Charm production in DIS and diffraction

Wolfram Erdmann, ETH-Zürich


Photon 2001

representing **H1** and **ZEUS**

- Introduction
- charm production in DIS
 - $D^{*\pm}$ in e^+p , double differential (H1)
 - $D^{*\pm}$ in e^+p, e^-p (Zeus)
 - semileptonic charm (Zeus)
 - $F_2^c(x, Q^2)$
- $D^{*\pm}$ in diffraction
 - photoproduction
 - DIS
- summary

charm production in ep scattering

dominant at HERA: Boson Gluon Fusion



- $Q^2 = -q^2$
Momentum transfer
DIS: $Q^2 > 1\text{GeV}^2$
- $x_B = \frac{Q^2}{2P \cdot q}$
Bjorken scaling variable
- $y = \frac{P \cdot q}{P \cdot k}$
inelasticity

$$q = p(e') - p(e), \quad P = p(p)$$

BGF probes directly the gluon content of the proton (or IP).

approaches in pQCD:

- NLO + DGLAP
- LO + CCFM

predictions depend on


- structure functions
- value of the charm quark mass m_c
- hadronization

charm tagging I

Work horse:

$$D^{*+} \rightarrow D^0 \pi_s^+ \quad \mathcal{B} = 67.7\%$$

$$\hookrightarrow K^- \pi^+ \pi_s^+ \quad \mathcal{B} = 3.38\%$$



$$|M(K^- \pi^+) - M(D^0)| < 70 \text{ MeV}$$

$$N = 973 \pm 40$$

- $f(c \rightarrow D^{*+}) = 0.233 \pm 0.010 \pm 0.011$ (Aleph)
(changed by 16% wrt previous H1 publication)

$D^{*\pm}$ in DIS: total cross section

H1 1996 – 1997 $e^+p, \mathcal{L} = 18.6\text{pb}^{-1}$ (hep-ex/0108039, subm. to Phys. Lett. B)

Kinematic region

$$1 < Q^2 < 100\text{GeV}^2 \quad (\text{new backward ECAL})$$

$$0.05 < y < 0.7$$

$$E(e') > 8\text{GeV}$$

$$\text{visible: } p_t(D^*) > 1.5\text{GeV} \quad \text{and} \quad |\eta(D^*)| < 1.5$$

$$\sigma_{vis}(e^+p \rightarrow e^+D^{*\pm}X) = 8.50 \pm 0.42(\text{stat.})_{-0.76}^{+1.02}(\text{syst.}) \pm 0.65(\text{mod.})\text{nb}$$

syst.: tracking efficiency, signal extraction

QCD

NLO DGLAP $5.17\text{nb} - 7.02\text{nb}$ HVQDIS, GRV98

CCFM $8.04\text{nb} - 10.77\text{nb}$ CASCADE, H1 Fit

Ranges: $(m_c, \epsilon_c) = (1.3\text{GeV}, 0.100) - (1.5\text{GeV}, 0.035)$

good agreement with published Zeus result.

(Eur. Phys. J. C12, 35-52(2000)).

similar kinematic region:





$$1 < Q^2 < 600\text{GeV}^2$$

$$0.02 < y < 0.7$$

$$\text{visible: } 1.5\text{GeV} < p_t(D^*) < 15\text{GeV} \quad \text{and} \quad |\eta(D^*)| < 1.5$$


$$\sigma_{vis}(e^+p \rightarrow e^+D^{*\pm}X) = 8.31 \pm 0.31(\text{stat.})_{-0.50}^{+0.30}(\text{syst})\text{nb}$$

$D^{*\pm}$ in DIS: differential cross section



- HVQDIS = NLO+DGLAP, CASCADE = LO+CCFM
- bands: $(m_c, \epsilon_c) = (1.3\text{GeV}, 0.035) \rightarrow (1.5\text{GeV}, 0.100)$
- shapes in $pt(D^*), Q^2, x$ well described

double differential cross sections



- $\eta > 0$ deviation not confined to a certain Q^2 region (no breakdown of “massive” approach for high Q^2)
- most prominent at very low transverse momenta

$D^{*\pm}$ in e^+p and e^-p

- 1998: Hera beam energy 820GeV \rightarrow 920GeV
expected cross section increase 5% (HVQDIS).
- 1998-1999: e^-p running ($\mathcal{L}_{\text{Zeus}} = 16.7\text{pb}^{-1}$)
- 1999-2000: e^+p running ($\mathcal{L}_{\text{Zeus}} = 65.2\text{pb}^{-1}$)

larger Q^2 range

$$1 < Q^2 < 1000\text{GeV}^2$$

$$0.02 < y < 0.8$$

Zeus preliminary:

$$\sigma_{vis}(e^-p \rightarrow e^+ D^{*\pm} X) = 10.20 \pm 0.48(\text{stat.})_{-0.54}^{+0.34}(\text{syst})\text{nb}$$
$$\sigma_{vis}(e^+p \rightarrow e^+ D^{*\pm} X) = 8.94 \pm 0.24(\text{stat.})_{-0.51}^{+0.27}(\text{syst})\text{nb}$$

- no lepton charge dependence expected from BGF
- no systematic effect identified
 - rates of random background 1.01 ± 0.05
 - no difference observed in photoproduction