

# Forward Jet Production in DIS



N. Vlasov

On behalf of the ZEUS Collaboration



ALBERT-LUDWIGS-  
UNIVERSITÄT FREIBURG

XIII International Workshop on  
Deep Inelastic Scattering


April 26 – May 1, 2005

Madison, Wisconsin, U.S.A.

## OUTLINE:

- QCD Dynamics at Low  $x$
- MC Modes and QCD Calculations
- Inclusive Forward Jet Measurement with ZEUS data 95-97
- Inclusive Forward Jet Measurement with ZEUS data 98-2000
- Conclusions

# QCD Dynamics at Low $x$




## DGLAP

Evolution & resummation  
in powers of  $\ln Q^2$

$$Q^2 \gg k_{T,n}^2 \gg \dots \gg k_{T,2}^2 \gg k_{T,1}^2$$

The DGLAP gluon cascade  
is strongly ordered in  $k_T$   
and ordered in  $x$



## BFKL

Evolution & resummation  
in powers of  $\ln(1/x)$

$$x_1 \gg x_2 \gg \dots \gg x_n \gg x$$

The BFKL is only  
strongly ordered in  $x$

- **DGLAP (Dokshitzer-Gribov-Lipatov-Altarelli-Parisi)** is expected to break down at low  $x$  and  $Q^2$  region
- **BFKL (Balitsky-Fadin-Kuraev-Lipatov)** can be applicable at low  $x$
- **CCFM (Ciafaloni-Catani-Fiorani-Marchesini)** describes an evolution in both  $Q^2$  and  $x$  and approaches BFKL at low  $x$  and DGLAP at high  $Q^2$ ; angular ordering

# MC Models and QCD Calculations

- **DISENT**: Fixed order QCD partonic cross section, on mass shell ME + DGLAP

- **LEPTO**: LO ME+PS , (DGLAP)



- Strong ordering in  $k_T$

- **ARIADNE**: LO, an implementation of Color Dipole Model (CDM)

- Independently radiating dipoles formed by quarks and emitted gluons
- Random walk in  $k_T$  like in BFKL

- **CASCADE**: LO off mass shell ME + parton shower based on  $k_T$  factorized CCFM evolution model

- Angular ordering in parton emission
- Unintegrated gluon densities fits *J2003 set 1* (with new treatment of soft region ) and *J2003 set 2* (fit form includes non singular terms)



# Inclusive Forward Jet Measurements

## DIS kinematical range

95-97 Data,  $L \cong 38 \text{ pb}^{-1}$

$$Q^2 > 25 \text{ GeV}^2$$

No restriction


$$y > 0.04$$

98-00 Data,  $L \cong 82 \text{ pb}^{-1}$

$$20 < Q^2 < 100 \text{ GeV}^2$$

$$0.0004 < x_{Bj} < 0.005$$

$$0.04 < y < 0.7$$



$$x_{jet} = k_{z,jet}/p \gg x_{Bj}$$

$$k_{T,jet}^2 \sim Q^2$$

## Forward Jet selection

Jet finding with inclusive  $K_T$  algorithm in Lab frame

$$E_{T,jet} > 6 \text{ GeV}$$

No restriction

$$0.5 < E_{T,jet}^2/Q^2 < 2$$

$$0 < \eta^{jet} < 3$$

$\cos\gamma_{had} < 0$  suppresses QPM

Jet finding with inclusive  $K_T$  algorithm in Breit frame

$$E_{T,jet} > 5 \text{ GeV}$$

$$x_{jet} > 0.036$$

$$0.5 < E_{T,jet}^2/Q^2 < 2$$


$$2 < \eta^{jet} < 3.5$$

- Jet carries a large fraction of longitudinal momentum of proton in order to maximise phase space for BFKL evolution

- DGLAP type of evolution is suppressed, leaving no room for strong ordering in transverse momenta

# Inclusive Forward Jet Measurement (DESY-05-017)


## ZEUS



- NLO gives a good description of  $E_T^{\text{jet}}$  dependence
- Discrepancy between data and NLO in the forward region  $1.5 < \eta^{\text{jet}} < 3$ ; this region is more sensitive to higher order radiations (estimation of uncertainty from higher orders is large)


# Inclusive Forward Jet Measurement (DESY-05-017)

## ZEUS



- NLO predictions lower than data at low  $x_{Bj}$  but still within theoretical uncertainties. Gives a good description of  $Q^2$  dependence
- CDM describes all measured cross sections
- ME+PS: LEPTO (DGLAP) fails for low  $x_{Bj}$  and  $Q^2$


## Forward Jet Measurement with Forward Plug Calorimeter (FPC)



- Forward Plug Calorimeter in the  $20 \times 20 \text{ cm}^2$  beam hole of FCAL for 98-2000
- Lead-scintillator sandwich with 60 EMC and 16 HAC cells
- Extend calorimeter acceptance by about 1 unit in pseudorapidity to  $\eta \leq 5$

# Inclusive Forward Jet Measurement

## ZEUS




## Measurement extended to $2 < \eta^{\text{jet}} < 3.5$

- CASCADE *set1* disagrees with all cross sections
- CASCADE *set2* is in a good agreement with data in  $Q^2$  and  $E_T^{\text{jet}}$  but fails to reproduce the shapes of  $x_{Bj}$  and  $\eta^{\text{jet}}$
- CDM (ARIADNE) gives a good description of data in all measured cross sections
- LEPTO underestimates data by a factor of 2



# Inclusive Forward Jet Measurement

## ZEUS



- Average hadronisation correction obtained with LEPTO and ARIADNE
- Proton PDF CTEQ5D
- NLO predictions lower than data but within theoretical uncertainties (except very low  $x_{Bj}$ )
- Theory has too large uncertainty
- No disagreement with NLO DGLAP has been observed for forward jets

## Conclusions

- ❑ Parton dynamics at low- $x$  has been studied in forward jet production in DIS by ZEUS
- ❑ CDM (ARIADNE) gives a good description of data in all measured cross sections
- ❑ Lowest-order DGLAP calculations fail to describe forward jet cross sections
- ❑ LO CCFM-based CASCADE does not describe shapes of forward jet cross sections
- ❑ NLO calculations fail to describe data at low Bjorken- $x$  but the agreement is good for very forward jets within theoretical uncertainties
- ❑ Large renormalisation uncertainty does not permit more detailed comparison