



Particle Spectra at ZEUS

Lydia Shcheglova

on behalf of the ZEUS Collaboration
Nuclear Physics Institute
Moscow State University

EPS 2009, Krakow
16-22 July 2009

Outline

- **Motivation**
- **Multiplicity studies in NC DIS:**
 - multiplicity distributions
 - KNO scaling
 - energy dependence of multiplicity with different energy scales
- **Scaled momentum distributions in dijet photoproduction:**
 - comparison with the MLLA predictions
 - Λ_{eff} , LPHD k_{ch}

Multiplicity Distributions in DIS

Data and motivation

- Luminosity 38.6 pb⁻¹ collection in 1996-7 with $E_{\text{proton}} = 820 \text{ GeV}$ and $E_{e^+} = 27.5 \text{ GeV}$
- NC DIS events with $Q^2 > 25 \text{ GeV}^2$
 $70 < W < 225 \text{ GeV}$


Comparison with e^+e^- in previous studies in Breit frame:

- a reasonable agreement at $Q > 8 \text{ GeV}$
- no agreement at $Q < 8 \text{ GeV}$
- explained by the asymmetric nature of γ^*p


Alternative energy scales to Q :

- the invariant mass of hadronic system $M_{\text{eff}}^{\text{Breit}}$ and $M_{\text{eff}}^{\text{HCM}}$
- the available energy in the current region of Breit frame E_B^{Cr} or of HCM $E_{\text{HCM}}^{\text{Cr}} \approx W/2$


$$M_{\text{eff}}^2 = \left(\sum_{i \neq e'} E^i \right)^2 - \left(\sum_{i \neq e'} p_x^i \right)^2 - \left(\sum_{i \neq e'} p_y^i \right)^2 - \left(\sum_{i \neq e'} p_z^i \right)^2$$




ZEUS




ZEUS



ZEUS



ZEUS



Multiplicity in

$2 E_B^{cr}$ bins in the Breit frame

W bins in HCM frame

M_{eff} bins in the Breit frame

M_{eff} bins in the CR of HCM frame

ARIADNE is the best


HERWIG
—longer tails for multiplicities

→ increase of the systematic uncertainties

KNO SCALING

W and 2 E_B^{cr} bins


ZEUS



Koba, Z.H.B.Nielson, P.Olsen
N.P.B40(1972)317

M_{eff} bins



ZEUS



$$\langle n \rangle P(n_{ch}) = \psi(n_{ch} / \langle n_{ch} \rangle)$$

Scaling behaviour observed for HCM and Breit except M_{eff} less than 4 GeV

KNO scaling comparison with e+e-



a) ep in bins
2 E_B^{cr}
W
2E_B^{cr} > 12 GeV

b) in bins
M_{eff}
M_{eff} > 8 GeV

e+e- data from **one hemisphere**


LEP (DELPHI, OPAL) at Z⁰
PETRA (TASSO)

e+e- data for **2 hemispheres**

LEP (DELPHI, OPAL, L3, ALEPH)
PETRA (TASSO)

Energy dependence of average multiplicity

ZEUS



scales used:

$$2 E_B^{cr}$$


$$W$$

$$M_{eff}$$

At low energy scales :
differences if $2E_B^{cr}$ or Q
used

M_{eff} energy scale

ZEUS




All three models describe the data reasonably well, in the last M_{eff} bin in C.R. HCM the Herwig prediction is too high

$\langle n_{\text{ch}} \rangle$ vs M_{eff} in the C.R. of the Breit frame shows the same behaviour as $2\langle n_{\text{ch}} \rangle$ vs $2E_{\text{B}}^{\text{cr}}$
 $\langle n_{\text{ch}} \rangle$ vs M_{eff} rises faster in HCM than in the Breit frame

Summary plot

ZEUS



The measurements show good overall agreement with the those from other experiments. Exhibit approximately the same dependence of the $\langle n_{ch} \rangle$ on the respective energy scale.

- At low value of energy $\langle n_{ch} \rangle$ vs $2 E_B^{cr}$ agrees well with e+e- in contrast the previous measurements vs Q.
- data in C.R. of HCM (W) agree with the LEP data, but systematically below them
- when using these scales, ep DIS data can be consistently compared with data from e+e-, μP and νP collisions over a wide energy region

Scaled momentum distributions of charged particles in dijet photoproduction

Data and motivation

Luminosity 359 pb^{-1} collected in 2005-7
with $E_{\text{proton}} = 920 \text{ GeV}$ and $E_e = 27.5 \text{ GeV}$

Photoproduction events (γp) were studied :

- required to have only two reconstructed jets
- energy scales in the range **19 to 38 GeV**
- cones of various opening angles θ_c around the jet axis
- jets were reconstructed using the k_T cluster algorithm

Tests of MLLA predictions

- comparison of the scaled momentum distribution in jets with MLLA , LPHD is assumed
- extract MLLA scale Λ_{eff} and the LPHD parameter κ_{ch}
- Λ_{eff} previously measured for ee , eP , $P\bar{P}$, never for γP
- is Λ_{eff} independent of interaction type? ee , ep , $P\bar{P}$, γP
- is Λ_{eff} independent of E_{jet} and θ_c as predicted?

The MLLA+LPHD Theoretical Framework

The Modified Leading Log Approximation

- All orders pQCD resummation
- Analytical description of parton evolution
- Predicts parton multiplicity and momenta
- MLLA describes fragmentation with 2 parameters:
 - Q_0 - self-imposed cut-off energy scale
 - Λ_{eff} - QCD scale effective parameter
- We study MLLA within jets, where fragmentation is well defined
- Λ_{eff} - predicted to be universal
- Assuming Local Parton Hadron Duality MLLA predictions are directly comparable to the data

The Local Parton Hadron Duality (LPHD)

- Assumes hadronization is local and occurs at the end of parton shower
- Relates the observed hadron distributions to calculated parton distributions via constant factor, κ_{ch}
- κ_{ch} is the ratio of the number of charged particles over the total number of partons produced during fragmentation
- 2 free parameters in MLLA + LPHD: Λ_{eff} and κ_{ch}

Measurement of Λ_{eff} and κ_{ch}


Strategy of the analysis

- Measure scaled track momentum within jets, $x_p = \frac{P_{\text{track}}}{P_{\text{Jet}}}$.
- Plot scaled momentum distributions, $\xi = \ln\left(\frac{1}{x_p}\right)$, in bins of jet energy $E_{\text{Jet}} = \frac{M_{2j}}{2}$ (the hard scale) and θ_c (the opening angle).

To check the validity of the MLLA predictions fitting of the measured ξ distributions was done by

- 2 methods:
 - Gaussian around mean
 - MLLA +LPHD theory
- ξ_{peak} , Λ_{eff} and κ_{ch} were extracted from fits

The measured ξ distributions



- The Gaussian fits are shown. $0.48 \leq \chi^2/\text{dof} \leq 1.33$
- Blue lines indicate fitted region, ± 1 around mean.

The Gaussian fit method

- Gives peak position of ξ distribution, ξ_{peak}
- ξ_{peak} gives Λ_{eff} - Only valid for Leading Order


The MLLA +LPHD fit method

- Gives Λ_{eff} and K (the normalization) directly from fit
- κ_{ch} is calculated from K
- Λ_{eff} has strong dependence on ambiguous fit range
- κ_{ch} only weakly dependant on the fitting range

The results presented here use

- The Gaussian method for ξ_{peak} and Λ_{eff}
- The MLLA+LPHD method for κ_{ch} and to cross check Λ_{eff}

The Gaussian fit method



Peak position, ξ_{peak}


- Fit Gaussian ± 1 around mean.
- $\forall \xi$, independently measure ξ_{peak} .

$$\Lambda_{\text{eff}} = \frac{E_{\text{jet}} \sin(\theta_c)}{e^{(\sqrt{0.87+2\xi_{\text{peak}}}-0.54)^2}} \quad (@ \text{ LO})$$

Measuring Λ_{eff}

- Only use $\theta_c = 0.23$ energy points:
 - Different θ_c values are correlated;
 - MLLA loses validity at large θ_c .
- Fit equation to all 5 energy points.

$$\Lambda_{\text{eff}} = 275 \pm 4 (\text{stat.})_{-8}^{+4} (\text{syst.}) \text{ MeV}$$



The MLLA+ LPHD fit method

Momentum distribution of partons from a gluon is given by:

- $$\bar{D}_{g\text{-Jet}}^{\text{lim}} \left(\ln \left(\frac{1}{x_p} \right), Y \right) = \frac{4C_f}{b} \Gamma(B) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-B\alpha} \left[\frac{\cosh \alpha + (1-2\zeta) \sinh \alpha}{\frac{4N_c}{b} Y \frac{\alpha}{\sinh \alpha}} \right]^{\frac{B}{2}} \cdot I_B \left(\sqrt{\frac{16N_c}{b} Y \frac{\alpha}{\sinh \alpha} [\cosh \alpha + (1-2\zeta) \sinh \alpha]} \right) \frac{d\tau}{\pi}$$
- Valid for: $\ln \left(\frac{1}{x_p \ll 1} \right) \leq \ln \left(\frac{1}{x_p} \right) \leq \ln \left(\frac{M_{2j}}{2P_0} \right)$ $P_0 =$ Upper bound

For number of flavours, $N_f = 3$, and number of colours, $N_c = 3$

- $C_f = \frac{9}{4}$, $b = 9$, $B = 1.247$.
- I_B is the modified Bessel function of order B.
- $\alpha = \alpha_0 + i\tau$, where α_0 is determined by $\tanh \alpha_0 = 2\zeta - 1$
- $\zeta = 1 - \frac{\ln \left(\frac{1}{x_p} \right)}{Y}$ and $Y = \ln \left(\frac{E_{\text{Jet}} \sin(\theta_c)}{\Lambda_{\text{eff}}} \right)$
 $\bar{D}_{q\text{-Jet}}^{\text{lim}} = \frac{1}{r} \bar{D}_{g\text{-Jet}}^{\text{lim}}$

Quarks, gluons and the next-to-MLLA predictions

Quark and gluon jet mixture

- In γP events there is a mix of quark and gluon jets.

$\bar{D}_{\text{mix}}^{\text{lim}} = \left(\epsilon_g + \frac{1-\epsilon_g}{r} \right) \bar{D}_{\text{g-Jet}}^{\text{lim}}$, where ϵ_g is the fraction of gluon jets.


Energy (GeV)	19	23	28	32	38
ϵ_g (From PYTHIA)	0.203	0.213	0.211	0.227	0.242

The so called "next-to-MLLA" predictions

- Not actually higher order calculation, but a modification of MLLA.
- In nMLLA, $\bar{D}_{\text{mix}}^{\text{lim}} = F_{\text{nMLLA}} \left(\epsilon_g + \frac{1-\epsilon_g}{r} \right) \bar{D}_{\text{g-Jet}}^{\text{lim}}$
- Where $r = 1.6 \pm 0.2$ and $F_{\text{nMLLA}} = 1.3 \pm 0.2$ (from theory).
- When fitting to data the normalisation can be expressed as:


$$K = \kappa_{\text{ch}} F_{\text{nMLLA}} \left(\epsilon_g + \frac{1-\epsilon_g}{r} \right)$$

Λ_{eff} - Comparison of extraction methods



Λ_{eff} extracted from 359pb¹ ZEUS data via both methods

- $\forall \xi$, independently extract Λ_{eff} : **Red = Gaussian**. Black = nMLLA.
- Λ_{eff} has a weak dependence on θ_c , no dependence on scale.
- nMLLA, $\theta_c = 0.23$: $\Lambda_{\text{eff}} = 304 \pm 6 \text{ (stat.)}_{-32}^{+8} \text{ (syst.) MeV}$
- Large nMLLA systematics come from ambiguous fitting range.
- nMLLA regularisation scheme \Rightarrow Parton cut-off at $p_T^{\text{rel,pl}} = \Lambda_{\text{eff}}$




κ_{ch} extracted from 359pb¹ ZEUS data via nMLLA method

- κ_{ch} comes from the normalisation of ξ
- κ_{ch} is insensitive to the ambiguous fitting range.
- κ_{ch} has a weak dependence on θ_c , no dependence on scale.
- Theoretical uncertainties dominate the overall uncertainty.

$$\kappa_{\text{ch}} = 0.55 \pm 0.01 \text{ (stat.)}_{-0.02}^{+0.03} \text{ (syst.)}_{-0.09}^{+0.11} \text{ (theo.)}$$


Global Comparisons - ξ_{peak}



Global fit gives $\Lambda_{\text{eff}} = 246 \pm 3 \text{ MeV}$ with $\chi^2/\text{dof} = 2.20$

- The fit assumes that Λ_{eff} is independent of scale and θ_c .
- Both ZEUS and CDF observe a weak θ_c dependence.
- CDF also observe a weak scale dependence:
 - Λ_{eff} observed to decrease with increasing energy.
- May explain why this is inconsistent with ZEUS only fit result.

Global Comparisons - Λ_{eff}



Λ_{eff} as a function of energy scale for different experiments

- 359pb^{-1} ZEUS data fills the gap from $19 \rightarrow 38$ GeV.
- First measurement of Λ_{eff} from γp process.

Summary

Summary

- Scaled momentum distributions have been measured in dijet events in 359pb^{-1} γp ZEUS data.
- Λ_{eff} and κ_{ch} have been extracted at energy scales from $19 \rightarrow 38$ GeV.

$$\Lambda_{\text{eff}} = 275 \pm 4 \text{ (stat.)}_{-8}^{+4} \text{ (syst.) MeV}$$

$$\kappa_{\text{ch}} = 0.55 \pm 0.01 \text{ (stat.)}_{-0.02}^{+0.03} \text{ (syst.)}_{-0.09}^{+0.11} \text{ (theo.)}$$

Publication

- Pre-print on arXiv : [hep-ex/0904.3466](https://arxiv.org/abs/hep-ex/0904.3466)

Conclusions

Hadronic spectra proved to be a powerful tool for study of various aspects of the multiparticle dynamics:

Multiplicity studies in NC DIS:

- detailed comparison multiplicity distributions with MC models
- KNO scaling was studied with different energy scales
- energy dependence of multiplicity was investigated for different energy scales and detailed comparison with e+e- was done

Scaled momentum distribution:

- scaled momentum distributions were measured
- Λ_{eff} and LPHD and κ_{ch} were extracted

Still rich program of studies with ZEUS : many results not shown, many studies in progress...

Thank you for your attention!

**Many thanks to Krakow
for
hospitality!**